



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2020 – 05:33 PM BST

PDB ID : 4MW5
Title : Trypanosoma brucei methionyl-tRNA synthetase in complex with inhibitor 1-{3-[(3-chloro-5-methoxybenzyl)amino]propyl}-3-phenylurea (Chem 1415)
Authors : Koh, C.Y.; Kim, J.E.; Wetzel, A.B.; de van der Schueren, W.J.; Shibata, S.; Liu, J.; Zhang, Z.; Fan, E.; Verlinde, C.L.M.J.; Hol, W.G.J.
Deposited on : 2013-09-24
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.14.4.dev1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	<div> <div>6%</div> <div> <div></div> <div>93%</div> <div>5%</div> </div> </div>
1	B	542	<div> <div>4%</div> <div> <div></div> <div>93%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9143 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionyl-tRNA synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	527	Total	As	C	N	O	S	0	3	0
			4243	1	2732	719	779	12			
1	B	530	Total	As	C	N	O	S	0	2	0
			4254	1	2740	722	780	11			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP Q38C91
A	-3	PRO	-	EXPRESSION TAG	UNP Q38C91
A	-2	GLY	-	EXPRESSION TAG	UNP Q38C91
A	-1	SER	-	EXPRESSION TAG	UNP Q38C91
A	0	MET	-	EXPRESSION TAG	UNP Q38C91
A	309	THR	ALA	CONFLICT	UNP Q38C91
A	452	ALA	LYS	ENGINEERED MUTATION	UNP Q38C91
A	453	ARG	LYS	ENGINEERED MUTATION	UNP Q38C91
A	454	ALA	GLU	ENGINEERED MUTATION	UNP Q38C91
A	499	VAL	ALA	CONFLICT	UNP Q38C91
A	503	ASN	SER	CONFLICT	UNP Q38C91
B	-4	GLY	-	EXPRESSION TAG	UNP Q38C91
B	-3	PRO	-	EXPRESSION TAG	UNP Q38C91
B	-2	GLY	-	EXPRESSION TAG	UNP Q38C91
B	-1	SER	-	EXPRESSION TAG	UNP Q38C91
B	0	MET	-	EXPRESSION TAG	UNP Q38C91
B	309	THR	ALA	CONFLICT	UNP Q38C91
B	452	ALA	LYS	ENGINEERED MUTATION	UNP Q38C91
B	453	ARG	LYS	ENGINEERED MUTATION	UNP Q38C91
B	454	ALA	GLU	ENGINEERED MUTATION	UNP Q38C91
B	499	VAL	ALA	CONFLICT	UNP Q38C91
B	503	ASN	SER	CONFLICT	UNP Q38C91

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is METHIONINE (three-letter code: MET) (formula: $C_5H_{11}NO_2S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			9	5	1	2	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 1-{3-[(3-chloro-5-methoxybenzyl)amino]propyl}-3-phenylurea (three-letter code: 415) (formula: C₁₈H₂₂ClN₃O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	Cl	N	O	0	0
			24	18	1	3	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	275	Total	O	0	0
			275	275		
7	B	269	Total	O	0	0
			269	269		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methionyl-tRNA synthetase



- Molecule 1: Methionyl-tRNA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.45Å 105.73Å 206.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.35 48.16 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.6 (30.00-2.35) 99.6 (48.16-2.35)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.34Å)	Xtriage
Refinement program	REFMAC refmac_5.7.0032	Depositor
R, R_{free}	0.189 , 0.217 0.179 , 0.205	Depositor DCC
R_{free} test set	4054 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.541	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9143	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, GOL, DMS, SO4, 415

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/4346	0.60	1/5905 (0.0%)
1	B	0.43	0/4354	0.61	0/5914
All	All	0.42	0/8700	0.61	1/11819 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	476	ASP	CB-CG-OD1	5.68	123.41	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4243	0	4185	18	0
1	B	4254	0	4209	17	0
2	A	18	0	24	2	0
2	B	30	0	40	1	0
3	A	8	0	12	0	0
3	B	8	0	12	0	0
4	A	9	0	8	0	0
5	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	24	0	22	4	0
7	A	275	0	0	0	0
7	B	269	0	0	0	0
All	All	9143	0	8512	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:487:LEU:HD22	1:B:495:GLU:HG3	1.63	0.81
1:A:487:LEU:HD22	1:A:495:GLU:HG3	1.67	0.76
1:A:548:TRP:CH2	1:A:581:LEU:HD22	2.22	0.74
1:B:474:TRP:HE3	6:B:909:415:H1	1.59	0.66
1:A:271:ARG:NH1	2:A:902:GOL:O2	2.27	0.66
1:A:548:TRP:CZ3	1:A:581:LEU:HD22	2.35	0.62
1:A:668:VAL:HG11	1:A:713:LEU:HG	1.85	0.58
1:B:462:PRO:HA	1:B:470:CAS:CE1	2.33	0.58
1:A:487:LEU:HD22	1:A:495:GLU:CG	2.35	0.57
1:A:665:ILE:HG12	1:A:716:ILE:HD13	1.89	0.55
1:B:361:GLY:HA2	1:B:399:TRP:CZ3	2.42	0.55
1:A:322:MET:HE3	1:A:566:VAL:HG22	1.87	0.55
1:B:474:TRP:CE3	6:B:909:415:H1	2.41	0.54
1:B:487:LEU:HD22	1:B:495:GLU:CG	2.36	0.53
1:B:462:PRO:CA	1:B:470:CAS:CE1	2.87	0.53
1:A:272:TYR:CE2	1:A:276:LYS:HD2	2.46	0.51
1:B:284:THR:HG22	1:B:326:ILE:HG21	1.93	0.50
1:B:478:LEU:HG	6:B:909:415:H2	1.92	0.50
1:B:668:VAL:HG11	1:B:713:LEU:HG	1.94	0.48
1:A:284:THR:HG22	1:A:326:ILE:HG21	1.96	0.47
1:A:583:ARG:HH22	2:A:901:GOL:H2	1.79	0.47
1:A:682:MET:O	1:B:434[A]:ARG:NH2	2.46	0.46
1:A:665:ILE:CG1	1:A:716:ILE:HD13	2.46	0.45
1:A:580:PHE:CE1	1:A:598[B]:MET:HA	2.53	0.44
1:B:456:LEU:HD12	1:B:470:CAS:HB3	2.00	0.43
1:B:427:ILE:HB	1:B:433:ARG:HG3	2.00	0.43
1:B:478:LEU:HG	6:B:909:415:CAA	2.49	0.43
1:A:580:PHE:CE1	1:A:598[A]:MET:HA	2.53	0.43
1:B:648:LEU:HB3	1:B:649:PRO:HD3	2.00	0.42
1:A:714:SER:N	1:A:715:PRO:CD	2.82	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:714:SER:N	1:B:715:PRO:CD	2.82	0.42
1:B:482:LEU:HD21	2:B:901:GOL:H32	2.03	0.41
1:A:687:LEU:HA	1:A:690:THR:HG22	2.02	0.40
1:A:261:TYR:CD1	1:A:548:TRP:CH2	3.09	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/542 (97%)	513 (98%)	12 (2%)	0	100	100
1	B	527/542 (97%)	515 (98%)	12 (2%)	0	100	100
All	All	1052/1084 (97%)	1028 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/468 (97%)	454 (100%)	1 (0%)	93	96
1	B	456/468 (97%)	450 (99%)	6 (1%)	69	79
All	All	911/936 (97%)	904 (99%)	7 (1%)	81	89

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	690	THR
1	B	377	THR
1	B	398	THR
1	B	597	ASN
1	B	681	ASP
1	B	760	VAL
1	B	767	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	B	470	1	5,8,9	0.89	0	1,9,11	0.43	0
1	CAS	A	470	1	5,8,9	0.93	0	1,9,11	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	B	470	1	-	0/0/7/9	-
1	CAS	A	470	1	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	470	CAS	3	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DMS	A	904	-	3,3,3	0.57	0	3,3,3	0.84	0
2	GOL	B	904	-	5,5,5	0.19	0	5,5,5	0.43	0
2	GOL	B	903	-	5,5,5	0.17	0	5,5,5	0.43	0
3	DMS	B	907	-	3,3,3	0.49	0	3,3,3	0.75	0
2	GOL	A	901	-	5,5,5	0.24	0	5,5,5	0.68	0
4	MET	A	906	-	4,8,8	0.39	0	2,9,9	0.60	0
2	GOL	A	902	-	5,5,5	0.30	0	5,5,5	0.66	0
5	SO4	B	908	-	4,4,4	0.40	0	6,6,6	0.24	0
3	DMS	A	905	-	3,3,3	0.49	0	3,3,3	0.53	0
2	GOL	B	902	-	5,5,5	0.20	0	5,5,5	0.55	0
2	GOL	B	901	-	5,5,5	0.38	0	5,5,5	0.69	0
3	DMS	B	906	-	3,3,3	0.48	0	3,3,3	0.85	0
2	GOL	A	903	-	5,5,5	0.33	0	5,5,5	0.54	0
2	GOL	B	905	-	5,5,5	0.33	0	5,5,5	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	415	B	909	-	25,25,25	0.23	0	31,31,31	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MET	A	906	-	-	0/4/8/8	-
2	GOL	B	904	-	-	2/4/4/4	-
2	GOL	B	903	-	-	2/4/4/4	-
2	GOL	A	901	-	-	2/4/4/4	-
2	GOL	A	902	-	-	2/4/4/4	-
2	GOL	B	902	-	-	3/4/4/4	-
2	GOL	B	901	-	-	2/4/4/4	-
2	GOL	A	903	-	-	0/4/4/4	-
2	GOL	B	905	-	-	4/4/4/4	-
6	415	B	909	-	-	0/15/15/15	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	903	GOL	C1-C2-C3-O3
2	B	903	GOL	O2-C2-C3-O3
2	B	905	GOL	O1-C1-C2-C3
2	B	905	GOL	C1-C2-C3-O3
2	B	905	GOL	O1-C1-C2-O2
2	B	904	GOL	O1-C1-C2-C3
2	A	902	GOL	O1-C1-C2-C3
2	B	902	GOL	C1-C2-C3-O3
2	B	901	GOL	O1-C1-C2-C3
2	A	902	GOL	O1-C1-C2-O2
2	B	905	GOL	O2-C2-C3-O3
2	B	901	GOL	O1-C1-C2-O2
2	B	902	GOL	O1-C1-C2-C3
2	A	901	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	904	GOL	O1-C1-C2-O2
2	B	902	GOL	O2-C2-C3-O3
2	A	901	GOL	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	GOL	1	0
2	A	902	GOL	1	0
2	B	901	GOL	1	0
6	B	909	415	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	526/542 (97%)	-0.08	31 (5%) 22 31	19, 31, 84, 116	0
1	B	529/542 (97%)	-0.12	23 (4%) 35 46	19, 31, 79, 125	0
All	All	1055/1084 (97%)	-0.10	54 (5%) 28 39	19, 31, 83, 125	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	548	TRP	7.0
1	A	552	ARG	6.4
1	B	755	ALA	6.2
1	B	380	VAL	6.0
1	B	758	GLY	6.0
1	A	756	VAL	5.9
1	A	767	THR	5.4
1	A	757	GLU	5.3
1	A	768	GLU	4.8
1	B	767	THR	4.8
1	A	551	ASP	4.4
1	A	766	SER	4.4
1	A	383	ASP	4.4
1	B	559	LEU	4.3
1	A	384	GLY	4.3
1	A	561	ASN	4.1
1	B	756	VAL	3.9
1	A	759	GLU	3.9
1	B	558	SER	3.7
1	A	560	GLY	3.6
1	A	758	GLY	3.5
1	B	377	THR	3.4
1	A	624	VAL	3.4
1	B	491	GLU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	555	ILE	3.2
1	A	547	TRP	3.1
1	A	685	TRP	3.1
1	A	382	LYS	3.1
1	A	562	VAL	3.1
1	A	621	LYS	3.1
1	A	760	VAL	3.0
1	B	757	GLU	2.9
1	A	553	LYS	2.9
1	A	592	ASP	2.7
1	A	299	GLN	2.7
1	B	395	HIS	2.7
1	B	391	LEU	2.7
1	A	559	LEU	2.6
1	B	394	GLY	2.6
1	A	751	ARG	2.6
1	A	590	ASP	2.5
1	B	389	VAL	2.5
1	B	624	VAL	2.5
1	B	392	GLU	2.4
1	B	751	ARG	2.4
1	B	557	LYS	2.3
1	B	381	ASP	2.3
1	B	759	GLU	2.2
1	A	755	ALA	2.2
1	B	397	VAL	2.2
1	A	421[A]	HIS	2.1
1	B	393	SER	2.1
1	A	301	GLN	2.1
1	B	688	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	A	470	9/10	0.95	0.14	27,30,36,38	3
1	CAS	B	470	9/10	0.97	0.12	27,29,35,36	3

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	903	6/6	0.80	0.20	52,55,56,57	0
3	DMS	A	904	4/4	0.81	0.23	74,79,80,86	0
2	GOL	B	903	6/6	0.84	0.27	43,51,53,56	0
3	DMS	B	907	4/4	0.86	0.17	77,77,77,82	0
2	GOL	B	904	6/6	0.86	0.20	47,48,50,50	0
2	GOL	B	901	6/6	0.88	0.20	44,52,53,57	0
3	DMS	B	906	4/4	0.89	0.27	62,73,77,82	0
3	DMS	A	905	4/4	0.91	0.14	84,86,87,90	0
5	SO4	B	908	5/5	0.92	0.25	64,71,75,75	0
2	GOL	A	902	6/6	0.92	0.19	47,48,50,51	0
2	GOL	B	902	6/6	0.92	0.21	47,48,50,55	0
2	GOL	B	905	6/6	0.92	0.26	56,59,63,65	0
2	GOL	A	901	6/6	0.93	0.14	49,51,52,52	0
6	415	B	909	24/24	0.96	0.12	22,30,39,42	0
4	MET	A	906	9/9	0.98	0.10	25,25,26,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.