



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 10, 2021 – 10:02 AM EST

PDB ID : 7MXH
Title : CD1c with antigen analogue 3
Authors : Cao, T.P.; Shahine, A.; Rossjohn, J.
Deposited on : 2021-05-19
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	<div> <div>7%</div> <div>89%</div> <div>9%</div> <div>.</div> </div>
2	B	108	<div> <div>5%</div> <div>84%</div> <div>11%</div> <div>5%</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1c/T-cell surface glycoprotein CD1b chimeric protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2233	1426	386	412	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	SER	GLN	conflict	UNP P29017
A	52	GLN	ASN	engineered mutation	UNP P29017
A	57	GLN	ASN	engineered mutation	UNP P29017
A	108	GLY	LYS	engineered mutation	UNP P29017
A	128	GLN	ASN	engineered mutation	UNP P29017
A	242	GLY	TRP	engineered mutation	UNP P29016
A	280	GLY	-	expression tag	UNP P29016
A	281	SER	-	expression tag	UNP P29016
A	282	LEU	-	expression tag	UNP P29016
A	283	VAL	-	expression tag	UNP P29016
A	284	PRO	-	expression tag	UNP P29016
A	285	ARG	-	expression tag	UNP P29016

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	103	Total	C	N	O	S	0	0	0
			851	542	144	162	3			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	ASP	-	expression tag	UNP P61769
B	0	ALA	-	expression tag	UNP P61769
B	101	GLY	-	expression tag	UNP P61769

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	102	SER	-	expression tag	UNP P61769
B	103	LEU	-	expression tag	UNP P61769
B	104	VAL	-	expression tag	UNP P61769
B	105	PRO	-	expression tag	UNP P61769
B	106	ARG	-	expression tag	UNP P61769

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 2,6-anhydro-1-deoxy-1,1-difluoro-1-[(R)-hydroxy{[(4S,8S,12S,16S,20S)-4,8,12,16,20-pentamethylheptacosyl]oxy}phosphoryl]-D-glycero-D-galacto-heptitol (three-letter code: ZQ7) (formula: $C_{39}H_{77}F_2O_8P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	O	P	0	0
			50	39	2	8	1		

- Molecule 5 is DODECANE (three-letter code: D12) (formula: $C_{12}H_{26}$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C	0	0
			12	12		

- Molecule 6 is (2S)-2,3-dihydroxypropyl hexadecanoate (three-letter code: ZP7) (formula: $C_{19}H_{38}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			23	19	4		

- Molecule 7 is MALONIC ACID (three-letter code: MLA) (formula: C₃H₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	3	4		
7	B	1	Total	C	O	0	0
			7	3	4		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	88	Total 88	O 88	0	0
8	B	33	Total 33	O 33	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-cell surface glycoprotein CD1c/T-cell surface glycoprotein CD1b chimeric protein



- Molecule 2: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.57Å 72.73Å 98.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.16 – 2.11 44.16 – 2.11	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.16-2.11) 100.0 (44.16-2.11)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874-000	Depositor
R, R_{free}	0.208 , 0.255 0.206 , 0.252	Depositor DCC
R_{free} test set	1187 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.5	Xtriage
Anisotropy	0.307	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3318	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZQ7, ZP7, NAG, MLA, D12

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2300	0.62	0/3126
2	B	0.47	0/874	0.71	0/1183
All	All	0.45	0/3174	0.65	0/4309

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2233	0	2119	17	0
2	B	851	0	818	8	0
3	A	14	0	13	0	0
4	A	50	0	0	0	0
5	A	12	0	26	0	0
6	A	23	0	0	0	0
7	A	7	0	2	0	0
7	B	7	0	2	0	0
8	A	88	0	0	0	0
8	B	33	0	0	0	0
All	All	3318	0	2980	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:PRO:HB3	1:A:212:PHE:HB3	1.75	0.68
2:B:16:ALA:HB1	2:B:98:ARG:HD3	1.87	0.56
1:A:138:GLY:O	1:A:142:GLN:HG3	2.08	0.53
2:B:16:ALA:CB	2:B:98:ARG:HD3	2.39	0.53
1:A:126:PHE:CE1	1:A:156:THR:HG23	2.44	0.52
1:A:11:HIS:CD2	1:A:113:PHE:HZ	2.29	0.51
1:A:45:GLY:O	1:A:71:ARG:NH1	2.44	0.50
1:A:34:GLU:HG3	1:A:244:TRP:CZ2	2.50	0.47
1:A:13:ILE:HG23	2:B:57:PHE:CE2	2.50	0.46
1:A:126:PHE:HE1	1:A:156:THR:HG23	1.80	0.46
1:A:7:HIS:HA	1:A:104:LEU:O	2.16	0.45
2:B:34:SER:HB3	2:B:63:PHE:CZ	2.52	0.45
2:B:55:LEU:HD23	2:B:65:LEU:HD21	1.98	0.45
1:A:190:ALA:HA	1:A:208:HIS:O	2.17	0.45
1:A:117:ALA:HB2	2:B:61:TRP:CE2	2.53	0.44
1:A:80:GLU:O	1:A:84:HIS:HD2	2.01	0.43
2:B:74:THR:HG22	2:B:76:LYS:H	1.84	0.43
1:A:86:SER:HB3	1:A:87:GLN:OE1	2.19	0.43
1:A:184:ARG:NE	1:A:184:ARG:HA	2.33	0.42
1:A:218:TRP:HB3	1:A:265:LYS:HB2	2.01	0.42
2:B:24:LEU:O	2:B:68:TYR:HA	2.19	0.42
1:A:11:HIS:ND1	1:A:99:LYS:HE3	2.34	0.41
1:A:89:TYR:HD1	1:A:89:TYR:HA	1.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/285 (98%)	267 (96%)	11 (4%)	1 (0%)	34	32
2	B	101/108 (94%)	99 (98%)	2 (2%)	0	100	100
All	All	380/393 (97%)	366 (96%)	13 (3%)	1 (0%)	41	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/242 (99%)	240 (100%)	0	100	100
2	B	96/100 (96%)	96 (100%)	0	100	100
All	All	336/342 (98%)	336 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	HIS
1	A	185	GLN
1	A	239	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	301	1	14,14,15	0.53	0	17,19,21	0.62	0
7	MLA	B	201	-	0,6,6	-	-	0,7,7	-	-
5	D12	A	303	-	11,11,11	0.27	0	10,10,10	0.41	0
7	MLA	A	305	-	0,6,6	-	-	0,7,7	-	-
6	ZP7	A	304	-	22,22,22	0.84	2 (9%)	23,23,23	0.89	1 (4%)
4	ZQ7	A	302	-	46,50,50	1.41	3 (6%)	55,66,66	1.55	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	301	1	-	0/6/23/26	0/1/1/1
7	MLA	B	201	-	-	0/0/4/4	-
5	D12	A	303	-	-	3/9/9/9	-
7	MLA	A	305	-	-	0/0/4/4	-
6	ZP7	A	304	-	-	12/22/22/22	-
4	ZQ7	A	302	-	-	26/52/76/76	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	ZQ7	P1-O5	6.42	1.67	1.57
4	A	302	ZQ7	P1-O8	-4.89	1.46	1.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	304	ZP7	O18-C16	2.58	1.40	1.33
4	A	302	ZQ7	P1-O7	-2.33	1.46	1.51
6	A	304	ZP7	O18-C19	-2.01	1.40	1.45

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	302	ZQ7	C29-O1-C31	4.98	122.60	113.19
4	A	302	ZQ7	O8-P1-O7	-4.22	101.17	111.34
6	A	304	ZP7	O18-C16-C15	3.02	121.37	111.91
4	A	302	ZQ7	C8-C7-C3	-2.56	107.64	115.92
4	A	302	ZQ7	O5-P1-O7	-2.49	111.07	115.48
4	A	302	ZQ7	C23-C32-C35	-2.38	105.74	113.27
4	A	302	ZQ7	O1-C31-C28	2.38	114.36	109.06
4	A	302	ZQ7	P1-O5-C35	-2.36	116.80	122.71
4	A	302	ZQ7	C13-C14-C15	-2.25	108.63	115.92

There are no chirality outliers.

All (41) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	302	ZQ7	C31-C1-P1-O5
4	A	302	ZQ7	F1-C1-P1-O7
4	A	302	ZQ7	F2-C1-P1-O5
4	A	302	ZQ7	F2-C1-P1-O7
6	A	304	ZP7	C19-C20-C21-O22
6	A	304	ZP7	C02-C03-C04-C05
4	A	302	ZQ7	O1-C29-C36-O6
4	A	302	ZQ7	C27-C29-C36-O6
4	A	302	ZQ7	C6-C12-C16-C17
4	A	302	ZQ7	C12-C16-C17-C18
4	A	302	ZQ7	C6-C10-C9-C5
6	A	304	ZP7	O18-C19-C20-O23
4	A	302	ZQ7	C11-C13-C14-C15
4	A	302	ZQ7	C18-C23-C32-C35
6	A	304	ZP7	O18-C19-C20-C21
4	A	302	ZQ7	C33-C34-C37-C38
6	A	304	ZP7	C05-C06-C07-C08
5	A	303	D12	C7-C8-C9-C10
5	A	303	D12	C2-C3-C4-C5
6	A	304	ZP7	O23-C20-C21-O22
4	A	302	ZQ7	C13-C11-C3-C21

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	304	ZP7	C03-C04-C05-C06
4	A	302	ZQ7	C35-O5-P1-O7
6	A	304	ZP7	C15-C16-O18-C19
4	A	302	ZQ7	C35-O5-P1-O8
4	A	302	ZQ7	C13-C11-C3-C7
4	A	302	ZQ7	C16-C17-C18-C30
6	A	304	ZP7	C07-C08-C09-C10
6	A	304	ZP7	O17-C16-O18-C19
6	A	304	ZP7	C04-C05-C06-C07
4	A	302	ZQ7	C16-C12-C6-C22
4	A	302	ZQ7	C24-C33-C34-C37
4	A	302	ZQ7	C16-C12-C6-C10
4	A	302	ZQ7	C16-C17-C18-C23
4	A	302	ZQ7	F1-C1-P1-O5
4	A	302	ZQ7	C17-C18-C23-C32
4	A	302	ZQ7	C30-C18-C23-C32
5	A	303	D12	C5-C6-C7-C8
4	A	302	ZQ7	C15-C20-C24-C33
4	A	302	ZQ7	C19-C2-C4-C8
6	A	304	ZP7	C01-C02-C03-C04

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/285 (98%)	0.52	21 (7%) 14 18	24, 35, 61, 85	0
2	B	103/108 (95%)	0.42	5 (4%) 29 35	27, 36, 53, 65	0
All	All	384/393 (97%)	0.49	26 (6%) 17 21	24, 36, 57, 85	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	107	GLY	4.8
1	A	87	GLN	4.7
1	A	90	SER	4.3
1	A	89	TYR	3.9
2	B	100	MET	3.8
1	A	108	GLY	3.8
1	A	86	SER	3.6
1	A	136	GLY	3.5
2	B	1	ALA	3.5
1	A	106	SER	3.5
1	A	5	SER	2.9
1	A	138	GLY	2.8
1	A	137	CYS	2.7
2	B	2	ILE	2.6
1	A	270	GLU	2.6
1	A	91	LYS	2.6
1	A	153	GLU	2.5
1	A	74	LEU	2.4
1	A	88	ASP	2.4
2	B	98	ARG	2.2
1	A	154	GLY	2.2
1	A	272	GLN	2.1
1	A	147	LEU	2.1
1	A	285	ARG	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	60	ASP	2.0
1	A	160	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	MLA	B	201	7/7	0.75	0.20	56,56,59,60	0
7	MLA	A	305	7/7	0.79	0.20	53,55,57,57	0
4	ZQ7	A	302	50/50	0.80	0.26	31,50,95,97	0
6	ZP7	A	304	23/23	0.80	0.36	42,51,68,70	0
3	NAG	A	301	14/15	0.91	0.13	38,42,44,47	0
5	D12	A	303	12/12	0.94	0.16	33,38,44,45	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ZQ7 A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ZP7 A 304:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.