



wwPDB NMR Structure Validation Summary Report ⓘ

May 29, 2020 – 12:05 am BST

PDB ID : 2N0A
Title : Atomic-resolution structure of alpha-synuclein fibrils
Authors : Tuttle, M.D.; Comellas, G.; Nieuwkoop, A.J.; Covell, D.J.; Berthold, D.A.;
Kloepper, K.D.; Courtney, J.M.; Kim, J.K.; Schwieters, C.D.; Lee, V.M.;
George, J.M.; Rienstra, C.M.
Deposited on : 2015-03-04

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLID-STATE NMR

The overall completeness of chemical shifts assignment is 2%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	140	83% 16% ..
1	B	140	88% 11% .
1	C	140	82% 17% .
1	D	140	82% 15% .
1	E	140	84% 16%
1	F	140	82% 16% .
1	G	140	74% 23% .
1	H	140	86% 14%
1	I	140	86% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	140	 <div>91%9%</div>

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 20160 atoms, of which 10030 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called Alpha-synuclein.


Mol	Chain	Residues	Atoms						Trace
1	A	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	B	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	C	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	D	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	E	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	F	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	G	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	H	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	I	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	
1	J	140	Total	C	H	N	O	S	0
			2016	627	1003	166	216	4	

4 Residue-property plots [i](#)


These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Alpha-synuclein

Chain A:  83% 16% ..





• Molecule 1: Alpha-synuclein

Chain B:  88% 11% .




• Molecule 1: Alpha-synuclein

Chain C:  82% 17% .





• Molecule 1: Alpha-synuclein

Chain D:  82% 15% .





• Molecule 1: Alpha-synuclein

Chain E:  84% 16%



• Molecule 1: Alpha-synuclein

Chain F:  82% 16% .




- Molecule 1: Alpha-synuclein

Chain G: 74% 23% .




- Molecule 1: Alpha-synuclein

Chain H: 86% 14%



- Molecule 1: Alpha-synuclein

Chain I: 86% 14%



- Molecule 1: Alpha-synuclein

Chain J: 91% 9%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 256 calculated structures, 1 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	geometry optimization	2.33.4
X-PLOR NIH	refinement	2.33.4
X-PLOR NIH	structure solution	2.33.4

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	277
Number of shifts mapped to atoms	277
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	2%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1013	1003	1003	17
1	B	1013	1003	1003	11
1	C	1013	1003	1003	16
1	D	1013	1003	1003	19
1	E	1013	1003	1003	14
1	F	1013	1003	1003	15
1	G	1013	1003	1003	23
1	H	1013	1003	1003	12
1	I	1013	1003	1003	11
1	J	1013	1003	1003	7
All	All	10130	10030	10030	114

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

5 of 114 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:G:116:MET:N	1:G:117:PRO:CD	0.59	2.65
1:D:137:GLU:N	1:D:138:PRO:CD	0.59	2.66
1:D:100:LEU:HD23	1:D:100:LEU:N	0.57	2.15
1:C:5:MET:SD	1:C:5:MET:N	0.56	2.77
1:D:100:LEU:N	1:D:100:LEU:CD2	0.54	2.71

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/140 (99%)	112 (81%)	21 (15%)	5 (4%)	6	34
1	B	138/140 (99%)	118 (86%)	17 (12%)	3 (2%)	10	49
1	C	138/140 (99%)	110 (80%)	19 (14%)	9 (7%)	2	18
1	D	138/140 (99%)	104 (75%)	23 (17%)	11 (8%)	2	14
1	E	138/140 (99%)	113 (82%)	18 (13%)	7 (5%)	4	24
1	F	138/140 (99%)	110 (80%)	18 (13%)	10 (7%)	2	16
1	G	138/140 (99%)	113 (82%)	15 (11%)	10 (7%)	2	16
1	H	138/140 (99%)	118 (86%)	12 (9%)	8 (6%)	3	21
1	I	138/140 (99%)	114 (83%)	18 (13%)	6 (4%)	5	29
1	J	138/140 (99%)	118 (86%)	15 (11%)	5 (4%)	6	34
All	All	1380/1400 (99%)	1130 (82%)	176 (13%)	74 (5%)	3	23

5 of 74 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	C	36	GLY
1	C	98	ASP
1	D	25	GLY
1	D	18	ALA
1	C	123	GLU

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/103 (100%)	99 (96%)	4 (4%)	36	84
1	B	103/103 (100%)	99 (96%)	4 (4%)	36	84
1	C	103/103 (100%)	102 (99%)	1 (1%)	77	96
1	D	103/103 (100%)	101 (98%)	2 (2%)	59	93
1	E	103/103 (100%)	102 (99%)	1 (1%)	77	96
1	F	103/103 (100%)	102 (99%)	1 (1%)	77	96
1	G	103/103 (100%)	100 (97%)	3 (3%)	45	89
1	H	103/103 (100%)	101 (98%)	2 (2%)	59	93
1	I	103/103 (100%)	102 (99%)	1 (1%)	77	96
1	J	103/103 (100%)	102 (99%)	1 (1%)	77	96
All	All	1030/1030 (100%)	1010 (98%)	20 (2%)	59	93

5 of 20 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	B	1	MET
1	G	116	MET
1	D	33	THR
1	D	100	LEU
1	G	22	THR

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 2% for the well-defined parts and 2% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	277
Number of shifts mapped to atoms	277
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	66	0.03 ± 0.23	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	47	-0.72 ± 0.26	Should be applied
$^{13}\text{C}'$	55	0.57 ± 0.14	Should be applied
^{15}N	63	-2.62 ± 0.38	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 2%, i.e. 277 atoms were assigned a chemical shift out of a possible 15290. 0 out of 230 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	230/6900 (3%)	46/2750 (2%)	121/2800 (4%)	63/1350 (5%)
Sidechain	47/7820 (1%)	0/4500 (0%)	47/3080 (2%)	0/240 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	0/570 (0%)	0/300 (0%)	0/260 (0%)	0/10 (0%)
Overall	277/15290 (2%)	46/7550 (1%)	168/6140 (3%)	63/1600 (4%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

