



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 10:35 pm BST

PDB ID : 4N0P  
Title : Crystal structure of a pilus assembly protein CpaE (CC\_2943) from *Caulobacter crescentus* CB15 at 1.75 Å resolution (PSI Community Target, Shapiro)  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2013-10-02  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	<div><div>%</div><div><div></div><div>85%</div><div>•</div><div>11%</div></div></div>
1	B	151	<div><div>%</div><div><div></div><div>82%</div><div>5%</div><div>13%</div></div></div>
1	C	151	<div><div></div><div><div></div><div>85%</div><div>•</div><div>12%</div></div></div>
1	D	151	<div><div>4%</div><div><div></div><div>78%</div><div>9%</div><div>13%</div></div></div>
1	E	151	<div><div>%</div><div><div></div><div>76%</div><div>5%</div><div>19%</div></div></div>
1	F	151	<div><div>5%</div><div><div></div><div>84%</div><div>5%</div><div>11%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	151	<div><div>%</div><div><div></div><div>83%</div><div>13%</div></div></div>
1	H	151	<div><div>3%</div><div><div></div><div>73%</div><div>9%</div><div>18%</div></div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pilus assembly protein CpaE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	S	0	7	0
			1061	669	187	199	6			
1	B	131	Total	C	N	O	S	0	8	0
			1046	662	185	193	6			
1	C	133	Total	C	N	O	S	0	5	0
			1045	658	186	196	5			
1	D	132	Total	C	N	O	S	0	6	0
			1040	661	185	188	6			
1	E	123	Total	C	N	O	S	0	6	0
			975	616	177	177	5			
1	F	134	Total	C	N	O	S	0	5	0
			1046	662	185	192	7			
1	G	132	Total	C	N	O	S	0	1	0
			1024	646	184	189	5			
1	H	124	Total	C	N	O	S	0	4	0
			967	610	172	179	6			

There are 152 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	leader sequence	UNP Q9A495
A	-17	GLY	-	leader sequence	UNP Q9A495
A	-16	SER	-	leader sequence	UNP Q9A495
A	-15	ASP	-	leader sequence	UNP Q9A495
A	-14	LYS	-	leader sequence	UNP Q9A495
A	-13	ILE	-	leader sequence	UNP Q9A495
A	-12	HIS	-	leader sequence	UNP Q9A495
A	-11	HIS	-	leader sequence	UNP Q9A495
A	-10	HIS	-	leader sequence	UNP Q9A495
A	-9	HIS	-	leader sequence	UNP Q9A495
A	-8	HIS	-	leader sequence	UNP Q9A495
A	-7	HIS	-	leader sequence	UNP Q9A495
A	-6	GLU	-	leader sequence	UNP Q9A495

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	ASN	-	leader sequence	UNP Q9A495
A	-4	LEU	-	leader sequence	UNP Q9A495
A	-3	TYR	-	leader sequence	UNP Q9A495
A	-2	PHE	-	leader sequence	UNP Q9A495
A	-1	GLN	-	leader sequence	UNP Q9A495
A	0	GLY	-	leader sequence	UNP Q9A495
B	-18	MET	-	leader sequence	UNP Q9A495
B	-17	GLY	-	leader sequence	UNP Q9A495
B	-16	SER	-	leader sequence	UNP Q9A495
B	-15	ASP	-	leader sequence	UNP Q9A495
B	-14	LYS	-	leader sequence	UNP Q9A495
B	-13	ILE	-	leader sequence	UNP Q9A495
B	-12	HIS	-	leader sequence	UNP Q9A495
B	-11	HIS	-	leader sequence	UNP Q9A495
B	-10	HIS	-	leader sequence	UNP Q9A495
B	-9	HIS	-	leader sequence	UNP Q9A495
B	-8	HIS	-	leader sequence	UNP Q9A495
B	-7	HIS	-	leader sequence	UNP Q9A495
B	-6	GLU	-	leader sequence	UNP Q9A495
B	-5	ASN	-	leader sequence	UNP Q9A495
B	-4	LEU	-	leader sequence	UNP Q9A495
B	-3	TYR	-	leader sequence	UNP Q9A495
B	-2	PHE	-	leader sequence	UNP Q9A495
B	-1	GLN	-	leader sequence	UNP Q9A495
B	0	GLY	-	leader sequence	UNP Q9A495
C	-18	MET	-	leader sequence	UNP Q9A495
C	-17	GLY	-	leader sequence	UNP Q9A495
C	-16	SER	-	leader sequence	UNP Q9A495
C	-15	ASP	-	leader sequence	UNP Q9A495
C	-14	LYS	-	leader sequence	UNP Q9A495
C	-13	ILE	-	leader sequence	UNP Q9A495
C	-12	HIS	-	leader sequence	UNP Q9A495
C	-11	HIS	-	leader sequence	UNP Q9A495
C	-10	HIS	-	leader sequence	UNP Q9A495
C	-9	HIS	-	leader sequence	UNP Q9A495
C	-8	HIS	-	leader sequence	UNP Q9A495
C	-7	HIS	-	leader sequence	UNP Q9A495
C	-6	GLU	-	leader sequence	UNP Q9A495
C	-5	ASN	-	leader sequence	UNP Q9A495
C	-4	LEU	-	leader sequence	UNP Q9A495
C	-3	TYR	-	leader sequence	UNP Q9A495
C	-2	PHE	-	leader sequence	UNP Q9A495

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLN	-	leader sequence	UNP Q9A495
C	0	GLY	-	leader sequence	UNP Q9A495
D	-18	MET	-	leader sequence	UNP Q9A495
D	-17	GLY	-	leader sequence	UNP Q9A495
D	-16	SER	-	leader sequence	UNP Q9A495
D	-15	ASP	-	leader sequence	UNP Q9A495
D	-14	LYS	-	leader sequence	UNP Q9A495
D	-13	ILE	-	leader sequence	UNP Q9A495
D	-12	HIS	-	leader sequence	UNP Q9A495
D	-11	HIS	-	leader sequence	UNP Q9A495
D	-10	HIS	-	leader sequence	UNP Q9A495
D	-9	HIS	-	leader sequence	UNP Q9A495
D	-8	HIS	-	leader sequence	UNP Q9A495
D	-7	HIS	-	leader sequence	UNP Q9A495
D	-6	GLU	-	leader sequence	UNP Q9A495
D	-5	ASN	-	leader sequence	UNP Q9A495
D	-4	LEU	-	leader sequence	UNP Q9A495
D	-3	TYR	-	leader sequence	UNP Q9A495
D	-2	PHE	-	leader sequence	UNP Q9A495
D	-1	GLN	-	leader sequence	UNP Q9A495
D	0	GLY	-	leader sequence	UNP Q9A495
E	-18	MET	-	leader sequence	UNP Q9A495
E	-17	GLY	-	leader sequence	UNP Q9A495
E	-16	SER	-	leader sequence	UNP Q9A495
E	-15	ASP	-	leader sequence	UNP Q9A495
E	-14	LYS	-	leader sequence	UNP Q9A495
E	-13	ILE	-	leader sequence	UNP Q9A495
E	-12	HIS	-	leader sequence	UNP Q9A495
E	-11	HIS	-	leader sequence	UNP Q9A495
E	-10	HIS	-	leader sequence	UNP Q9A495
E	-9	HIS	-	leader sequence	UNP Q9A495
E	-8	HIS	-	leader sequence	UNP Q9A495
E	-7	HIS	-	leader sequence	UNP Q9A495
E	-6	GLU	-	leader sequence	UNP Q9A495
E	-5	ASN	-	leader sequence	UNP Q9A495
E	-4	LEU	-	leader sequence	UNP Q9A495
E	-3	TYR	-	leader sequence	UNP Q9A495
E	-2	PHE	-	leader sequence	UNP Q9A495
E	-1	GLN	-	leader sequence	UNP Q9A495
E	0	GLY	-	leader sequence	UNP Q9A495
F	-18	MET	-	leader sequence	UNP Q9A495
F	-17	GLY	-	leader sequence	UNP Q9A495

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	-16	SER	-	leader sequence	UNP Q9A495
F	-15	ASP	-	leader sequence	UNP Q9A495
F	-14	LYS	-	leader sequence	UNP Q9A495
F	-13	ILE	-	leader sequence	UNP Q9A495
F	-12	HIS	-	leader sequence	UNP Q9A495
F	-11	HIS	-	leader sequence	UNP Q9A495
F	-10	HIS	-	leader sequence	UNP Q9A495
F	-9	HIS	-	leader sequence	UNP Q9A495
F	-8	HIS	-	leader sequence	UNP Q9A495
F	-7	HIS	-	leader sequence	UNP Q9A495
F	-6	GLU	-	leader sequence	UNP Q9A495
F	-5	ASN	-	leader sequence	UNP Q9A495
F	-4	LEU	-	leader sequence	UNP Q9A495
F	-3	TYR	-	leader sequence	UNP Q9A495
F	-2	PHE	-	leader sequence	UNP Q9A495
F	-1	GLN	-	leader sequence	UNP Q9A495
F	0	GLY	-	leader sequence	UNP Q9A495
G	-18	MET	-	leader sequence	UNP Q9A495
G	-17	GLY	-	leader sequence	UNP Q9A495
G	-16	SER	-	leader sequence	UNP Q9A495
G	-15	ASP	-	leader sequence	UNP Q9A495
G	-14	LYS	-	leader sequence	UNP Q9A495
G	-13	ILE	-	leader sequence	UNP Q9A495
G	-12	HIS	-	leader sequence	UNP Q9A495
G	-11	HIS	-	leader sequence	UNP Q9A495
G	-10	HIS	-	leader sequence	UNP Q9A495
G	-9	HIS	-	leader sequence	UNP Q9A495
G	-8	HIS	-	leader sequence	UNP Q9A495
G	-7	HIS	-	leader sequence	UNP Q9A495
G	-6	GLU	-	leader sequence	UNP Q9A495
G	-5	ASN	-	leader sequence	UNP Q9A495
G	-4	LEU	-	leader sequence	UNP Q9A495
G	-3	TYR	-	leader sequence	UNP Q9A495
G	-2	PHE	-	leader sequence	UNP Q9A495
G	-1	GLN	-	leader sequence	UNP Q9A495
G	0	GLY	-	leader sequence	UNP Q9A495
H	-18	MET	-	leader sequence	UNP Q9A495
H	-17	GLY	-	leader sequence	UNP Q9A495
H	-16	SER	-	leader sequence	UNP Q9A495
H	-15	ASP	-	leader sequence	UNP Q9A495
H	-14	LYS	-	leader sequence	UNP Q9A495
H	-13	ILE	-	leader sequence	UNP Q9A495

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
H	-12	HIS	-	leader sequence	UNP Q9A495
H	-11	HIS	-	leader sequence	UNP Q9A495
H	-10	HIS	-	leader sequence	UNP Q9A495
H	-9	HIS	-	leader sequence	UNP Q9A495
H	-8	HIS	-	leader sequence	UNP Q9A495
H	-7	HIS	-	leader sequence	UNP Q9A495
H	-6	GLU	-	leader sequence	UNP Q9A495
H	-5	ASN	-	leader sequence	UNP Q9A495
H	-4	LEU	-	leader sequence	UNP Q9A495
H	-3	TYR	-	leader sequence	UNP Q9A495
H	-2	PHE	-	leader sequence	UNP Q9A495
H	-1	GLN	-	leader sequence	UNP Q9A495
H	0	GLY	-	leader sequence	UNP Q9A495

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	79	Total O 79 79	0	0
2	B	73	Total O 73 73	0	0
2	C	89	Total O 90 90	0	1
2	D	90	Total O 90 90	0	0
2	E	75	Total O 75 75	0	0
2	F	43	Total O 43 43	0	0
2	G	79	Total O 79 79	0	0
2	H	75	Total O 75 75	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pilus assembly protein CpaE





- Molecule 1: Pilus assembly protein CpaE





- Molecule 1: Pilus assembly protein CpaE




- Molecule 1: Pilus assembly protein CpaE




- Molecule 1: Pilus assembly protein CpaE





- Molecule 1: Pilus assembly protein CpaE



- Molecule 1: Pilus assembly protein CpaE



- Molecule 1: Pilus assembly protein CpaE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.60 Å   112.78 Å   73.88 Å 90.00°   94.12°   90.00°	Depositor
Resolution (Å)	47.27 – 1.75 47.27 – 1.75	Depositor EDS
% Data completeness (in resolution range)	97.8 (47.27-1.75) 97.8 (47.27-1.75)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 1.75 Å)	Xtriage
Refinement program	BUSTER-TNT 2.10.0, BUSTER 2.10.0	Depositor
R, $R_{free}$	0.181   ,   0.205 0.188   ,   0.217	Depositor DCC
$R_{free}$ test set	5254 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.4	Xtriage
Anisotropy	0.304	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33   ,   43.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8808	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/1093	0.61	0/1487
1	B	0.51	0/1083	0.54	0/1474
1	C	0.50	0/1070	0.56	0/1456
1	D	0.50	0/1074	0.58	0/1461
1	E	0.53	0/1006	0.55	0/1367
1	F	0.46	0/1077	0.57	0/1465
1	G	0.48	0/1044	0.55	0/1420
1	H	0.50	0/993	0.56	0/1350
All	All	0.50	0/8440	0.57	0/11480

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1061	0	1083	3	0
1	B	1046	0	1083	4	0
1	C	1045	0	1062	3	0
1	D	1040	0	1087	16	0
1	E	975	0	1028	5	0
1	F	1046	0	1077	4	0
1	G	1024	0	1046	5	0
1	H	967	0	1005	14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	79	0	0	1	0
2	B	73	0	0	0	0
2	C	90	0	0	1	0
2	D	90	0	0	1	0
2	E	75	0	0	0	0
2	F	43	0	0	0	0
2	G	79	0	0	0	0
2	H	75	0	0	0	0
All	All	8808	0	8471	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:143:GLU:HB2	1:H:157:ILE:HD11	1.37	1.02
1:D:143:GLU:CB	1:H:157:ILE:HD11	1.93	0.97
1:D:143:GLU:HG3	1:H:157:ILE:HD11	1.49	0.95
1:D:143:GLU:CG	1:H:157:ILE:HD11	1.98	0.91
1:D:155:ALA:HB2	1:H:152:SER:O	1.82	0.80
1:D:143:GLU:HB2	1:H:157:ILE:CD1	2.16	0.73
1:F:133[B]:CYS:SG	1:F:135:ARG:O	2.47	0.71
1:E:183:THR:HG21	1:E:190:LEU:HD13	1.73	0.70
1:D:143:GLU:HG3	1:H:157:ILE:CD1	2.22	0.68
1:E:192:HIS:CD2	1:F:234:PRO:HG2	2.33	0.64
1:D:129:ILE:HB	1:D:156[B]:THR:HG22	1.83	0.60
1:H:183:THR:HG21	1:H:190:LEU:HD13	1.86	0.57
1:H:128[B]:THR:HG23	1:H:177:SER:H	1.74	0.52
1:D:138:THR:OG1	1:D:233:GLN:NE2	2.42	0.52
1:A:233:GLN:HG2	1:A:235:LEU:HD12	1.92	0.51
1:E:178[A]:LEU:HD11	1:E:208:VAL:HG23	1.95	0.48
1:D:157:ILE:HG12	1:H:146:ALA:HB1	1.95	0.48
1:A:143:GLU:HB3	1:B:149:ARG:HH12	1.79	0.48
2:A:348:HOH:O	1:B:143:GLU:HG2	2.14	0.47
1:H:178:LEU:HD11	1:H:208:VAL:HG23	1.98	0.46
1:D:178[A]:LEU:HD11	1:D:208:VAL:HG23	1.99	0.45
1:D:143:GLU:CG	1:H:157:ILE:CD1	2.83	0.44
1:H:179:VAL:HG12	1:H:181:VAL:HG13	2.01	0.43
1:D:124:ILE:HD13	1:D:151:MET:SD	2.59	0.43
1:G:134:ALA:HB1	1:G:184:LEU:HG	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:134:ALA:HB1	1:F:184:LEU:HG	2.00	0.42
1:C:126:ARG:CZ	1:G:126:ARG:HD2	2.50	0.42
1:D:203:PRO:HB3	1:G:153:ARG:CZ	2.50	0.42
1:C:134:ALA:HB1	1:C:184:LEU:HG	2.01	0.42
1:F:130:HIS:CG	1:F:176:PRO:HG3	2.55	0.42
1:H:130:HIS:CG	1:H:176:PRO:HG3	2.55	0.41
1:D:200:VAL:HA	1:G:149:ARG:HG3	2.02	0.41
1:E:137:GLU:HA	1:G:157:ILE:HD11	2.02	0.41
1:A:178:LEU:HD11	1:A:208:VAL:HG23	2.03	0.41
1:B:179:VAL:HG12	1:B:181:VAL:HG13	2.03	0.41
1:C:137:GLU:HG3	2:C:344:HOH:O	2.21	0.41
1:B:178:LEU:HD11	1:B:208:VAL:HG23	2.03	0.41
1:D:139:ALA:HB2	2:D:356:HOH:O	2.21	0.40
1:E:178[B]:LEU:HD11	1:E:247:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/151 (92%)	138 (99%)	1 (1%)	0	100	100
1	B	137/151 (91%)	135 (98%)	2 (2%)	0	100	100
1	C	136/151 (90%)	134 (98%)	2 (2%)	0	100	100
1	D	136/151 (90%)	134 (98%)	2 (2%)	0	100	100
1	E	127/151 (84%)	126 (99%)	1 (1%)	0	100	100
1	F	137/151 (91%)	136 (99%)	1 (1%)	0	100	100
1	G	131/151 (87%)	130 (99%)	1 (1%)	0	100	100
1	H	126/151 (83%)	125 (99%)	1 (1%)	0	100	100
All	All	1069/1208 (88%)	1058 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/122 (93%)	112 (99%)	1 (1%)	78	67
1	B	112/122 (92%)	111 (99%)	1 (1%)	78	67
1	C	110/122 (90%)	110 (100%)	0	100	100
1	D	111/122 (91%)	111 (100%)	0	100	100
1	E	105/122 (86%)	104 (99%)	1 (1%)	76	63
1	F	111/122 (91%)	111 (100%)	0	100	100
1	G	108/122 (88%)	108 (100%)	0	100	100
1	H	104/122 (85%)	103 (99%)	1 (1%)	76	63
All	All	874/976 (90%)	870 (100%)	4 (0%)	88	83

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-6	GLU
1	B	141	LEU
1	E	141	LEU
1	H	126	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	233	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	134/151 (88%)	-0.10	2 (1%) 73 80	20, 28, 52, 81	0
1	B	131/151 (86%)	-0.10	1 (0%) 86 90	19, 27, 51, 83	0
1	C	133/151 (88%)	-0.24	0 100 100	19, 27, 42, 53	0
1	D	132/151 (87%)	0.07	6 (4%) 33 39	17, 28, 55, 68	0
1	E	123/151 (81%)	-0.11	1 (0%) 86 90	19, 27, 47, 63	0
1	F	134/151 (88%)	0.34	8 (5%) 21 27	23, 37, 63, 91	0
1	G	132/151 (87%)	-0.02	2 (1%) 73 80	19, 31, 56, 65	0
1	H	124/151 (82%)	0.10	5 (4%) 38 45	18, 28, 70, 90	0
All	All	1043/1208 (86%)	-0.00	25 (2%) 59 65	17, 29, 56, 91	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	249	ALA	10.5
1	H	152	SER	6.0
1	H	149	ARG	5.1
1	A	249	ALA	4.7
1	D	235	LEU	4.3
1	H	126	ARG	4.1
1	D	-5	ASN	3.8
1	H	153	ARG	3.6
1	D	147	ALA	3.5
1	F	172	ASN	3.3
1	H	125	PRO	3.1
1	G	248	TYR	3.0
1	F	202	ASP	2.9
1	G	234	PRO	2.8
1	F	201[A]	CYS	2.7
1	B	250	ASP	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	248	TYR	2.5
1	F	192	HIS	2.5
1	D	149	ARG	2.5
1	F	171	GLN	2.3
1	D	238	LEU	2.3
1	F	234	PRO	2.2
1	A	172	ASN	2.2
1	F	168	ASP	2.2
1	D	234	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.