



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2020 – 05:54 AM BST

PDB ID : 6N3R
Title : Sheep Galectin-11 (LGALS11) complex with galactose
Authors : Beddoe, T.C.; Sakthivel, D.
Deposited on : 2018-11-16
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>
1	B	139	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>5%</div> <div>.</div> </div> </div>
1	C	139	<div> <div>4%</div> <div> <div></div> <div>93%</div> <div>5%</div> <div>.</div> </div> </div>
1	D	139	<div> <div>3%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>.</div> </div> </div>
1	E	139	<div> <div>4%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>.</div> </div> </div>
1	F	139	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	139	<div><div></div><div>4%</div><div>80%</div><div>14%</div><div>6%</div></div>
1	H	139	<div><div></div><div>%</div><div>88%</div><div>9%</div><div>•</div></div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	1	0
			1097	712	178	200	7			
1	B	135	Total	C	N	O	S	0	0	0
			1064	691	173	194	6			
1	C	136	Total	C	N	O	S	0	0	0
			1061	687	173	195	6			
1	D	135	Total	C	N	O	S	0	0	0
			1064	691	173	194	6			
1	E	135	Total	C	N	O	S	0	0	0
			1054	682	172	194	6			
1	F	135	Total	C	N	O	S	0	1	0
			1067	692	174	195	6			
1	G	131	Total	C	N	O	S	0	1	0
			1043	681	168	188	6			
1	H	135	Total	C	N	O	S	0	1	0
			1069	693	173	197	6			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	PRO	-	expression tag	UNP A0A0A7EMW6
A	0	GLY	-	expression tag	UNP A0A0A7EMW6
B	-1	PRO	-	expression tag	UNP A0A0A7EMW6
B	0	GLY	-	expression tag	UNP A0A0A7EMW6
C	-1	PRO	-	expression tag	UNP A0A0A7EMW6
C	0	GLY	-	expression tag	UNP A0A0A7EMW6
D	-1	PRO	-	expression tag	UNP A0A0A7EMW6
D	0	GLY	-	expression tag	UNP A0A0A7EMW6
E	-1	PRO	-	expression tag	UNP A0A0A7EMW6
E	0	GLY	-	expression tag	UNP A0A0A7EMW6
F	-1	PRO	-	expression tag	UNP A0A0A7EMW6
F	0	GLY	-	expression tag	UNP A0A0A7EMW6
G	-1	PRO	-	expression tag	UNP A0A0A7EMW6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	GLY	-	expression tag	UNP A0A0A7EMW6
H	-1	PRO	-	expression tag	UNP A0A0A7EMW6
H	0	GLY	-	expression tag	UNP A0A0A7EMW6

- Molecule 2 is beta-D-galactopyranose (three-letter code: GAL) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			12	6	6		
2	F	1	Total	C	O	0	0
			12	6	6		
2	H	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	155	Total	O	0	0
			155	155		
3	B	145	Total	O	0	0
			145	145		
3	C	119	Total	O	0	0
			119	119		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	134	Total 134	O 134	0	0
3	E	133	Total 133	O 133	0	0
3	F	128	Total 128	O 128	0	0
3	G	124	Total 124	O 124	0	0
3	H	126	Total 126	O 126	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Galectin




- Molecule 1: Galectin





- Molecule 1: Galectin




- Molecule 1: Galectin





- Molecule 1: Galectin




- Molecule 1: Galectin



● Molecule 1: Galectin



● Molecule 1: Galectin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.11Å 127.43Å 140.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.17 – 2.40 47.17 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.17-2.40) 100.0 (47.17-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.26 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.10 _2148	Depositor
R, R_{free}	0.160 , 0.209 0.160 , 0.209	Depositor DCC
R_{free} test set	3454 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9643	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1128	0.56	0/1531
1	B	0.42	0/1091	0.56	0/1483
1	C	0.39	0/1087	0.56	0/1479
1	D	0.39	0/1091	0.57	0/1483
1	E	0.42	0/1080	0.58	0/1470
1	F	0.40	0/1094	0.58	0/1489
1	G	0.40	0/1072	0.57	0/1456
1	H	0.37	0/1096	0.55	0/1491
All	All	0.40	0/8739	0.57	0/11882

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1097	0	1098	7	1
1	B	1064	0	1059	3	0
1	C	1061	0	1047	4	0
1	D	1064	0	1059	8	0
1	E	1054	0	1041	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1067	0	1050	5	0
1	G	1043	0	1041	12	0
1	H	1069	0	1053	8	1
2	B	12	0	11	0	0
2	C	12	0	12	0	0
2	D	12	0	12	0	0
2	F	12	0	12	1	0
2	H	12	0	12	1	0
3	A	155	0	0	1	1
3	B	145	0	0	1	0
3	C	119	0	0	2	0
3	D	134	0	0	4	0
3	E	133	0	0	4	0
3	F	128	0	0	1	0
3	G	124	0	0	3	0
3	H	126	0	0	0	1
All	All	9643	0	8507	54	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 54 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:23:LYS:HE3	3:C:301:HOH:O	1.70	0.90
1:D:3:SER:N	3:D:301:HOH:O	2.08	0.85
1:E:46:GLN:O	3:E:201:HOH:O	2.00	0.78
1:B:3:SER:N	3:B:301:HOH:O	2.19	0.76
1:D:3:SER:N	3:D:303:HOH:O	2.21	0.72

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:GLY:O	1:H:28:SER:OG[4_577]	2.13	0.07
3:A:320:HOH:O	3:H:417:HOH:O[4_577]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/139 (99%)	135 (98%)	3 (2%)	0	100	100
1	B	133/139 (96%)	130 (98%)	3 (2%)	0	100	100
1	C	134/139 (96%)	131 (98%)	3 (2%)	0	100	100
1	D	133/139 (96%)	131 (98%)	2 (2%)	0	100	100
1	E	133/139 (96%)	129 (97%)	4 (3%)	0	100	100
1	F	134/139 (96%)	132 (98%)	2 (2%)	0	100	100
1	G	128/139 (92%)	125 (98%)	3 (2%)	0	100	100
1	H	134/139 (96%)	132 (98%)	2 (2%)	0	100	100
All	All	1067/1112 (96%)	1045 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/124 (100%)	123 (99%)	1 (1%)	81	91
1	B	120/124 (97%)	118 (98%)	2 (2%)	60	78
1	C	118/124 (95%)	117 (99%)	1 (1%)	81	91
1	D	120/124 (97%)	118 (98%)	2 (2%)	60	78
1	E	118/124 (95%)	117 (99%)	1 (1%)	81	91
1	F	119/124 (96%)	118 (99%)	1 (1%)	81	91

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	118/124 (95%)	117 (99%)	1 (1%)	81	91
1	H	120/124 (97%)	119 (99%)	1 (1%)	81	91
All	All	957/992 (96%)	947 (99%)	10 (1%)	76	88

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	66	LEU
1	D	77	LEU
1	F	77	LEU
1	C	110	GLN
1	E	77	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GAL	H	201	-	12,12,12	0.55	0	17,17,17	1.37	4 (23%)
2	GAL	B	201	-	12,12,12	0.86	0	17,17,17	1.26	3 (17%)
2	GAL	C	201	-	12,12,12	0.78	0	17,17,17	1.35	2 (11%)
2	GAL	D	201	-	12,12,12	0.79	1 (8%)	17,17,17	1.27	1 (5%)
2	GAL	F	201	-	12,12,12	0.71	0	17,17,17	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	H	201	-	-	1/2/22/22	0/1/1/1
2	GAL	B	201	-	-	0/2/22/22	0/1/1/1
2	GAL	C	201	-	-	0/2/22/22	0/1/1/1
2	GAL	D	201	-	-	2/2/22/22	0/1/1/1
2	GAL	F	201	-	-	2/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	201	GAL	C4-C5	2.07	1.57	1.53

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	GAL	O5-C5-C4	2.78	114.75	109.69
2	C	201	GAL	O5-C1-C2	-2.78	105.33	110.28
2	B	201	GAL	O5-C5-C4	2.39	114.03	109.69
2	H	201	GAL	O5-C1-C2	-2.36	106.08	110.28
2	H	201	GAL	C1-C2-C3	-2.34	105.45	110.31

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	201	GAL	O5-C5-C6-O6
2	D	201	GAL	C4-C5-C6-O6
2	F	201	GAL	O5-C5-C6-O6
2	F	201	GAL	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	H	201	GAL	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	201	GAL	1	0
2	F	201	GAL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	139/139 (100%)	-0.04	6 (4%) 35 33	13, 21, 50, 65	0
1	B	135/139 (97%)	-0.17	2 (1%) 73 72	13, 19, 33, 54	0
1	C	136/139 (97%)	0.02	5 (3%) 41 41	15, 22, 50, 72	0
1	D	135/139 (97%)	-0.05	4 (2%) 50 49	14, 22, 38, 59	0
1	E	135/139 (97%)	-0.09	5 (3%) 41 41	12, 20, 40, 59	0
1	F	135/139 (97%)	-0.21	4 (2%) 50 49	13, 21, 40, 58	0
1	G	131/139 (94%)	-0.11	5 (3%) 40 39	14, 23, 40, 44	0
1	H	135/139 (97%)	-0.08	2 (1%) 73 72	15, 25, 40, 53	0
All	All	1081/1112 (97%)	-0.09	33 (3%) 49 47	12, 22, 41, 72	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	30	PHE	7.0
1	E	30	PHE	5.6
1	C	31	GLY	4.8
1	A	0	GLY	4.6
1	F	44	THR	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GAL	F	201	12/12	0.85	0.25	39,46,52,62	0
2	GAL	D	201	12/12	0.87	0.21	37,40,46,54	0
2	GAL	H	201	12/12	0.91	0.23	38,42,55,56	0
2	GAL	B	201	12/12	0.93	0.16	29,35,46,53	0
2	GAL	C	201	12/12	0.94	0.22	29,39,43,61	0

6.5 Other polymers

There are no such residues in this entry.