



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 02:25 PM BST

PDB ID : 3N43
Title : Crystal structures of the mature envelope glycoprotein complex (trypsin cleavage) of Chikungunya virus.
Authors : Voss, J.; Vaney, M.C.; Duquerroy, S.; Rey, F.A.
Deposited on : 2010-05-21
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	65	<div> <div>38%</div> <div> <div>77%</div> <div>22%</div> </div> </div>
2	B	360	<div> <div>9%</div> <div> <div>89%</div> <div>7%</div> </div> </div>
3	F	473	<div> <div>2%</div> <div> <div>80%</div> <div>17%</div> </div> </div>
4	C	2	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	B	2001	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	51	Total	C	N	O	S	0	0	0
			400	250	63	78	9			

- Molecule 2 is a protein called E2 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	336	Total	C	N	O	S	0	1	0
			2650	1653	480	497	20			

- Molecule 3 is a protein called E1 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	393	Total	C	N	O	S	0	0	0
			2989	1890	501	574	24			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-1	GLY	-	SEE REMARK 999	UNP Q1H8W5
F	0	GLY	-	SEE REMARK 999	UNP Q1H8W5

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	F	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	46	Total 46	O 46	0	0
7	F	107	Total 107	O 107	0	0


- Molecule 1: E3 envelope glycoprotein



- Chain B:  9% 89% 7%



- Chain F: 



- Chain C:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.41Å 90.81Å 179.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.00 – 2.58 48.95 – 2.58	Depositor EDS
% Data completeness (in resolution range)	97.8 (44.00-2.58) 97.7 (48.95-2.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.9.3	Depositor
R, R_{free}	0.245 , 0.257 0.260 , 0.272	Depositor DCC
R_{free} test set	1532 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtriage
Anisotropy	0.692	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6238	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, NDG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/409	0.54	0/557
2	B	0.36	0/2712	0.56	0/3691
3	F	0.36	0/3066	0.54	0/4182
All	All	0.36	0/6187	0.55	0/8430

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	400	0	381	0	0
2	B	2650	0	2572	4	0
3	F	2989	0	2893	7	0
4	C	28	0	24	0	0
5	B	14	0	13	0	0
6	F	4	0	3	0	0
7	B	46	0	0	0	0
7	F	107	0	0	0	0
All	All	6238	0	5886	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:60:VAL:HG22	3:F:102:GLN:HG3	1.88	0.55
3:F:42:LEU:HD11	3:F:266:VAL:HG22	1.89	0.53
3:F:38:LEU:HB2	3:F:268:ALA:HB3	1.90	0.52
2:B:38:ARG:HB2	2:B:47:LYS:HB3	1.91	0.51
2:B:248:LEU:HD13	2:B:251:ARG:HB2	1.94	0.50
3:F:28:VAL:HG23	3:F:329:ALA:HB1	1.94	0.48
3:F:263:THR:HG22	3:F:266:VAL:HG12	2.00	0.43
3:F:41:THR:HB	3:F:125:HIS:HB2	2.00	0.42
2:B:13:ARG:HE	2:B:236:GLN:HE21	1.68	0.42
3:F:19:VAL:HB	3:F:27:MET:HB3	2.00	0.42
2:B:172:PRO:HG3	2:B:236:GLN:HB2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	49/65 (75%)	46 (94%)	2 (4%)	1 (2%)	7	13
2	B	334/360 (93%)	317 (95%)	16 (5%)	1 (0%)	41	62
3	F	391/473 (83%)	383 (98%)	8 (2%)	0	100	100
All	All	774/898 (86%)	746 (96%)	26 (3%)	2 (0%)	41	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	ALA
2	B	72	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	48/61 (79%)	48 (100%)	0	100	100
2	B	297/319 (93%)	290 (98%)	7 (2%)	49	72
3	F	328/378 (87%)	326 (99%)	2 (1%)	86	94
All	All	673/758 (89%)	664 (99%)	9 (1%)	69	85

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	28	CYS
2	B	160	THR
2	B	178	ARG
2	B	206	SER
2	B	208	GLU
2	B	220	CYS
2	B	235	TRP
3	F	192	PHE
3	F	370	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	236	GLN
2	B	282	GLN
3	F	102	GLN
3	F	373	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	C	1	3,4	14,14,15	1.19	1 (7%)	17,19,21	0.88	0
4	NDG	C	2	4	14,14,15	1.45	1 (7%)	17,19,21	1.36	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	3,4	-	0/6/23/26	0/1/1/1
4	NDG	C	2	4	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	2	NDG	C1-C2	3.86	1.58	1.52
4	C	1	NAG	C1-C2	3.01	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	2	NDG	C1-C2-N2	3.60	116.64	110.49

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	2	NDG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	2001	2	14,14,15	1.24	1 (7%)	17,19,21	0.91	0
6	ACT	F	452	-	1,3,3	1.81	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	2001	2	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	2001	NAG	C1-C2	2.97	1.56	1.52

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	2001	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	51/65 (78%)	2.51	25 (49%) 0 0	78, 92, 110, 116	0
2	B	336/360 (93%)	0.77	31 (9%) 9 7	27, 49, 71, 95	0
3	F	393/473 (83%)	0.21	8 (2%) 65 62	19, 32, 55, 71	0
All	All	780/898 (86%)	0.61	64 (8%) 11 9	19, 41, 83, 116	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	LEU	10.0
1	A	10	LEU	6.1
2	B	206	SER	5.7
1	A	13	THR	5.3
1	A	11	ALA	5.2
2	B	205	GLY	5.1
2	B	250	ASP	4.8
1	A	15	PHE	4.7
1	A	8	CYS	4.6
1	A	22	CYS	4.0
2	B	248	LEU	3.9
1	A	14	THR	3.8
1	A	7	MET	3.7
2	B	185	SER	3.6
2	B	93	ILE	3.6
1	A	12	ASN	3.6
1	A	49	GLN	3.5
1	A	21	PRO	3.5
1	A	31	PRO	3.4
2	B	163	THR	3.2
3	F	212	ASP	3.2
1	A	19	GLN	3.0
1	A	23	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	-1	GLY	2.8
2	B	158	GLN	2.8
1	A	26	CYS	2.8
1	A	27	TYR	2.8
1	A	44	ARG	2.8
2	B	219	ASN	2.7
2	B	249	GLY	2.7
2	B	160	THR	2.6
3	F	126	THR	2.6
2	B	288	TYR	2.6
2	B	231	ASN	2.5
2	B	122	SER	2.5
2	B	247	GLU	2.5
2	B	67	LEU	2.5
1	A	53	ALA	2.4
2	B	142	HIS	2.4
2	B	120	LYS	2.4
2	B	25	GLY	2.3
1	A	30	GLU	2.3
2	B	223	ASP	2.3
3	F	108	VAL	2.3
2	B	84	PHE	2.2
2	B	113	VAL	2.2
2	B	65	THR	2.2
2	B	276	VAL	2.2
1	A	33	GLU	2.2
2	B	56	ILE	2.2
2	B	202	ASN	2.2
3	F	291	VAL	2.1
3	F	297	THR	2.1
1	A	17	CYS	2.1
3	F	213	VAL	2.1
2	B	82	GLY	2.1
2	B	269	PRO	2.1
1	A	25	CYS	2.1
2	B	121	ILE	2.1
1	A	35	LEU	2.1
2	B	184	GLN	2.1
3	F	387	ILE	2.1
1	A	38	LEU	2.0
2	B	24	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NDG	C	2	14/15	0.78	0.27	52,56,58,58	0
4	NAG	C	1	14/15	0.92	0.14	45,49,51,51	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	B	2001	14/15	0.70	0.44	73,77,80,80	0
6	ACT	F	452	4/4	0.97	0.15	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.