



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 02:12 pm BST

PDB ID : 3N5I  
Title : Crystal structure of the precursor (S250A mutant) of the N-terminal beta-aminopeptidase BapA  
Authors : Merz, T.; Heck, T.; Geueke, B.; Kohler, H.-P.; Gruetter, M.G.  
Deposited on : 2010-05-25  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div>••</div> </div> </div>
1	B	373	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>•</div> </div> </div>
1	C	373	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>5%</div> <div>•</div> </div> </div>
1	D	373	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>8%</div> <div>•</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-peptidyl aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	4	0
			2651	1663	484	494	10			
1	B	360	Total	C	N	O	S	0	3	0
			2639	1655	479	495	10			
1	C	362	Total	C	N	O	S	0	2	0
			2637	1651	477	499	10			
1	D	371	Total	C	N	O	S	0	3	0
			2712	1698	492	512	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	250	ALA	SER	ENGINEERED MUTATION	UNP Q52VH2
B	250	ALA	SER	ENGINEERED MUTATION	UNP Q52VH2
C	250	ALA	SER	ENGINEERED MUTATION	UNP Q52VH2
D	250	ALA	SER	ENGINEERED MUTATION	UNP Q52VH2

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	266	Total O 266 266	0	0
4	B	278	Total O 278 278	0	0
4	C	290	Total O 290 290	0	0
4	D	290	Total O 290 290	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-peptidyl aminopeptidase




- Molecule 1: Beta-peptidyl aminopeptidase




- Molecule 1: Beta-peptidyl aminopeptidase



- Molecule 1: Beta-peptidyl aminopeptidase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.70 Å 113.70 Å 126.00 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.39 – 1.80 29.39 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.39-1.80) 99.2 (29.39-1.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.38 (at 1.80 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, $R_{free}$	0.154 , 0.180 0.152 , 0.176	Depositor DCC
$R_{free}$ test set	2634 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.1	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 46.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11812	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/2709	0.55	0/3675
1	B	0.42	1/2690 (0.0%)	0.54	0/3652
1	C	0.38	0/2688	0.55	0/3650
1	D	0.36	1/2767 (0.0%)	0.54	0/3758
All	All	0.38	2/10854 (0.0%)	0.54	0/14735

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	233	ASN	C-N	-5.80	1.22	1.33
1	B	198	TYR	CD1-CE1	-5.40	1.31	1.39

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2651	0	2687	26	0
1	B	2639	0	2671	28	0
1	C	2637	0	2658	26	0
1	D	2712	0	2730	31	0
2	A	12	0	16	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	6	0	8	0	0
2	C	6	0	8	0	0
3	B	5	0	0	0	0
3	D	20	0	0	0	0
4	A	266	0	0	2	0
4	B	278	0	0	1	0
4	C	290	0	0	1	0
4	D	290	0	0	6	0
All	All	11812	0	10778	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:268:ARG:HH21	1:D:74:ASN:HD21	1.14	0.95
1:C:74:ASN:HD21	1:D:268:ARG:HH21	1.24	0.86
1:A:268:ARG:HH21	1:B:74:ASN:HD21	1.24	0.84
1:A:74:ASN:HD21	1:B:268:ARG:HH21	1.22	0.84
1:B:248:LYS:HB2	1:D:124[A]:PHE:CE1	2.16	0.81
1:B:248:LYS:HB2	1:D:124[A]:PHE:HE1	1.46	0.80
1:C:248:LYS:HE3	4:C:697:HOH:O	1.85	0.75
1:C:326:ARG:HG3	1:C:326:ARG:HH11	1.56	0.71
1:A:124[A]:PHE:CZ	1:C:248:LYS:HB2	2.25	0.71
1:A:233:ASN:ND2	1:A:351:ASN:HD22	1.90	0.69
1:A:226:LYS:NZ	1:A:364[A]:ARG:HH12	1.93	0.66
1:A:364[B]:ARG:HG2	1:A:364[B]:ARG:HH11	1.62	0.64
1:D:120:GLU:CD	1:D:120:GLU:H	2.02	0.63
1:C:326:ARG:NH1	1:D:283:THR:HG21	2.14	0.62
1:A:233:ASN:HD21	1:A:351:ASN:HD22	1.48	0.61
1:A:124[A]:PHE:CE1	1:C:248:LYS:HB2	2.37	0.60
1:B:248:LYS:CB	1:D:124[A]:PHE:HE1	2.14	0.58
1:C:326:ARG:HH12	1:D:283:THR:CG2	2.18	0.55
1:C:326:ARG:NH1	1:D:283:THR:CG2	2.70	0.55
1:B:122:ALA:HA	1:B:124:PHE:CE2	2.42	0.54
1:C:149:ARG:HG2	1:C:153:PHE:CE1	2.42	0.54
1:D:208:HIS:HA	1:D:249:ASN:HB3	1.90	0.54
1:A:74:ASN:ND2	1:B:272:ARG:HH12	2.06	0.53
1:C:74:ASN:ND2	1:D:272:ARG:HH12	2.07	0.53
1:B:215[A]:ARG:HD2	1:B:220:GLN:HG3	1.90	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124[A]:PHE:HZ	1:C:248:LYS:HB2	1.71	0.53
1:A:120:GLU:HA	1:A:123:LEU:HG	1.92	0.52
1:A:160:LYS:HE2	4:A:509:HOH:O	2.09	0.52
1:A:121:GLU:HG3	4:A:739:HOH:O	2.10	0.51
1:C:149:ARG:HG2	1:C:153:PHE:CZ	2.45	0.51
1:B:185:GLY:O	1:B:203[A]:LEU:HD22	2.11	0.51
1:C:326:ARG:HH12	1:D:283:THR:HG23	1.75	0.50
1:C:326:ARG:HG3	1:C:326:ARG:NH1	2.21	0.49
1:A:272:ARG:HH12	1:B:74:ASN:ND2	2.11	0.49
1:D:211:ARG:HD3	1:D:229:TRP:CZ3	2.47	0.49
1:B:208:HIS:HA	1:B:249:ASN:HB3	1.94	0.49
1:B:263:PRO:HG2	1:C:264:HIS:CE1	2.49	0.48
1:A:364[B]:ARG:NH1	1:A:364[B]:ARG:HG2	2.27	0.48
1:D:306:LYS:HG2	4:D:915:HOH:O	2.15	0.47
1:B:226:LYS:HG2	1:B:361:GLN:OE1	2.15	0.46
1:C:79:TRP:CD2	1:C:132:ALA:HB2	2.51	0.46
1:A:304:GLY:HA2	4:D:737:HOH:O	2.16	0.46
1:D:364:ARG:NH2	4:D:701:HOH:O	2.48	0.46
1:B:215[A]:ARG:CD	1:B:220:GLN:HG3	2.45	0.46
1:D:149:ARG:HD2	1:D:153:PHE:CZ	2.50	0.46
1:B:79:TRP:CD2	1:B:132:ALA:HB2	2.51	0.46
1:B:301:ILE:N	1:B:301:ILE:HD12	2.32	0.45
1:A:208:HIS:HA	1:A:249:ASN:HB3	1.98	0.45
1:D:249:ASN:O	1:D:288:SER:HA	2.17	0.45
1:D:92:LEU:HD22	1:D:92:LEU:N	2.31	0.45
1:D:64:ASP:OD1	1:D:302:PRO:HA	2.16	0.45
1:C:74:ASN:HB3	1:C:288:SER:O	2.17	0.45
1:D:136:ASP:O	1:D:140:ASN:HB3	2.17	0.44
1:C:265:GLN:HG2	1:C:315:ASP:OD1	2.17	0.44
1:B:186:THR:HB	1:B:203[A]:LEU:CD2	2.48	0.44
1:A:79:TRP:CG	1:A:132:ALA:HB2	2.52	0.44
1:A:79:TRP:CD2	1:A:132:ALA:HB2	2.53	0.44
1:A:6:ASP:N	1:A:6:ASP:OD1	2.51	0.44
1:B:264:HIS:CE1	1:C:263:PRO:HG2	2.53	0.43
1:B:113:TRP:O	1:B:117:LYS:HB2	2.19	0.43
1:B:213:ASP:O	1:B:215[A]:ARG:CD	2.66	0.43
1:C:193:ALA:HB1	1:C:320:THR:HG23	2.01	0.43
1:C:260:PRO:HG3	1:C:311:ALA:HB3	2.00	0.42
1:A:226:LYS:HZ2	1:A:364[A]:ARG:HH12	1.64	0.42
1:C:79:TRP:CG	1:C:132:ALA:HB2	2.54	0.42
1:D:369[A]:ARG:NH1	4:D:1108:HOH:O	2.52	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:ASP:O	1:A:140:ASN:HB3	2.20	0.42
1:B:213:ASP:O	1:B:215[A]:ARG:HD3	2.20	0.42
1:B:248:LYS:HE3	4:B:645:HOH:O	2.20	0.42
1:D:264:HIS:O	1:D:268:ARG:HG3	2.19	0.42
1:C:272:ARG:HH12	1:D:74:ASN:ND2	2.18	0.42
1:C:224:GLU:HG3	1:C:369:ARG:HH12	1.84	0.41
1:B:124:PHE:HB2	1:D:248:LYS:HD3	2.02	0.41
1:D:234:GLY:HA2	4:D:666:HOH:O	2.19	0.41
1:A:284:ALA:HB3	1:B:272:ARG:NE	2.36	0.41
1:B:186:THR:HB	1:B:203[A]:LEU:HD23	2.01	0.41
1:D:246:GLN:NE2	4:D:856:HOH:O	2.47	0.41
1:A:135:LEU:HD23	1:A:248:LYS:HB3	2.02	0.40
1:B:248:LYS:HB2	1:D:124[A]:PHE:CZ	2.52	0.40
1:D:79:TRP:CD2	1:D:132:ALA:HB2	2.56	0.40
1:A:249:ASN:O	1:A:288:SER:HA	2.21	0.40
1:A:318:SER:O	1:A:322:ASN:HB2	2.22	0.40
1:B:138:ARG:HH12	1:B:230:PRO:CB	2.34	0.40
1:C:264:HIS:O	1:C:268:ARG:HG3	2.22	0.40
1:D:123:LEU:HA	1:D:123:LEU:HD23	1.97	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/373 (96%)	355 (99%)	5 (1%)	0	100	100
1	B	359/373 (96%)	352 (98%)	7 (2%)	0	100	100
1	C	360/373 (96%)	355 (99%)	5 (1%)	0	100	100
1	D	372/373 (100%)	367 (99%)	5 (1%)	0	100	100
All	All	1451/1492 (97%)	1429 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/276 (99%)	270 (99%)	2 (1%)	84	81
1	B	271/276 (98%)	271 (100%)	0	100	100
1	C	270/276 (98%)	270 (100%)	0	100	100
1	D	277/276 (100%)	274 (99%)	3 (1%)	73	68
All	All	1090/1104 (99%)	1085 (100%)	5 (0%)	91	87

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASP
1	A	233	ASN
1	D	179	THR
1	D	369[A]	ARG
1	D	369[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	233	ASN
1	B	74	ASN
1	B	108	GLN
1	B	220	GLN
1	C	74	ASN
1	C	90	GLN
1	C	220	GLN
1	C	233	ASN
1	D	74	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	D	374	-	4,4,4	0.19	0	6,6,6	0.21	0
2	GOL	A	375	-	5,5,5	0.37	0	5,5,5	0.60	0
3	SO4	D	377	-	4,4,4	0.15	0	6,6,6	0.08	0
3	SO4	D	375	-	4,4,4	0.14	0	6,6,6	0.09	0
3	SO4	B	375	-	4,4,4	0.15	0	6,6,6	0.09	0
2	GOL	B	374	-	5,5,5	0.36	0	5,5,5	0.46	0
2	GOL	A	374	-	5,5,5	0.36	0	5,5,5	0.73	0
2	GOL	C	374	-	5,5,5	0.39	0	5,5,5	0.54	0
3	SO4	D	376	-	4,4,4	0.13	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	375	-	-	0/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	374	-	-	0/4/4/4	-
2	GOL	A	374	-	-	0/4/4/4	-
2	GOL	C	374	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	360/373 (96%)	-0.41	6 (1%) 70 66	12, 18, 35, 52	0
1	B	360/373 (96%)	-0.33	5 (1%) 75 72	14, 20, 37, 63	0
1	C	362/373 (97%)	-0.44	4 (1%) 80 78	13, 19, 35, 52	0
1	D	371/373 (99%)	-0.40	3 (0%) 86 84	13, 19, 35, 55	2 (0%)
All	All	1453/1492 (97%)	-0.40	18 (1%) 79 76	12, 19, 36, 63	2 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	195	ASP	5.3
1	B	116	GLY	4.4
1	A	235	ILE	4.4
1	B	194	GLY	4.1
1	B	195	ASP	4.1
1	C	371	PRO	4.0
1	A	234	GLY	3.5
1	A	371	PRO	3.5
1	D	371	PRO	3.1
1	A	233	ASN	3.0
1	A	195	ASP	2.8
1	C	124	PHE	2.6
1	C	195	ASP	2.5
1	D	254	VAL	2.5
1	C	194	GLY	2.5
1	B	124	PHE	2.5
1	B	121	GLU	2.3
1	A	232	VAL	2.2



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	D	375	5/5	0.86	0.24	35,42,60,60	5
2	GOL	B	374	6/6	0.93	0.11	25,27,28,35	0
2	GOL	A	374	6/6	0.93	0.12	23,25,26,26	0
2	GOL	C	374	6/6	0.93	0.10	24,26,27,29	0
2	GOL	A	375	6/6	0.94	0.10	23,25,26,29	0
3	SO4	B	375	5/5	0.96	0.27	46,47,56,60	0
3	SO4	D	377	5/5	0.97	0.22	45,51,53,55	0
3	SO4	D	374	5/5	0.97	0.17	34,34,43,44	0
3	SO4	D	376	5/5	0.97	0.19	37,43,51,51	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.