



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 12:46 am BST

PDB ID : 5N7G  
Title : MAGI-1 complexed with a synthetic pRSK1 peptide  
Authors : Gogl, G.; Nyitray, L.  
Deposited on : 2017-02-20  
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	
1	B	427	
2	C	7	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5939 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1,Annexin A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	423	Total	C	N	O	S	0	0	0
			3334	2100	573	647	14			
1	B	318	Total	C	N	O	S	0	0	0
			2521	1584	431	495	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	452	GLY	-	expression tag	UNP Q96QZ7
A	453	SER	-	expression tag	UNP Q96QZ7
A	454	MET	-	expression tag	UNP Q96QZ7
A	559	GLY	-	linker	UNP Q96QZ7
A	560	SER	-	linker	UNP Q96QZ7
A	605	GLU	ALA	conflict	UNP P07355
B	452	GLY	-	expression tag	UNP Q96QZ7
B	453	SER	-	expression tag	UNP Q96QZ7
B	454	MET	-	expression tag	UNP Q96QZ7
B	559	GLY	-	linker	UNP Q96QZ7
B	560	SER	-	linker	UNP Q96QZ7
B	605	GLU	ALA	conflict	UNP P07355

- Molecule 2 is a protein called Ribosomal protein S6 kinase alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	P	0	0	0
			45	25	6	13	1			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Ca	0	0
			3	3		
4	A	6	Total	Ca	0	0
			6	6		



- Molecule 1: Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1,Annexin A2

GLY	D861	T862	Q867	D878	F482	T495	D490	E491	E494	H532	L558	G559	S560	L579	V590	I598	R602	R607	L641	N676	T702	R707	R718	I725	K766	I783	R784	R785	N793	A794	P795	L796	I797	S816	R823
-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

- Molecule 1: Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1,Annexin A2

Chain B:

Amino Acid	Frequency
GLY	67%
SER	7%
MET	26%
GLU	
THR	
GLY	
LYS	
PRO	
PHE	
THR	
ARG	
ASN	
PRO	
SER	
LEU	
LYS	
LYS	
PHE	
ILE	
THR	
LYS	
LEU	
ARG	
ARG	
SER	
GLY	
PHE	
GLY	
PHE	
THR	
VAL	
VAL	
GLY	
ASP	
GLU	
PRO	
ASP	
GLU	
PHE	
LEU	
GLN	
ILE	
LYS	
SER	
VAL	
LEU	
ASP	
LEU	
THR	
MET	
THR	
LYS	
MET	
GLU	
THR	
GLY	
ASP	
VAL	
ILE	
VAL	
SER	
VAL	
ASN	
ASP	
THR	
CYS	
VAL	
LEU	
GLY	
HIS	
THR	
HIS	
ALA	
GLN	
VAL	
VAL	
ILE	
PHE	
GLN	
SER	
ILE	
PRO	
ILE	
GLY	
ALA	
SER	
VAL	
ASP	
LEU	
LEU	
CYS	
ARG	
GLY	
THR	
PRO	
LEU	
GLY	
SER	
S61	
I58	
R62	
L36	
L42	
K63	
T84	

- Molecule 2: Ribosomal protein S6 kinase alpha-1

Chain C:

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	194.93 Å   60.23 Å   99.35 Å 90.00°   98.69°   90.00°	Depositor
Resolution (Å)	49.10 – 2.95 49.10 – 2.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.10-2.95) 98.8 (49.10-2.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 2.96 Å)	Xtriage
Refinement program	PHENIX (dev_2420: ???)	Depositor
R, $R_{free}$	0.210   ,   0.264 0.210   ,   0.264	Depositor DCC
$R_{free}$ test set	1195 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.1	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 23.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	5939	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/3385	0.41	0/4554
1	B	0.23	0/2555	0.39	0/3436
2	C	0.34	0/34	0.65	0/44
All	All	0.29	0/5974	0.40	0/8034

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3334	0	3346	16	0
1	B	2521	0	2511	20	0
2	C	45	0	36	3	0
3	A	18	0	24	0	0
3	B	12	0	16	0	0
4	A	6	0	0	0	0
4	B	3	0	0	0	0
All	All	5939	0	5933	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:643:LYS:HB3	1:B:647:GLN:HG3	1.72	0.71
1:B:783:ILE:HD13	1:B:795:PHE:HB3	1.79	0.64
1:A:532:HIS:NE2	2:C:733:THR:OG1	2.31	0.63
1:A:558:LEU:O	1:A:560:SER:N	2.32	0.63
1:B:679:LEU:O	1:B:683:ASN:ND2	2.31	0.61
1:A:491:GLU:HB3	1:A:494:GLU:HB2	1.85	0.59
1:B:793:ASN:O	1:B:797:ASN:ND2	2.34	0.58
1:B:709:LEU:HD13	1:B:753:ILE:HG13	1.86	0.58
1:A:456:LYS:N	1:A:457:PRO:HD3	2.18	0.58
1:B:626:LEU:HD12	1:B:642:LEU:HD11	1.84	0.58
1:B:698:ILE:O	1:B:702:THR:OG1	2.19	0.57
1:A:783:ILE:HD13	1:A:795:PHE:HB3	1.86	0.55
1:B:674:ARG:HB3	1:B:679:LEU:HG	1.89	0.53
1:B:693:ASP:HB2	1:B:696:LYS:HG3	1.90	0.53
1:A:485:THR:HG22	2:C:734:THR:HA	1.91	0.53
1:B:822:THR:OG1	1:B:861:ASP:OD2	2.27	0.52
1:B:728:GLU:OE1	1:B:728:GLU:N	2.44	0.51
1:A:793:ASN:O	1:A:797:ASN:ND2	2.40	0.49
1:B:656:SER:O	1:B:661:GLY:HA2	2.13	0.48
1:B:787:VAL:HG21	1:B:795:PHE:HD1	1.79	0.48
1:A:702:THR:O	1:A:707:ARG:NH2	2.44	0.48
1:A:725:ILE:HD13	1:A:766:LYS:HE2	1.96	0.47
1:B:661:GLY:C	1:B:663:ASP:H	2.19	0.46
1:B:665:ASP:HB3	1:B:812:ARG:HH22	1.79	0.46
1:A:816:SER:HB2	1:A:823:ARG:HB2	1.99	0.45
1:B:598:ILE:O	1:B:602:ARG:HG2	2.16	0.45
1:B:644:THR:OG1	1:B:647:GLN:HG2	2.18	0.44
1:B:643:LYS:HB2	1:B:648:TYR:HB2	1.99	0.44
1:A:482:PHE:N	2:C:735:LEU:OXT	2.50	0.43
1:A:607:ARG:NH1	1:A:641:LEU:O	2.51	0.43
1:A:862:THR:O	1:A:867:GLN:NE2	2.42	0.42
1:A:676:ASN:ND2	1:A:718:ARG:O	2.48	0.42
1:B:787:VAL:O	1:B:788:LYS:HD2	2.20	0.42
1:A:598:ILE:O	1:A:602:ARG:HG2	2.20	0.41
1:B:740:ALA:HB1	1:B:748:ASP:HB3	2.03	0.41
1:A:785:LYS:HE2	1:A:785:LYS:HB3	1.90	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	421/427 (99%)	414 (98%)	6 (1%)	1 (0%)	47	79
1	B	316/427 (74%)	309 (98%)	7 (2%)	0	100	100
2	C	3/7 (43%)	3 (100%)	0	0	100	100
All	All	740/861 (86%)	726 (98%)	13 (2%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	590	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/370 (98%)	358 (99%)	5 (1%)	67	86
1	B	270/370 (73%)	269 (100%)	1 (0%)	91	96
2	C	4/6 (67%)	4 (100%)	0	100	100
All	All	637/746 (85%)	631 (99%)	6 (1%)	78	91

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	456	LYS
1	A	490	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	558	LEU
1	A	579	LEU
1	A	861	ASP
1	B	861	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SEP	C	732	2	8,9,10	1.64	1 (12%)	8,12,14	2.37	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	C	732	2	-	4/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	732	SEP	P-O1P	3.49	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	C	732	SEP	P-OG-CB	-4.91	104.78	118.30
2	C	732	SEP	OG-CB-CA	4.16	112.20	108.14

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	732	SEP	N-CA-CB-OG
2	C	732	SEP	CB-OG-P-O2P
2	C	732	SEP	CB-OG-P-O3P
2	C	732	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 9 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	902	-	5,5,5	0.37	0	5,5,5	0.31	0
3	GOL	A	901	-	5,5,5	0.39	0	5,5,5	0.35	0
3	GOL	A	903	-	5,5,5	0.35	0	5,5,5	0.27	0
3	GOL	B	901	-	5,5,5	0.37	0	5,5,5	0.26	0
3	GOL	B	902	-	5,5,5	0.37	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	902	-	-	2/4/4/4	-
3	GOL	A	901	-	-	2/4/4/4	-
3	GOL	A	903	-	-	2/4/4/4	-
3	GOL	B	901	-	-	2/4/4/4	-
3	GOL	B	902	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	GOL	O1-C1-C2-C3
3	A	902	GOL	O1-C1-C2-C3
3	B	902	GOL	O1-C1-C2-C3
3	B	901	GOL	O1-C1-C2-C3
3	A	903	GOL	O1-C1-C2-C3
3	B	902	GOL	O1-C1-C2-O2
3	B	901	GOL	O1-C1-C2-O2
3	A	903	GOL	O1-C1-C2-O2
3	A	901	GOL	O1-C1-C2-O2
3	A	902	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	423/427 (99%)	-0.38	0 <span>100</span> <span>100</span>	10, 29, 54, 102	0
1	B	318/427 (74%)	0.13	1 (0%) <span>94</span> <span>87</span>	34, 60, 98, 117	0
2	C	5/7 (71%)	1.08	1 (20%) <span>1</span> <span>0</span>	89, 92, 105, 116	0
All	All	746/861 (86%)	-0.15	2 (0%) <span>94</span> <span>87</span>	10, 42, 89, 117	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	730	LEU	3.0
1	B	717	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SEP	C	732	10/11	0.80	0.22	111,126,133,138	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CA	B	904	1/1	0.87	0.18	90,90,90,90	0
4	CA	B	903	1/1	0.88	0.09	89,89,89,89	0
3	GOL	B	902	6/6	0.90	0.18	54,66,71,81	0
4	CA	A	904	1/1	0.93	0.21	44,44,44,44	0
4	CA	A	908	1/1	0.93	0.14	76,76,76,76	0
3	GOL	B	901	6/6	0.94	0.21	20,41,49,50	0
3	GOL	A	903	6/6	0.95	0.21	26,27,32,34	0
3	GOL	A	901	6/6	0.96	0.19	12,17,18,20	0
3	GOL	A	902	6/6	0.96	0.15	9,32,39,40	0
4	CA	B	905	1/1	0.96	0.10	38,38,38,38	0
4	CA	A	907	1/1	0.97	0.11	49,49,49,49	0
4	CA	A	909	1/1	0.98	0.10	34,34,34,34	0
4	CA	A	905	1/1	0.98	0.06	34,34,34,34	0
4	CA	A	906	1/1	0.98	0.10	35,35,35,35	0

## 6.5 Other polymers

There are no such residues in this entry.