



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 30, 2021 – 02:04 PM EDT

PDB ID : 7N79
Title : O2-, PLP-dependent desaturase Plu4 holo-enzyme
Authors : Hoffarth, E.R.; Ryan, K.S.
Deposited on : 2021-06-09
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>• •</div> </div> </div>
1	B	389	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	403	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aminotran_1_2 domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	P	S	0	4	0
			3018	1938	486	582	1	11			
1	B	378	Total	C	N	O	P	S	0	2	0
			3011	1926	490	584	1	10			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	382	LEU	-	expression tag	UNP A0A0C1MLE8
A	383	GLU	-	expression tag	UNP A0A0C1MLE8
A	384	HIS	-	expression tag	UNP A0A0C1MLE8
A	385	HIS	-	expression tag	UNP A0A0C1MLE8
A	386	HIS	-	expression tag	UNP A0A0C1MLE8
A	387	HIS	-	expression tag	UNP A0A0C1MLE8
A	388	HIS	-	expression tag	UNP A0A0C1MLE8
A	389	HIS	-	expression tag	UNP A0A0C1MLE8
B	382	LEU	-	expression tag	UNP A0A0C1MLE8
B	383	GLU	-	expression tag	UNP A0A0C1MLE8
B	384	HIS	-	expression tag	UNP A0A0C1MLE8
B	385	HIS	-	expression tag	UNP A0A0C1MLE8
B	386	HIS	-	expression tag	UNP A0A0C1MLE8
B	387	HIS	-	expression tag	UNP A0A0C1MLE8
B	388	HIS	-	expression tag	UNP A0A0C1MLE8
B	389	HIS	-	expression tag	UNP A0A0C1MLE8

- Molecule 2 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	143	Total	O	0	0
			143	143		
5	B	187	Total	O	0	0
			187	187		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aminotran_1_2 domain-containing protein



- Molecule 1: Aminotran_1_2 domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.02Å 72.19Å 76.27Å 90.00° 112.09° 90.00°	Depositor
Resolution (Å)	37.75 – 1.98 37.75 – 1.98	Depositor EDS
% Data completeness (in resolution range)	98.4 (37.75-1.98) 98.4 (37.75-1.98)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.166 , 0.215 0.166 , 0.215	Depositor DCC
R_{free} test set	2642 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6411	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BTB, EDO, ACT, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/3070	0.68	5/4167 (0.1%)
1	B	0.51	0/3057	0.64	6/4153 (0.1%)
All	All	0.53	0/6127	0.66	11/8320 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	73[A]	SER	CA-C-O	6.63	134.01	120.10
1	B	73[B]	SER	CA-C-O	6.63	134.01	120.10
1	B	73[A]	SER	CA-C-N	-5.62	104.84	117.20
1	B	73[B]	SER	CA-C-N	-5.62	104.84	117.20
1	A	238[A]	SER	CA-C-O	5.58	131.82	120.10
1	A	238[B]	SER	CA-C-O	5.58	131.82	120.10
1	B	239[A]	SER	CA-C-O	5.58	131.81	120.10
1	B	239[B]	SER	CA-C-O	5.58	131.81	120.10
1	A	358[A]	MET	CA-C-O	5.36	131.36	120.10
1	A	358[B]	MET	CA-C-O	5.36	131.36	120.10
1	A	302	ASP	N-CA-C	5.17	124.94	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3018	0	2878	42	0
1	B	3011	0	2866	25	0
2	A	14	0	18	6	0
2	B	14	0	19	3	0
3	A	12	0	9	4	0
4	B	12	0	18	3	0
5	A	143	0	0	9	0
5	B	187	0	0	4	0
All	All	6411	0	5808	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:ALA:O	1:B:372:LYS:HD3	1.44	1.13
1:B:290:ALA:O	1:B:372:LYS:CD	2.13	0.96
1:A:26:GLY:O	1:A:358[A]:MET:HE3	1.65	0.94
1:A:26:GLY:O	1:A:358[A]:MET:CE	2.16	0.92
1:B:68:GLU:OE1	5:B:501:HOH:O	1.94	0.86
1:B:145:SER:HA	1:B:180:HIS:CD2	2.12	0.84
1:B:226:LEU:HD12	4:B:402:EDO:H21	1.66	0.78
1:A:152:LEU:HG	1:A:154:LEU:HD21	1.68	0.75
1:B:138:HIS:O	1:B:142:GLN:HG3	1.91	0.70
1:A:290:ALA:O	1:A:372:LYS:HD2	1.93	0.67
1:A:26:GLY:O	1:A:358[A]:MET:HE2	1.95	0.67
1:A:59:GLU:OE2	2:A:401:BTB:H12	1.97	0.64
1:A:140:LEU:HD23	1:A:140:LEU:O	2.01	0.61
3:A:403:ACT:H3	5:B:591:HOH:O	2.02	0.59
1:B:87:ALA:HB3	1:B:219:LLP:H5'1	1.83	0.59
2:A:401:BTB:H52	5:A:505:HOH:O	2.02	0.59
1:A:331:HIS:O	1:A:332:ASP:HB2	2.01	0.58
1:B:85:PRO:HB3	4:B:402:EDO:H12	1.84	0.58
1:A:308:PRO:HB2	1:A:358[B]:MET:HG2	1.87	0.56
1:B:56:PRO:O	1:B:60:LYS:HG3	2.05	0.56
1:A:137:GLU:HB3	1:A:172:TYR:CZ	2.41	0.56
1:A:375:VAL:O	1:A:379:LEU:HG	2.05	0.55
1:B:226:LEU:CD1	4:B:402:EDO:H21	2.35	0.54
1:B:152:LEU:HG	1:B:154:LEU:HD21	1.90	0.52
1:A:72:SER:HA	1:A:199:ASP:OD2	2.10	0.52
1:B:144:VAL:HG23	1:B:182:ILE:HG13	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145:SER:HA	1:B:180:HIS:HD2	1.72	0.52
2:A:401:BTB:C5	5:A:505:HOH:O	2.57	0.52
1:A:60:LYS:NZ	5:A:505:HOH:O	2.42	0.52
1:A:296:LEU:N	1:A:296:LEU:HD23	2.26	0.51
5:A:584:HOH:O	1:B:51:LYS:HE2	2.11	0.51
1:A:134:LEU:HB3	1:A:165:MET:HE3	1.92	0.50
2:A:401:BTB:H52	2:A:401:BTB:H11	1.44	0.49
1:A:116:TYR:OH	3:A:404:ACT:H1	2.13	0.49
1:A:134:LEU:HD22	1:A:165:MET:HE1	1.95	0.48
1:A:5:ARG:HA	5:A:637:HOH:O	2.12	0.48
1:A:307:LEU:C	1:A:309[B]:LEU:H	2.17	0.47
1:A:48:GLN:NE2	5:A:502:HOH:O	1.90	0.47
1:B:290:ALA:O	1:B:372:LYS:NZ	2.43	0.47
1:A:27:HIS:CD2	1:A:219:LLP:HE2	2.50	0.47
1:B:300:ASP:OD2	1:B:301:ASN:N	2.48	0.47
2:B:404:BTB:H32	2:B:404:BTB:H52	1.48	0.46
1:A:135:LYS:HG2	1:A:169:GLU:OE1	2.16	0.46
1:A:153:PHE:HA	1:A:185:LEU:HB2	1.98	0.46
2:B:404:BTB:H11	2:B:404:BTB:H71	1.63	0.46
1:A:32:ILE:HG23	1:A:36:TYR:HB2	1.98	0.45
1:A:307:LEU:C	1:A:309[A]:LEU:H	2.18	0.45
1:B:135:LYS:HE2	1:B:169:GLU:OE2	2.16	0.45
1:A:69:LEU:HD22	1:A:264:VAL:HG13	1.98	0.45
1:A:202:GLN:HG2	1:A:206:GLN:NE2	2.31	0.45
1:A:209:ILE:C	1:A:209:ILE:CD1	2.84	0.45
3:A:403:ACT:H2	5:B:527:HOH:O	2.17	0.45
1:B:356:SER:OG	1:B:359:LYS:HE3	2.17	0.44
2:A:401:BTB:H62	2:A:401:BTB:H71	1.37	0.44
1:B:193:TYR:HE1	1:B:280:VAL:HG21	1.83	0.44
1:A:87:ALA:HB3	1:A:219:LLP:H5'1	1.99	0.44
1:A:304:LYS:HE2	1:A:305:CYS:O	2.18	0.43
1:B:371:LEU:O	1:B:375:VAL:HG23	2.18	0.43
1:B:273:ASP:HA	1:B:277:LYS:HB2	2.01	0.43
1:B:308:PRO:HB2	1:B:358:MET:HG2	2.01	0.43
3:A:403:ACT:H2	5:A:636:HOH:O	2.19	0.42
1:A:63:LYS:HB3	1:A:63:LYS:HE3	1.80	0.42
2:B:404:BTB:H61	5:B:624:HOH:O	2.20	0.42
1:A:137:GLU:HG2	5:A:629:HOH:O	2.19	0.42
1:B:69:LEU:HD22	1:B:264:VAL:HG13	2.02	0.42
1:B:106:ALA:HB2	1:B:149:ILE:HG21	2.02	0.42
1:A:226:LEU:HD21	1:A:256:ALA:HA	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:BTB:H32	2:A:401:BTB:H72	1.24	0.41
1:A:248:ILE:HG23	1:A:249:PHE:CD2	2.55	0.41
1:B:286:THR:HG22	1:B:368:ILE:HG21	2.01	0.41
1:A:140:LEU:C	1:A:140:LEU:CD2	2.89	0.41
1:A:372:LYS:O	1:A:376:THR:HG23	2.20	0.41
1:A:308:PRO:HB2	1:A:358[A]:MET:HG2	2.02	0.41
1:A:129:PHE:CE2	1:A:152:LEU:HD11	2.56	0.41
1:A:274:ALA:HA	5:A:501:HOH:O	2.21	0.41
1:A:156:ASN:O	1:A:188:ARG:HD2	2.21	0.40
1:A:300:ASP:OD1	1:A:301:ASN:N	2.55	0.40
1:A:178:LYS:HA	1:A:209:ILE:CG2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/389 (97%)	366 (97%)	13 (3%)	0	100	100
1	B	377/389 (97%)	369 (98%)	8 (2%)	0	100	100
All	All	756/778 (97%)	735 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/344 (91%)	304 (97%)	10 (3%)	39	28
1	B	316/344 (92%)	311 (98%)	5 (2%)	62	56
All	All	630/688 (92%)	615 (98%)	15 (2%)	49	41

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASP
1	A	5	ARG
1	A	25	ASP
1	A	77	SER
1	A	140	LEU
1	A	153	PHE
1	A	209	ILE
1	A	243	LEU
1	A	282	ARG
1	A	304	LYS
1	B	79	ASN
1	B	129	PHE
1	B	153	PHE
1	B	377	ARG
1	B	379	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	GLN
1	A	331	HIS
1	B	331	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	B	219	1	23,24,25	2.50	6 (26%)	25,32,34	1.40	4 (16%)
1	LLP	A	219	1	23,24,25	2.48	6 (26%)	25,32,34	1.49	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	219	1	-	4/16/17/19	0/1/1/1
1	LLP	A	219	1	-	6/16/17/19	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	219	LLP	C4-C4'	8.12	1.62	1.46
1	A	219	LLP	C4-C4'	7.74	1.61	1.46
1	B	219	LLP	C4'-NZ	4.66	1.42	1.27
1	A	219	LLP	C4'-NZ	4.58	1.42	1.27
1	A	219	LLP	C4-C5	-3.95	1.37	1.42
1	A	219	LLP	C2'-C2	3.39	1.56	1.50
1	B	219	LLP	C2'-C2	3.27	1.55	1.50
1	B	219	LLP	C4-C5	-2.93	1.38	1.42
1	A	219	LLP	C6-N1	2.85	1.40	1.34
1	B	219	LLP	C6-N1	2.79	1.40	1.34
1	B	219	LLP	C5'-C5	2.45	1.57	1.50
1	A	219	LLP	C5'-C5	2.16	1.56	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	LLP	CE-NZ-C4'	-4.08	106.39	118.90
1	B	219	LLP	OP4-P-OP1	3.06	115.06	106.47
1	A	219	LLP	C4-C4'-NZ	-2.92	110.92	124.31
1	B	219	LLP	C4-C4'-NZ	-2.89	111.06	124.31
1	A	219	LLP	OP4-P-OP1	2.70	114.06	106.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	219	LLP	CE-NZ-C4'	-2.66	110.74	118.90
1	B	219	LLP	C5-C6-N1	-2.25	120.06	123.82
1	A	219	LLP	C3-C4-C5	2.14	119.91	118.26

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	219	LLP	C-CA-CB-CG
1	B	219	LLP	C4-C5-C5'-OP4
1	B	219	LLP	C6-C5-C5'-OP4
1	B	219	LLP	C4-C4'-NZ-CE
1	A	219	LLP	C6-C5-C5'-OP4
1	A	219	LLP	C4-C4'-NZ-CE
1	B	219	LLP	C3-C4-C4'-NZ
1	A	219	LLP	C4-C5-C5'-OP4
1	A	219	LLP	C3-C4-C4'-NZ
1	A	219	LLP	N-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	219	LLP	1	0
1	A	219	LLP	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	A	402	-	1,3,3	2.97	1 (100%)	0,3,3	-	-
2	BTB	A	401	-	13,13,13	1.47	2 (15%)	7,16,16	1.32	1 (14%)
4	EDO	B	402	-	3,3,3	0.48	0	2,2,2	0.30	0
4	EDO	B	403	-	3,3,3	0.54	0	2,2,2	0.15	0
4	EDO	B	401	-	3,3,3	0.52	0	2,2,2	0.45	0
2	BTB	B	404	-	13,13,13	0.61	0	7,16,16	0.48	0
3	ACT	A	403	-	1,3,3	2.61	1 (100%)	0,3,3	-	-
3	ACT	A	404	-	1,3,3	2.26	1 (100%)	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTB	A	401	-	-	13/21/21/21	-
2	BTB	B	404	-	-	5/21/21/21	-
4	EDO	B	402	-	-	1/1/1/1	-
4	EDO	B	403	-	-	1/1/1/1	-
4	EDO	B	401	-	-	0/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	BTB	C7-N	-2.99	1.43	1.48
3	A	402	ACT	CH3-C	2.97	1.52	1.48
3	A	403	ACT	CH3-C	2.61	1.52	1.48
3	A	404	ACT	CH3-C	2.26	1.51	1.48
2	A	401	BTB	C4-C2	-2.07	1.50	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	BTB	O4-C4-C2	-2.53	104.52	111.44

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	BTB	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401	BTB	O1-C1-C2-C4
2	A	401	BTB	O1-C1-C2-N
2	A	401	BTB	C1-C2-C3-O3
2	A	401	BTB	C4-C2-C3-O3
2	A	401	BTB	N-C2-C3-O3
2	A	401	BTB	C1-C2-N-C7
2	A	401	BTB	C3-C2-N-C7
2	A	401	BTB	C4-C2-N-C7
2	A	401	BTB	C6-C5-N-C7
2	B	404	BTB	N-C7-C8-O8
4	B	403	EDO	O1-C1-C2-O2
2	A	401	BTB	C1-C2-N-C5
2	A	401	BTB	C4-C2-N-C5
2	B	404	BTB	C1-C2-N-C5
2	B	404	BTB	C1-C2-N-C7
2	B	404	BTB	C3-C2-N-C5
2	B	404	BTB	C4-C2-N-C7
2	A	401	BTB	N-C5-C6-O6
4	B	402	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	BTB	6	0
4	B	402	EDO	3	0
2	B	404	BTB	3	0
3	A	403	ACT	3	0
3	A	404	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/389 (96%)	0.04	15 (3%) 38 40	30, 53, 79, 105	0
1	B	377/389 (96%)	-0.13	15 (3%) 38 40	29, 45, 71, 98	0
All	All	754/778 (96%)	-0.05	30 (3%) 38 40	29, 49, 76, 105	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	379	LEU	3.8
1	A	248	ILE	3.7
1	B	4	ASP	3.3
1	A	249	PHE	3.2
1	A	86	THR	3.1
1	A	84	CYS	3.0
1	B	85	PRO	3.0
1	B	86	THR	3.0
1	B	228	ILE	3.0
1	B	318	GLY	2.7
1	B	89	ASN	2.7
1	B	248	ILE	2.6
1	A	85	PRO	2.5
1	B	88	SER	2.5
1	A	250	LEU	2.5
1	B	227	LYS	2.5
1	A	319	TYR	2.5
1	A	87	ALA	2.4
1	B	218	GLY	2.3
1	A	228	ILE	2.3
1	A	251	CYS	2.3
1	A	4	ASP	2.3
1	B	319	TYR	2.2
1	B	249	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	378	ILE	2.2
1	B	226	LEU	2.2
1	A	88	SER	2.1
1	A	295	GLY	2.1
1	A	381	LYS	2.1
1	A	227	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	219	24/25	0.96	0.21	34,44,61,63	0
1	LLP	B	219	24/25	0.98	0.18	28,39,48,51	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BTB	B	404	14/14	0.77	0.21	62,85,95,96	0
3	ACT	A	404	4/4	0.77	0.21	102,108,110,111	0
3	ACT	A	402	4/4	0.83	0.31	72,83,83,84	0
4	EDO	B	403	4/4	0.84	0.12	54,55,59,64	0
2	BTB	A	401	14/14	0.86	0.20	60,77,102,103	0
3	ACT	A	403	4/4	0.88	0.17	98,101,102,102	0
4	EDO	B	401	4/4	0.95	0.09	50,51,51,55	0
4	EDO	B	402	4/4	0.96	0.16	35,44,46,46	0

6.5 Other polymers ⓘ

There are no such residues in this entry.