



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2021 – 10:36 AM EDT

PDB ID : 7N7B  
Title : crystal structure of the N-formyltransferase HCAN\_0200 from *Helicobacter canadensis* on complex with folinic acid and dTDP-3-aminofucose  
Authors : Heisdorf, C.J.; Thoden, J.B.; Holden, H.M.  
Deposited on : 2021-06-10  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	 81% 14% . .
1	B	280	 2% 86% 11% .

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5104 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Formyl\_trans\_N domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	0	0	0
			2235	1457	363	409	6			
1	B	271	Total	C	N	O	S	0	0	0
			2245	1463	365	411	6			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	273	LEU	-	expression tag	UNP C5ZW02
A	274	GLU	-	expression tag	UNP C5ZW02
A	275	HIS	-	expression tag	UNP C5ZW02
A	276	HIS	-	expression tag	UNP C5ZW02
A	277	HIS	-	expression tag	UNP C5ZW02
A	278	HIS	-	expression tag	UNP C5ZW02
A	279	HIS	-	expression tag	UNP C5ZW02
A	280	HIS	-	expression tag	UNP C5ZW02
B	273	LEU	-	expression tag	UNP C5ZW02
B	274	GLU	-	expression tag	UNP C5ZW02
B	275	HIS	-	expression tag	UNP C5ZW02
B	276	HIS	-	expression tag	UNP C5ZW02
B	277	HIS	-	expression tag	UNP C5ZW02
B	278	HIS	-	expression tag	UNP C5ZW02
B	279	HIS	-	expression tag	UNP C5ZW02
B	280	HIS	-	expression tag	UNP C5ZW02

- Molecule 2 is N-{[4-({[(6R)-2-amino-5-formyl-4-oxo-1,4,5,6,7,8-hexahydropteridin-6-yl]methyl}amino)phenyl]carbonyl}-L-glutamic acid (three-letter code: FON) (formula: C<sub>20</sub>H<sub>23</sub>N<sub>7</sub>O<sub>7</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			34	20	7	7		
2	B	1	Total	C	N	O	0	0
			34	20	7	7		

- Molecule 3 is (3R,4S,5R,6R)-4-amino-3,5-dihydroxy-6-methyloxan-2-yl][hydroxy-[[[(2R,3S,5R)-3-hydroxy-5-(5-methyl-2,4-dioxypyrimidin-1-yl)oxolan-2-yl]methoxy]phosphoryl]hydrogen phosphate (three-letter code: T3F) (formula:  $C_{16}H_{27}N_3O_{14}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			35	16	3	14	2		
3	B	1	Total	C	N	O	P	0	0
			35	16	3	14	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	255	Total	O	0	0
			255	255		
6	B	199	Total	O	0	0
			199	199		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Formyl\_trans\_N domain-containing protein



- Molecule 1: Formyl\_trans\_N domain-containing protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.26Å 74.02Å 83.00Å 90.00° 114.01° 90.00°	Depositor
Resolution (Å)	29.64 – 2.00 29.62 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.6 (29.64-2.00) 96.6 (29.62-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, $R_{free}$	0.192 , 0.253 0.197 , 0.251	Depositor DCC
$R_{free}$ test set	2330 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5104	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, FON, T3F, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	0/2288	0.83	1/3085 (0.0%)
1	B	0.67	0/2298	0.79	0/3099
All	All	0.68	0/4586	0.81	1/6184 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	ARG	NE-CZ-NH2	-5.47	117.56	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2235	0	2216	29	0
1	B	2245	0	2223	17	0
2	A	34	0	21	1	0
2	B	34	0	21	1	0
3	A	35	0	25	0	0
3	B	35	0	25	0	0
4	A	12	0	13	2	0
5	A	16	0	24	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	4	0	6	0	0
6	A	255	0	0	3	0
6	B	199	0	0	2	0
All	All	5104	0	4574	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:ASN:HD22	1:A:179:ASN:ND2	1.74	0.86
1:B:184:ASN:HD22	1:B:184:ASN:H	1.37	0.72
1:A:184:ASN:H	1:A:184:ASN:HD22	1.37	0.72
1:A:82:VAL:H	2:A:301:FON:HN1	1.45	0.64
1:A:18:GLU:OE1	4:A:303:MES:H32	1.98	0.63
1:A:135:ASN:HD22	1:A:179:ASN:HD22	1.44	0.62
1:A:184:ASN:HD22	1:A:184:ASN:N	1.98	0.61
1:B:50:HIS:ND1	6:B:402:HOH:O	2.31	0.58
1:A:122:VAL:HG23	1:A:164:LEU:HD22	1.86	0.56
1:B:14:VAL:HG11	1:B:46:LYS:HE3	1.88	0.56
1:B:20:ILE:HD11	1:B:169:ILE:HG12	1.88	0.55
1:B:128:ASP:O	2:B:301:FON:NA2	2.38	0.55
1:A:168:ASN:OD1	1:A:171:ARG:NH1	2.41	0.54
1:B:140:TYR:CE2	1:B:160:TYR:CE1	2.98	0.52
1:A:115:ASN:O	5:A:305:EDO:H22	2.10	0.52
1:A:5:CYS:HB2	1:A:71:ILE:HD11	1.92	0.51
1:B:111:THR:OG1	1:B:112:PRO:HD3	2.13	0.49
1:A:70:GLN:NE2	1:A:70:GLN:HA	2.29	0.48
1:A:184:ASN:H	1:A:184:ASN:ND2	2.09	0.47
1:B:5:CYS:SG	1:B:33:ILE:HG13	2.55	0.47
1:A:18:GLU:OE2	1:A:22:LYS:HE3	2.14	0.47
1:A:18:GLU:OE2	4:A:303:MES:H82	2.15	0.47
1:A:135:ASN:HD22	1:A:179:ASN:HD21	1.61	0.46
1:B:4:LEU:O	1:B:30:VAL:HA	2.14	0.46
1:A:197:ILE:HG21	6:A:440:HOH:O	2.15	0.46
1:B:44:TRP:CD1	1:B:221:GLU:HG2	2.51	0.46
1:B:184:ASN:ND2	6:B:417:HOH:O	2.50	0.45
1:A:209:GLU:O	1:A:213:GLN:HG3	2.16	0.45
1:A:7:ALA:HB1	1:A:78:PHE:CD1	2.53	0.44
1:A:126:CYS:O	1:A:134:GLY:HA3	2.19	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:ASN:ND2	1:A:179:ASN:HD22	2.14	0.43
1:B:5:CYS:HB2	1:B:71:ILE:HD11	2.00	0.43
1:B:184:ASN:HD22	1:B:184:ASN:N	2.10	0.42
1:A:24:HIS:HE1	1:A:170:GLN:NE2	2.17	0.42
1:A:58:LYS:HE2	6:A:425:HOH:O	2.19	0.42
1:A:174:ASN:O	1:A:176:THR:HG23	2.21	0.41
1:A:62:LEU:HD23	1:A:62:LEU:HA	1.97	0.41
1:A:135:ASN:CB	1:A:179:ASN:HD22	2.34	0.41
1:A:220:LYS:HD3	6:A:618:HOH:O	2.20	0.41
1:A:204:LYS:O	1:A:204:LYS:HG3	2.21	0.41
1:B:62:LEU:HD23	1:B:62:LEU:HA	1.89	0.40
1:A:148:ASP:O	1:A:215:ARG:HD2	2.21	0.40
1:A:225:PRO:HG2	1:A:232:ILE:HB	2.04	0.40
1:B:44:TRP:CE3	1:B:45:GLN:HG2	2.56	0.40
1:B:78:PHE:CE2	1:B:80:ARG:HB2	2.56	0.40
1:B:111:THR:N	1:B:112:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/280 (95%)	256 (96%)	11 (4%)	0	100	100
1	B	269/280 (96%)	258 (96%)	11 (4%)	0	100	100
All	All	536/560 (96%)	514 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/260 (96%)	237 (95%)	12 (5%)	25	22
1	B	249/260 (96%)	239 (96%)	10 (4%)	31	29
All	All	498/520 (96%)	476 (96%)	22 (4%)	28	25

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	62	LEU
1	A	74	PHE
1	A	85	GLU
1	A	175	ASN
1	A	184	ASN
1	A	200	LYS
1	A	204	LYS
1	A	219	PHE
1	A	228	ASN
1	A	234	LYS
1	A	239	ASN
1	B	26	LYS
1	B	28	ASP
1	B	37	ASN
1	B	74	PHE
1	B	82	VAL
1	B	85	GLU
1	B	175	ASN
1	B	184	ASN
1	B	200	LYS
1	B	219	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	70	GLN
1	A	170	GLN
1	A	174	ASN
1	A	179	ASN
1	A	184	ASN
1	A	228	ASN
1	A	239	ASN
1	A	267	GLN
1	B	37	ASN
1	B	54	ASN
1	B	94	ASN
1	B	184	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	EDO	B	303	-	3,3,3	0.17	0	2,2,2	0.20	0
4	MES	A	303	-	12,12,12	0.79	0	14,16,16	0.55	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	A	306	-	3,3,3	0.28	0	2,2,2	0.44	0
5	EDO	A	305	-	3,3,3	0.22	0	2,2,2	0.35	0
2	FON	B	301	-	28,36,36	0.93	2 (7%)	28,50,50	1.31	5 (17%)
2	FON	A	301	-	28,36,36	1.00	3 (10%)	28,50,50	1.49	7 (25%)
5	EDO	A	304	-	3,3,3	0.05	0	2,2,2	0.01	0
3	T3F	B	302	-	31,37,37	0.84	1 (3%)	42,57,57	2.46	4 (9%)
5	EDO	A	307	-	3,3,3	0.30	0	2,2,2	0.26	0
3	T3F	A	302	-	31,37,37	0.88	2 (6%)	42,57,57	2.43	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	303	-	-	0/1/1/1	-
4	MES	A	303	-	-	4/6/14/14	0/1/1/1
5	EDO	A	306	-	-	0/1/1/1	-
5	EDO	A	305	-	-	1/1/1/1	-
2	FON	B	301	-	-	3/18/37/37	0/2/3/3
2	FON	A	301	-	-	1/18/37/37	0/2/3/3
5	EDO	A	304	-	-	0/1/1/1	-
3	T3F	B	302	-	-	4/18/53/53	0/3/3/3
5	EDO	A	307	-	-	1/1/1/1	-
3	T3F	A	302	-	-	0/18/53/53	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	T3F	C4-N3	3.31	1.38	1.33
2	B	301	FON	C4-N3	3.16	1.38	1.33
3	A	302	T3F	C4-N3	3.13	1.38	1.33
2	A	301	FON	C4-N3	3.01	1.38	1.33
2	A	301	FON	C4-C4A	-2.18	1.38	1.41
2	B	301	FON	C4A-N5	2.02	1.44	1.41
2	A	301	FON	C4A-N5	2.01	1.44	1.41
3	A	302	T3F	C6-C5	-2.00	1.34	1.40

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	T3F	C4-N3-C2	14.75	127.60	115.14
3	A	302	T3F	C4-N3-C2	14.46	127.35	115.14
2	B	301	FON	C4-C4A-C8A	3.17	116.92	114.44
2	A	301	FON	C4-C4A-C8A	3.13	116.89	114.44
2	A	301	FON	N3-C2-N1	-2.96	120.78	125.42
2	B	301	FON	C2-N1-C8A	2.90	121.04	114.54
2	A	301	FON	C2-N1-C8A	2.89	121.02	114.54
3	B	302	T3F	PA-O3A-PB	2.81	142.47	132.83
2	B	301	FON	C4-N3-C2	2.55	119.98	115.93
2	A	301	FON	C4-N3-C2	2.54	119.96	115.93
2	A	301	FON	O3-CP1-N5	-2.50	121.74	125.36
2	B	301	FON	N3-C2-N1	-2.31	121.80	125.42
2	A	301	FON	CB-CA-N	2.30	113.55	110.19
3	A	302	T3F	O1A-PA-O2A	2.29	123.55	112.24
2	A	301	FON	NA2-C2-N3	2.25	120.75	117.25
3	B	302	T3F	C5M-C5-C6	2.11	123.13	118.68
2	B	301	FON	O3-CP1-N5	-2.10	122.32	125.36
3	B	302	T3F	O2B-PB-O1B	2.07	122.46	112.24
3	A	302	T3F	C5M-C5-C6	2.05	123.01	118.68

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	301	FON	O3-CP1-N5-C6
4	A	303	MES	C7-C8-S-O1S
4	A	303	MES	C7-C8-S-O3S
2	B	301	FON	N-CA-CB-CG
3	B	302	T3F	PB-O3A-PA-O2A
3	B	302	T3F	PB-O3A-PA-O5'
2	B	301	FON	O3-CP1-N5-C4A
4	A	303	MES	C7-C8-S-O2S
2	A	301	FON	C15-C14-N10-C9
3	B	302	T3F	PA-O3A-PB-O2B
3	B	302	T3F	C5'-O5'-PA-O1A
4	A	303	MES	N4-C7-C8-S
5	A	305	EDO	O1-C1-C2-O2
5	A	307	EDO	O1-C1-C2-O2



There are no ring outliers.


4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	303	MES	2	0
5	A	305	EDO	1	0
2	B	301	FON	1	0
2	A	301	FON	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	269/280 (96%)	-0.37	1 (0%) 92 92	15, 26, 48, 68	0
1	B	271/280 (96%)	-0.11	7 (2%) 56 54	15, 33, 58, 70	0
All	All	540/560 (96%)	-0.24	8 (1%) 73 72	15, 30, 56, 70	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	81	ILE	6.0
1	A	269	TYR	4.8
1	B	269	TYR	2.6
1	B	1	MET	2.5
1	B	28	ASP	2.4
1	B	83	LYS	2.4
1	B	39	PHE	2.2
1	B	180	PHE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	A	306	4/4	0.79	0.19	47,49,51,52	0
2	FON	B	301	34/34	0.84	0.17	45,52,58,59	0
5	EDO	A	307	4/4	0.90	0.14	33,33,35,35	4
4	MES	A	303	12/12	0.91	0.18	68,71,76,76	0
5	EDO	B	303	4/4	0.91	0.11	29,30,30,35	0
5	EDO	A	305	4/4	0.93	0.12	38,42,44,44	0
2	FON	A	301	34/34	0.94	0.09	14,17,30,35	0
3	T3F	B	302	35/35	0.96	0.09	23,31,37,40	0
3	T3F	A	302	35/35	0.96	0.09	16,21,23,24	0
5	EDO	A	304	4/4	0.97	0.14	35,37,38,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around FON A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around T3F B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around T3F A 302:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.