



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 01:48 AM BST

PDB ID : 4N97  
Title : E. coli sliding clamp in complex with 5-nitroindole  
Authors : Yin, Z.; Oakley, A.J.  
Deposited on : 2013-10-19  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>4%</div> <div>84%</div> <div>14%</div> <div>..</div> </div>
1	B	366	<div> <div>4%</div> <div>84%</div> <div>15%</div> <div>..</div> </div>

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase III subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	4	0
			2821	1776	489	536	20			
1	B	364	Total	C	N	O	S	0	5	0
			2850	1793	501	537	19			

- Molecule 2 is 5-nitro-1H-indole (three-letter code: 2HU) (formula: C<sub>8</sub>H<sub>6</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			12	8	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	339	Total	O	0	0
			339	339		
5	B	336	Total	O	0	0
			336	336		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase III subunit beta



- Molecule 1: DNA polymerase III subunit beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.88Å 64.78Å 72.59Å 73.91° 84.51° 84.89°	Depositor
Resolution (Å)	28.90 – 1.97 28.90 – 1.97	Depositor EDS
% Data completeness (in resolution range)	84.7 (28.90-1.97) 84.7 (28.90-1.97)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.38 (at 1.96Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.232 , 0.306 0.235 , 0.305	Depositor DCC
$R_{free}$ test set	2150 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	6387	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PEG, 2HU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/2881	0.68	1/3897 (0.0%)
1	B	0.42	0/2914	0.65	0/3943
All	All	0.42	0/5795	0.67	1/7840 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	256	LEU	CA-CB-CG	6.75	130.83	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2821	0	2830	35	0
1	B	2850	0	2868	40	0
2	A	12	0	6	0	0
3	A	21	0	30	5	0
3	B	7	0	10	0	0
4	A	1	0	0	0	0
5	A	339	0	0	5	0
5	B	336	0	0	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6387	0	5744	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:282[B]:ARG:HG2	1:B:282[B]:ARG:HH11	1.15	1.11
1:B:282[B]:ARG:HH11	1:B:282[B]:ARG:CG	1.78	0.95
1:B:156:ASN:O	1:B:197:ARG:HG3	1.74	0.87
1:A:344:VAL:HG12	1:A:365:ARG:NH2	2.04	0.71
1:B:42:LEU:HB3	1:B:57:VAL:HG13	1.71	0.71
1:B:282[B]:ARG:HG2	1:B:282[B]:ARG:NH1	1.94	0.70
1:B:143:GLN:HG3	1:B:146:MET:CE	2.23	0.67
1:B:281:VAL:HG12	1:B:294:ALA:HB2	1.78	0.66
1:B:143:GLN:HG3	1:B:146:MET:HE2	1.79	0.65
1:B:282[B]:ARG:NH1	1:B:282[B]:ARG:CG	2.49	0.64
1:A:27:LEU:HD12	1:A:28:PRO:HD2	1.79	0.63
1:A:118:ASN:OD1	1:A:118:ASN:N	2.32	0.62
1:B:73:ARG:NH1	5:B:785:HOH:O	2.30	0.62
1:A:184:ILE:HD11	1:A:188:LEU:HD11	1.82	0.62
1:A:249:PRO:HD2	1:A:348:GLN:HE21	1.67	0.60
1:B:126:VAL:HG13	1:B:218:ILE:HB	1.84	0.59
1:B:184:ILE:HD11	1:B:188:LEU:HD11	1.83	0.59
1:A:56:ARG:HD2	5:A:812:HOH:O	2.03	0.58
1:B:282[B]:ARG:CB	1:B:282[B]:ARG:HH11	2.18	0.56
1:B:214:LEU:HD11	1:B:225:ALA:HB1	1.87	0.55
1:B:282[B]:ARG:NH1	1:B:282[B]:ARG:HB3	2.21	0.55
1:B:211:ASP:OD1	1:B:211:ASP:N	2.39	0.54
1:A:214:LEU:HD11	1:A:225:ALA:HB1	1.88	0.53
1:A:365:ARG:HD2	5:A:832:HOH:O	2.09	0.53
1:A:33:LEU:O	1:A:69:THR:HA	2.11	0.50
1:A:365:ARG:NH2	5:A:576:HOH:O	2.44	0.50
1:A:338[A]:MET:CG	1:A:347:VAL:HG11	2.42	0.49
1:B:98:LEU:HD21	1:B:100:ARG:HD3	1.94	0.49
1:B:282[B]:ARG:CB	1:B:282[B]:ARG:NH1	2.76	0.49
1:A:177:LEU:HD21	1:A:179:VAL:HG22	1.95	0.49
1:B:12:LYS:HB3	1:B:13:PRO:HD3	1.95	0.48
1:B:126:VAL:CG1	1:B:218:ILE:HB	2.43	0.48
1:A:144:PHE:CD1	1:A:326:ASP:HB3	2.49	0.48

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:SER:H	3:A:404:PEG:H42	1.78	0.48
1:B:2:LYS:HB3	1:B:64:GLU:HG2	1.94	0.48
1:B:161:GLU:OE2	1:B:168:ARG:NH2	2.42	0.47
1:A:170:VAL:HG22	1:A:179:VAL:HG13	1.95	0.47
1:B:129:THR:H	1:B:186:GLN:HE22	1.61	0.47
1:A:299:GLN:HE21	1:A:299:GLN:HA	1.80	0.47
1:A:44:LEU:HD12	1:A:44:LEU:N	2.30	0.47
1:B:243:ASP:OD1	1:B:245[B]:ARG:HG2	2.14	0.47
1:B:48:ASP:O	1:B:49:LEU:HB2	2.15	0.47
1:B:284[A]:TYR:HD1	5:B:828:HOH:O	1.98	0.47
1:A:281:VAL:HG12	1:A:294:ALA:HB2	1.97	0.46
1:A:140:GLU:HG3	1:A:204:MET:CE	2.46	0.46
1:B:323:TYR:O	1:B:327:VAL:HG23	2.16	0.46
1:A:95:GLU:O	1:A:95:GLU:OE1	2.33	0.46
1:A:129:THR:HG23	1:A:215:ARG:HH11	1.81	0.45
1:B:198:LYS:NZ	5:B:824:HOH:O	2.47	0.45
1:A:338[A]:MET:HG3	1:A:347:VAL:HG11	1.99	0.45
1:A:193:VAL:HB	1:A:236:LEU:HD13	1.98	0.45
1:A:163:GLU:OE1	1:A:168:ARG:NH2	2.39	0.45
1:B:249:PRO:HD2	1:B:348:GLN:HE21	1.82	0.44
1:B:285:VAL:HG22	1:B:310:TYR:CE2	2.52	0.44
1:B:2:LYS:HB3	1:B:64:GLU:CG	2.47	0.44
1:A:271:ALA:HB2	1:A:321:VAL:HG21	2.00	0.44
3:A:403:PEG:H31	5:A:830:HOH:O	2.17	0.44
1:B:277:LYS:HE3	1:B:278:PHE:CZ	2.53	0.44
1:B:348:GLN:OE1	1:B:350:GLU:OE1	2.36	0.44
1:A:124:SER:H	3:A:404:PEG:C4	2.31	0.44
1:A:256:LEU:HA	1:A:309:THR:O	2.17	0.43
1:B:243:ASP:OD1	1:B:245[B]:ARG:CG	2.66	0.43
1:A:298:GLU:OE1	1:A:298:GLU:N	2.51	0.43
1:B:77:ASP:OD1	1:B:80:ARG:NH2	2.52	0.43
1:A:254:LYS:HE3	1:A:313:ALA:HB3	2.01	0.43
1:A:269:ARG:HH22	3:A:403:PEG:C3	2.32	0.43
1:B:168:ARG:HD3	1:B:181:SER:OG	2.19	0.42
1:A:27:LEU:CD1	1:A:28:PRO:HD2	2.50	0.41
1:B:317:ILE:HD11	1:B:363:PRO:HB3	2.02	0.41
1:B:191:HIS:CD2	5:B:646:HOH:O	2.73	0.41
1:B:33:LEU:HG	1:B:72:ALA:HB2	2.02	0.41
1:A:124:SER:N	3:A:404:PEG:H42	2.36	0.41
1:B:282[B]:ARG:HH11	1:B:282[B]:ARG:HB3	1.80	0.41
1:A:212:ASN:N	5:A:504:HOH:O	2.53	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:PRO:HB2	1:A:47:THR:HA	2.03	0.40
1:A:344:VAL:HG12	1:A:365:ARG:HH21	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/366 (99%)	350 (97%)	11 (3%)	0	100	100
1	B	365/366 (100%)	353 (97%)	11 (3%)	1 (0%)	41	29
All	All	726/732 (99%)	703 (97%)	22 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	252	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/313 (99%)	300 (97%)	10 (3%)	39	28
1	B	313/313 (100%)	305 (97%)	8 (3%)	46	37
All	All	623/626 (100%)	605 (97%)	18 (3%)	42	31

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	ARG
1	A	118	ASN
1	A	176	ARG
1	A	179	VAL
1	A	207	LEU
1	A	256	LEU
1	A	267	PHE
1	A	277	LYS
1	A	340	LEU
1	A	350	GLU
1	B	24	ARG
1	B	44	LEU
1	B	176	ARG
1	B	197	ARG
1	B	211	ASP
1	B	227	VAL
1	B	240	ARG
1	B	354	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	HIS
1	A	123	GLN
1	A	191	HIS
1	A	226	HIS
1	A	295	ASN
1	A	299	GLN
1	A	348	GLN
1	A	355	GLN
1	B	9	HIS
1	B	186	GLN
1	B	191	HIS
1	B	295	ASN
1	B	335	ASN
1	B	348	GLN
1	B	355	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	PEG	A	404	-	6,6,6	0.49	0	5,5,5	0.58	0
2	2HU	A	401	-	10,13,13	1.15	0	12,18,18	1.69	3 (25%)
3	PEG	A	403	-	6,6,6	0.59	0	5,5,5	0.83	0
3	PEG	B	401	-	6,6,6	0.45	0	5,5,5	0.37	0
3	PEG	A	402	-	6,6,6	0.46	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	404	-	-	2/4/4/4	-
2	2HU	A	401	-	-	0/2/4/4	0/2/2/2
3	PEG	A	403	-	-	2/4/4/4	-
3	PEG	B	401	-	-	3/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	402	-	-	1/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	2HU	C10-C11-C7	4.21	109.79	106.20
2	A	401	2HU	C5-C4-N2	2.48	121.25	119.38
2	A	401	2HU	C10-C11-C12	-2.16	128.81	136.62

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	PEG	O2-C3-C4-O4
3	A	404	PEG	O2-C3-C4-O4
3	A	402	PEG	O2-C3-C4-O4
3	A	403	PEG	C4-C3-O2-C2
3	B	401	PEG	C1-C2-O2-C3
3	A	404	PEG	C4-C3-O2-C2
3	B	401	PEG	O2-C3-C4-O4
3	B	401	PEG	C4-C3-O2-C2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	PEG	3	0
3	A	403	PEG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	361/366 (98%)	0.33	14 (3%)	39 42	11, 19, 36, 51	6 (1%)
1	B	364/366 (99%)	0.39	15 (4%)	37 39	10, 19, 36, 44	10 (2%)
All	All	725/732 (99%)	0.36	29 (4%)	38 40	10, 19, 36, 51	16 (2%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	209	GLY	5.2
1	A	153	TYR	4.3
1	B	23	GLY	4.2
1	B	210	GLY	3.7
1	B	209	GLY	3.6
1	A	20	PRO	3.6
1	B	299	GLN	3.5
1	A	19	GLY	3.3
1	B	24	ARG	3.1
1	A	238	ASP	3.0
1	B	284[A]	TYR	2.9
1	A	208	ASP	2.9
1	B	211	ASP	2.9
1	B	19	GLY	2.7
1	B	297	PRO	2.7
1	B	26	THR	2.6
1	B	252	PRO	2.5
1	A	176	ARG	2.5
1	A	27	LEU	2.4
1	A	5	VAL	2.4
1	A	151	VAL	2.3
1	A	284[A]	TYR	2.2
1	A	336	VAL	2.2
1	B	17	VAL	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	26	THR	2.2
1	B	240	ARG	2.2
1	B	212	ASN	2.1
1	B	248	LEU	2.1
1	A	211	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PEG	A	404	7/7	0.75	0.21	33,33,34,34	0
3	PEG	A	403	7/7	0.84	0.24	21,23,25,26	0
2	2HU	A	401	12/12	0.84	0.18	33,35,35,36	0
3	PEG	B	401	7/7	0.85	0.14	27,28,30,30	0
3	PEG	A	402	7/7	0.87	0.13	27,28,30,31	0
4	CA	A	405	1/1	0.95	0.11	42,42,42,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.