



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 28, 2020 – 08:03 pm BST

PDB ID : 1NB7  
Title : HC-J4 RNA polymerase complexed with short RNA template strand  
Authors : O'Farrell, D.J.; Trowbridge, R.; Rowlands, D.J.; Jaeger, J.  
Deposited on : 2002-12-02  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RNA backbone	3102	1007 (3.16-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	E	4	<div> <div>75%</div> <div>25%</div> </div>
1	F	4	<div> <div>75%</div> <div>25%</div> </div>
2	A	570	<div> <div>44%</div> <div>51%</div> <div>5% •</div> </div>
2	B	570	<div> <div>44%</div> <div>50%</div> <div>5% •</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8927 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(\*UP\*UP\*UP\*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	4	Total	C	N	O	P	0	0	0
			77	36	8	30	3			
1	F	4	Total	C	N	O	P	0	0	0
			77	36	8	30	3			

- Molecule 2 is a protein called polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	566	Total	C	N	O	S	0	0	0
			4388	2763	777	816	32			
2	B	565	Total	C	N	O	S	0	0	0
			4381	2758	776	815	32			

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mn	0	0
			2	2		
3	A	2	Total	Mn	0	0
			2	2		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-R(\*UP\*UP\*UP\*U)-3'

Chain E: 




- Molecule 1: 5'-R(\*UP\*UP\*UP\*U)-3'

Chain F: 



- Molecule 2: polypeptide


Chain A: 





- Molecule 2: polyprotein

Chain B:  44%  50%  5% •



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.66Å 108.50Å 134.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.00 – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) (22.00-2.90)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.14	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.229 , 0.295	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8927	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	E	1.39	0/84	1.33	2/128 (1.6%)
1	F	1.23	0/84	1.44	1/128 (0.8%)
2	A	0.49	0/4484	0.74	1/6087 (0.0%)
2	B	0.52	1/4477 (0.0%)	0.73	0/6077
All	All	0.53	1/9129 (0.0%)	0.75	4/12420 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	142	SER	CB-OG	7.89	1.52	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	3	U	O4'-C1'-N1	7.28	114.03	108.20
1	E	4	U	C2'-C3'-O3'	5.29	122.17	113.70
2	A	351	GLY	N-CA-C	-5.12	100.29	113.10
1	E	4	U	N1-C1'-C2'	5.03	120.53	114.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	4	U	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	77	0	42	22	0
1	F	77	0	42	23	0
2	A	4388	0	4383	326	0
2	B	4381	0	4374	302	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
All	All	8927	0	8841	641	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 36.

The worst 5 of 641 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:515:GLY:HA2	2:B:519:ALA:HB2	1.33	1.06
2:B:144:VAL:HG22	2:B:394:ARG:HG2	1.35	1.03
2:B:160:ILE:HD12	2:B:282:SER:HG	1.27	1.00
2:A:195:TYR:HB3	2:A:199:GLN:HB2	1.43	1.00
1:F:1:U:HO5'	1:F:1:U:H6	1.03	1.00

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	564/570 (99%)	466 (83%)	74 (13%)	24 (4%)	2	10
2	B	563/570 (99%)	466 (83%)	76 (14%)	21 (4%)	3	13
All	All	1127/1140 (99%)	932 (83%)	150 (13%)	45 (4%)	3	11

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	23	ILE
2	A	505	ARG
2	A	565	SER
2	B	109	ARG
2	B	113	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	475/485 (98%)	450 (95%)	25 (5%)	22	54
2	B	474/485 (98%)	443 (94%)	31 (6%)	17	45
All	All	949/970 (98%)	893 (94%)	56 (6%)	19	49

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	29	SER
2	B	98	LYS
2	B	513	SER
2	B	55	ASP
2	B	68	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
2	A	483	ASN
2	A	514	GLN
2	B	483	ASN
2	A	502	HIS
2	B	24	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	E	4/4 (100%)	3 (75%)	3 (75%)
1	F	4/4 (100%)	3 (75%)	3 (75%)
All	All	8/8 (100%)	6 (75%)	6 (75%)

5 of 6 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	E	2	U
1	E	3	U
1	E	4	U
1	F	2	U
1	F	3	U

5 of 6 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	E	3	U
1	F	3	U
1	F	1	U
1	E	2	U
1	F	2	U

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.