



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2022 – 04:11 AM EST

PDB ID : 6NBW  
Title : Ternary Complex of Beta/Gamma-Actin with Profilin and AnCoA-NAA80  
Authors : Rebowski, G.; Boczkowska, M.; Dominguez, R.  
Deposited on : 2018-12-10  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>..</div> </div> </div>
2	N	235	<div> <div>6%</div> <div> <div></div> <div>63%</div> <div>8%</div> <div>29%</div> </div> </div>
3	P	140	<div> <div>12%</div> <div> <div></div> <div>92%</div> <div>6%</div> <div>.</div> </div> </div>

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 10938 atoms, of which 5345 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, cytoplasmic 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	367	Total	C	H	N	O	S	0	1	0
			5721	1822	2842	485	552	20			

- Molecule 2 is a protein called N-alpha-acetyltransferase 80.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	N	167	Total	C	H	N	O	S	0	0	0
			2692	852	1358	249	227	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	74	ALA	-	expression tag	UNP Q93015
N	75	GLY	-	expression tag	UNP Q93015
N	76	HIS	-	expression tag	UNP Q93015
N	77	MET	-	expression tag	UNP Q93015


- Molecule 3 is a protein called Profilin-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	P	138	Total	C	H	N	O	S	0	0	0
			2081	654	1040	179	201	7			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	P	
			43	10	12	5	13	3	

- Molecule 6 is LATRUNCULIN B (three-letter code: LAB) (formula:  $C_{20}H_{29}NO_5S$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	S	
			56	20	29	1	5	1	

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			14	3	8	3		
7	A	1	Total	C	H	O	0	0
			13	3	7	3		

- Molecule 8 is [(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-4-HYDROXY-3-(PHOSPHONOXY)TETRAHYDROFURAN-2-YL]METHYL (3R)-3-HYDROXY-2,2-DIMETHYL-4-OXO-4-{[3-OXO-3-({2-[(2-OXOPROPYL)THIO]ETHYL}AMINO)PROPYL]AMINO}BUTYL DIHYDROGEN DIPHOSPHATE (three-letter code: SOP) (formula:  $C_{24}H_{40}N_7O_{17}P_3S$ ).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
8	N	1	Total	C	H	N	O	P	S	0	0
			88	24	36	7	17	3	1		

- Molecule 9 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
9	N	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	116	Total	O	0	0
			116	116		
10	N	57	Total	O	0	0
			57	57		
10	P	31	Total	O	0	0
			31	31		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Actin, cytoplasmic 1



- Molecule 2: N-alpha-acetyltransferase 80



- Molecule 3: Profilin-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.41Å 115.59Å 132.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.10 – 2.50 26.10 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (26.10-2.50) 99.4 (26.10-2.50)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.09 (at 2.50Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.178 , 0.209 0.178 , 0.208	Depositor DCC
$R_{free}$ test set	1399 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.817	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 37.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10938	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SOP, CA, LAB, GOL, MES, HIC, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/2931	0.46	0/3966
2	N	0.26	0/1368	0.46	0/1854
3	P	0.27	0/1059	0.49	0/1430
All	All	0.27	0/5358	0.46	0/7250

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2879	2842	2842	21	0
2	N	1334	1358	1357	10	0
3	P	1041	1040	1040	5	0
4	A	1	0	0	0	0
5	A	31	12	12	0	0
6	A	27	29	29	2	0
7	A	12	15	16	0	0
8	N	52	36	35	0	0
9	N	12	13	13	0	0
10	A	116	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	N	57	0	0	0	0
10	P	31	0	0	0	0
All	All	5593	5345	5344	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:403:LAB:C16	6:A:403:LAB:N1	1.70	1.40
1:A:273:GLY:O	1:A:277:THR:HG23	1.88	0.73
2:N:107:ARG:NH1	2:N:111:SER:OG	2.31	0.63
1:A:372[A]:ARG:NH1	3:P:82:GLU:OE1	2.32	0.62
2:N:82:LEU:HD12	2:N:177:ARG:CZ	2.37	0.55
6:A:403:LAB:N1	6:A:403:LAB:C15	2.61	0.55
1:A:348:SER:HB3	2:N:222:THR:HG22	1.91	0.52
1:A:286:ASP:OD2	3:P:90:LYS:NZ	2.38	0.51
1:A:272:CYS:HB2	1:A:277:THR:HG22	1.94	0.49
1:A:345:ILE:HD11	2:N:227:PRO:HD2	1.95	0.49
1:A:4:ASP:O	1:A:5:ILE:HG22	2.12	0.48
1:A:351:THR:O	1:A:354:GLN:HG2	2.13	0.48
1:A:109:PRO:HB3	1:A:175:ILE:HD13	1.96	0.47
1:A:190:MET:HG2	1:A:209:VAL:HG21	1.96	0.47
1:A:352:PHE:HE2	1:A:356:TRP:CH2	2.31	0.47
2:N:301:ILE:HD12	2:N:301:ILE:O	2.16	0.46
1:A:25:ASP:HA	2:N:226:LEU:HD21	1.98	0.45
1:A:303:THR:HG22	1:A:303:THR:O	2.16	0.45
3:P:25:LYS:HE2	3:P:106:ASP:O	2.17	0.44
2:N:122:ALA:O	2:N:123:PHE:HB2	2.18	0.43
1:A:318:THR:HA	1:A:327:ILE:CD1	2.49	0.43
1:A:321:ALA:HB3	1:A:327:ILE:HD11	2.01	0.43
1:A:132:MET:HE3	1:A:132:MET:HB2	1.91	0.43
1:A:70:PRO:HG3	1:A:81:ASP:HB3	2.01	0.42
3:P:47:VAL:O	3:P:51:VAL:HG22	2.19	0.42
2:N:227:PRO:O	2:N:228:ALA:HB3	2.19	0.42
3:P:80:ASP:HA	3:P:83:PHE:CZ	2.54	0.42
1:A:349:LEU:HB3	1:A:351:THR:HG22	2.02	0.41
1:A:198:TYR:CE1	1:A:248:ILE:HG22	2.55	0.41
2:N:93:LEU:HD21	2:N:140:PRO:HG2	2.02	0.40
2:N:123:PHE:HA	2:N:124:PRO:C	2.41	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:ASP:O	1:A:4:ASP:CB	2.69	0.40
1:A:172:PRO:HA	1:A:175:ILE:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/374 (97%)	356 (98%)	6 (2%)	1 (0%)	41	61
2	N	159/235 (68%)	153 (96%)	5 (3%)	1 (1%)	25	43
3	P	136/140 (97%)	132 (97%)	4 (3%)	0	100	100
All	All	658/749 (88%)	641 (97%)	15 (2%)	2 (0%)	41	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
2	N	123	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/316 (99%)	303 (97%)	9 (3%)	42	69

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	N	147/201 (73%)	142 (97%)	5 (3%)	37	63
3	P	113/114 (99%)	112 (99%)	1 (1%)	78	92
All	All	572/631 (91%)	557 (97%)	15 (3%)	47	72

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	62	ARG
1	A	116	ARG
1	A	132	MET
1	A	225	GLN
1	A	281	SER
1	A	324	THR
1	A	372[A]	ARG
1	A	372[B]	ARG
2	N	78	SER
2	N	187	ARG
2	N	214	GLU
2	N	263	LEU
2	N	271	GLU
3	P	58	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	HIC	A	73	1	8,11,12	1.66	1 (12%)	6,14,16	1.14	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	73	1	-	0/5/6/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	HIC	CD2-CG	3.62	1.41	1.36

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	HIC	CB-CA-C	-2.22	107.31	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GOL	A	405	-	5,5,5	0.86	0	5,5,5	0.94	0
5	ATP	A	402	4	26,33,33	0.89	1 (3%)	31,52,52	1.37	4 (12%)
6	LAB	A	403	-	28,29,29	5.01	15 (53%)	30,41,41	4.53	6 (20%)
8	SOP	N	400	-	46,54,54	2.60	12 (26%)	56,80,80	2.11	10 (17%)
7	GOL	A	404	-	5,5,5	1.06	0	5,5,5	0.99	0
9	MES	N	401	-	12,12,12	2.21	1 (8%)	14,16,16	1.49	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	405	-	-	2/4/4/4	-
5	ATP	A	402	4	-	2/18/38/38	0/3/3/3
6	LAB	A	403	-	-	2/21/49/49	0/2/3/3
8	SOP	N	400	-	-	12/48/68/68	0/3/3/3
7	GOL	A	404	-	-	2/4/4/4	-
9	MES	N	401	-	-	0/6/14/14	0/1/1/1

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	403	LAB	C16-N1	17.76	1.70	1.46
6	A	403	LAB	C2-C3	10.86	1.54	1.33
8	N	400	SOP	C3'-C4'	-8.04	1.31	1.52
8	N	400	SOP	O4'-C4'	7.70	1.62	1.45
9	N	401	MES	C8-S	-7.32	1.67	1.77
8	N	400	SOP	O4'-C1'	-6.93	1.31	1.41
6	A	403	LAB	C7-C6	6.20	1.58	1.31
8	N	400	SOP	CP6-NP2	5.72	1.46	1.33
6	A	403	LAB	C18-S1	-5.67	1.65	1.77
6	A	403	LAB	C17-S1	-5.66	1.69	1.81
8	N	400	SOP	CP3-NP1	5.46	1.45	1.33
6	A	403	LAB	O2-C1	5.03	1.45	1.34
6	A	403	LAB	C18-N1	4.86	1.40	1.34
6	A	403	LAB	C12-C11	4.42	1.63	1.52
6	A	403	LAB	C10-C11	-4.21	1.37	1.51
6	A	403	LAB	C17-C16	-4.03	1.46	1.53
6	A	403	LAB	C2-C1	3.96	1.56	1.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	403	LAB	O3-C11	3.54	1.52	1.44
8	N	400	SOP	P3-O3'	3.18	1.65	1.59
8	N	400	SOP	C6-N6	3.16	1.45	1.34
6	A	403	LAB	C9-C10	2.91	1.61	1.53
6	A	403	LAB	O3-C15	2.79	1.48	1.43
8	N	400	SOP	O2'-C2'	-2.79	1.36	1.43
6	A	403	LAB	C4-C3	2.53	1.56	1.51
8	N	400	SOP	C5-C4	-2.35	1.34	1.40
5	A	402	ATP	C5-C4	2.25	1.46	1.40
8	N	400	SOP	C2-N3	2.24	1.35	1.32
8	N	400	SOP	OP2-CP6	-2.13	1.19	1.23
8	N	400	SOP	OP1-CP3	-2.13	1.18	1.23

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	403	LAB	C17-S1-C18	23.56	104.64	92.00
8	N	400	SOP	C1'-N9-C4	-9.07	110.70	126.64
8	N	400	SOP	C5-C6-N6	7.25	131.36	120.35
8	N	400	SOP	N3-C2-N1	-5.48	120.12	128.68
8	N	400	SOP	N6-C6-N1	-4.83	108.54	118.57
5	A	402	ATP	N3-C2-N1	-3.46	123.27	128.68
8	N	400	SOP	CP1-CP2-NP1	-3.28	105.53	112.42
5	A	402	ATP	C4-C5-N7	-2.99	106.28	109.40
6	A	403	LAB	C4-C5-C6	-2.97	106.17	112.59
5	A	402	ATP	PA-O3A-PB	-2.82	123.14	132.83
9	N	401	MES	O3S-S-C8	2.73	110.19	105.77
8	N	400	SOP	CPA-CP8-CP7	2.47	113.11	108.82
9	N	401	MES	O2S-S-C8	2.35	109.75	106.92
6	A	403	LAB	C19-C3-C4	2.32	119.17	115.27
6	A	403	LAB	O5-C18-S1	2.30	126.77	122.73
6	A	403	LAB	O5-C18-N1	-2.23	124.37	126.81
8	N	400	SOP	CP2-NP1-CP3	-2.20	118.75	122.84
5	A	402	ATP	C2-N1-C6	2.19	122.49	118.75
6	A	403	LAB	C9-C10-C11	-2.09	110.14	114.03
8	N	400	SOP	C3'-C2'-C1'	2.09	104.52	99.89
8	N	400	SOP	CP1-S-CA1	2.06	105.15	101.71
8	N	400	SOP	P2-O6-P1	-2.03	125.86	132.83
9	N	401	MES	O1S-S-C8	2.02	109.35	106.92

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	402	ATP	PB-O3B-PG-O3G
7	A	404	GOL	C1-C2-C3-O3
7	A	405	GOL	C1-C2-C3-O3
8	N	400	SOP	C5'-O5'-P1-O11
8	N	400	SOP	C5'-O5'-P1-O12
8	N	400	SOP	CP9-O7-P2-O6
8	N	400	SOP	CP9-O7-P2-O21
8	N	400	SOP	CP9-O7-P2-O22
7	A	404	GOL	O2-C2-C3-O3
7	A	405	GOL	O2-C2-C3-O3
8	N	400	SOP	OP3-CP7-CP8-CP9
6	A	403	LAB	O2-C1-C2-C3
6	A	403	LAB	O1-C1-C2-C3
8	N	400	SOP	CP2-CP1-S-CA1
5	A	402	ATP	PB-O3B-PG-O1G
8	N	400	SOP	C3'-O3'-P3-O33
8	N	400	SOP	C5'-O5'-P1-O6
8	N	400	SOP	OP3-CP7-CP8-CPB
8	N	400	SOP	CP8-CP9-O7-P2
8	N	400	SOP	CP6-CP7-CP8-CP9


There are no ring outliers.

1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	403	LAB	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






## Ligand LAB A 403



## Ligand SOP N 400



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	366/374 (97%)	0.11	20 (5%) 25 26	27, 35, 49, 76	0
2	N	167/235 (71%)	0.26	15 (8%) 9 9	29, 36, 84, 103	0
3	P	138/140 (98%)	0.54	17 (12%) 4 3	33, 41, 65, 77	0
All	All	671/749 (89%)	0.24	52 (7%) 13 13	27, 36, 63, 103	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	N	228	ALA	7.9
2	N	227	PRO	5.6
2	N	271	GLU	5.2
1	A	40	HIS	5.1
3	P	93	GLY	5.0
1	A	2	ASP	4.7
3	P	111	LEU	4.2
2	N	270	PRO	4.1
2	N	269	LEU	4.0
1	A	3	ASP	4.0
2	N	132	PRO	4.0
2	N	76	HIS	3.9
3	P	101	THR	3.9
1	A	48	GLY	3.9
1	A	4	ASP	3.8
2	N	230	LEU	3.7
3	P	102	VAL	3.7
2	N	225	ARG	3.6
3	P	110	VAL	3.3
3	P	95	ALA	3.3
2	N	300	PRO	3.2
2	N	226	LEU	3.2
3	P	94	GLY	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	N	268	PRO	3.2
1	A	104	LEU	3.1
3	P	20	ALA	3.0
1	A	360	GLN	3.0
2	N	133	HIS	3.0
3	P	80	ASP	3.0
3	P	96	PRO	2.9
3	P	112	LEU	2.9
1	A	10	VAL	2.8
1	A	105	LEU	2.8
3	P	26	ASP	2.7
3	P	79	GLN	2.7
2	N	229	THR	2.6
3	P	92	THR	2.6
1	A	135	ALA	2.6
1	A	244	ASP	2.5
1	A	350	SER	2.5
3	P	27	SER	2.4
3	P	127	CYS	2.4
1	A	106	THR	2.4
2	N	146	ALA	2.3
3	P	100	VAL	2.3
1	A	64	ILE	2.3
1	A	49	GLN	2.2
1	A	185	LEU	2.2
1	A	134	VAL	2.2
1	A	324	THR	2.1
1	A	5	ILE	2.1
1	A	323	SER	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	HIC	A	73	11/12	0.96	0.13	28,31,39,39	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	GOL	A	404	6/6	0.69	0.31	34,40,45,45	0
9	MES	N	401	12/12	0.82	0.22	41,50,55,69	0
7	GOL	A	405	6/6	0.94	0.28	44,52,55,59	0
6	LAB	A	403	27/27	0.95	0.16	26,31,34,36	0
8	SOP	N	400	52/52	0.96	0.13	29,35,40,43	0
5	ATP	A	402	31/31	0.98	0.15	24,29,35,38	0
4	CA	A	401	1/1	0.98	0.15	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around LAB A 403:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SOP N 400:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ATP A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.