



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2021 – 07:21 PM EDT

PDB ID : 1NEY  
Title : Triosephosphate Isomerase in Complex with DHAP  
Authors : Jogl, G.; Rozovsky, S.; McDermott, A.E.; Tong, L.  
Deposited on : 2002-12-12  
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	<div> <div>2%</div> <div> <div></div> <div>94%</div> <div>5%</div> </div> </div>
1	B	247	<div> <div></div> <div> <div>93%</div> <div>6%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8430 atoms, of which 3842 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called triosephosphate isomerase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	247	Total	C	F	H	N	O	S	0	5	0
			3804	1198	1	1911	324	368	2			
1	B	247	Total	C	F	H	N	O	S	0	11	0
			3826	1203	1	1921	326	373	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	TYR	TRP	engineered mutation	UNP P00942
A	157	PHE	TRP	engineered mutation	UNP P00942
A	168	FTR	TRP	engineered mutation	UNP P00942
B	90	TYR	TRP	engineered mutation	UNP P00942
B	157	PHE	TRP	engineered mutation	UNP P00942
B	168	FTR	TRP	engineered mutation	UNP P00942

- Molecule 2 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula:  $C_3H_7O_6P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	O	P	0	0
			15	3	5	6	1		
2	B	1	Total	C	H	O	P	0	0
			15	3	5	6	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	377	Total	O	0	0
			377	377		
3	B	393	Total	O	0	0
			393	393		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: triosephosphate isomerase

Chain A:  94% 5% 2%



- Molecule 1: triosephosphate isomerase

Chain B:  93% 6% 1%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.62Å 82.80Å 38.21Å 90.00° 101.75° 90.00°	Depositor
Resolution (Å)	30.00 – 1.20 28.82 – 1.20	Depositor EDS
% Data completeness (in resolution range)	95.7 (30.00-1.20) 93.2 (28.82-1.20)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.38 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.125 , 0.150 0.134 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	5.9	Xtriage
Anisotropy	0.637	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 64.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	8430	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FTR, 13P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/1932	1.21	17/2606 (0.7%)
1	B	0.69	0/1979	1.16	11/2668 (0.4%)
All	All	0.67	0/3911	1.18	28/5274 (0.5%)

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	26[A]	ARG	NE-CZ-NH2	9.99	125.30	120.30
1	A	26[B]	ARG	NE-CZ-NH2	9.99	125.30	120.30
1	A	247	ARG	CG-CD-NE	9.79	132.35	111.80
1	A	145[A]	ARG	NE-CZ-NH1	9.36	124.98	120.30
1	A	145[B]	ARG	NE-CZ-NH1	9.36	124.98	120.30
1	B	26[A]	ARG	NE-CZ-NH2	8.57	124.59	120.30
1	B	26[B]	ARG	NE-CZ-NH2	8.57	124.59	120.30
1	B	183	ASP	CB-CG-OD1	8.55	126.00	118.30
1	A	26[A]	ARG	CD-NE-CZ	8.26	135.17	123.60
1	A	26[B]	ARG	CD-NE-CZ	8.26	135.17	123.60
1	B	247	ARG	NE-CZ-NH1	7.21	123.91	120.30
1	A	141	ASP	CB-CG-OD1	6.64	124.28	118.30
1	A	101	TYR	CB-CG-CD2	6.48	124.89	121.00
1	B	183	ASP	CB-CG-OD2	-6.36	112.58	118.30
1	B	247	ARG	CG-CD-NE	6.30	125.04	111.80
1	A	145[A]	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	A	145[B]	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	B	132	GLU	OE1-CD-OE2	-5.50	116.70	123.30
1	B	26[A]	ARG	CG-CD-NE	5.37	123.09	111.80
1	B	26[B]	ARG	CG-CD-NE	5.37	123.09	111.80
1	B	98	ARG	NE-CZ-NH2	5.34	122.97	120.30
1	A	98	ARG	CG-CD-NE	5.27	122.86	111.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	208	TYR	CB-CG-CD2	5.24	124.14	121.00
1	A	98	ARG	CD-NE-CZ	5.21	130.89	123.60
1	A	26[A]	ARG	NH1-CZ-NH2	-5.03	113.87	119.40
1	A	26[B]	ARG	NH1-CZ-NH2	-5.03	113.87	119.40
1	A	173	GLY	C-N-CA	5.02	134.24	121.70
1	B	157	PHE	CB-CG-CD1	5.02	124.31	120.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1893	1911	1902	5	0
1	B	1905	1921	1904	6	0
2	A	10	5	5	0	0
2	B	10	5	5	0	0
3	A	377	0	0	1	0
3	B	393	0	0	2	0
All	All	4588	3842	3816	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:LYS:HE2	3:A:5174:HOH:O	2.08	0.54
1:B:140:LEU:HD21	1:B:187[B]:SER:OG	2.11	0.51
1:B:132:GLU:HA	1:B:135:LYS:NZ	2.27	0.50
1:A:41:CYS:HA	1:A:62:GLY:O	2.17	0.45
1:B:236:LEU:HB3	3:B:6392:HOH:O	2.18	0.43
1:B:41:CYS:HA	1:B:62:GLY:O	2.18	0.43
1:A:6:PHE:O	1:A:228:GLY:HA3	2.19	0.42
1:B:6:PHE:O	1:B:228:GLY:HA3	2.19	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:PHE:CE1	1:A:228:GLY:HA2	2.55	0.42
1:B:55:LYS:NZ	3:B:6390:HOH:O	2.51	0.42
1:A:26[B]:ARG:HH21	1:A:26[B]:ARG:HD2	1.70	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/247 (101%)	241 (97%)	8 (3%)	0	100	100
1	B	255/247 (103%)	245 (96%)	10 (4%)	0	100	100
All	All	504/494 (102%)	486 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/199 (102%)	201 (98%)	3 (2%)	65	29
1	B	210/199 (106%)	207 (99%)	3 (1%)	67	32
All	All	414/398 (104%)	408 (99%)	6 (1%)	69	32

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LYS
1	A	98	ARG
1	A	159	ASN
1	B	98	ARG
1	B	159[A]	ASN
1	B	159[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	245	ASN
1	B	148	ASN
1	B	245	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	FTR	A	168	1	14,16,17	1.04	0	14,22,24	1.32	3 (21%)
1	FTR	B	168	1	14,16,17	1.02	1 (7%)	14,22,24	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FTR	A	168	1	-	0/4/6/8	0/2/2/2
1	FTR	B	168	1	-	0/4/6/8	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	168	FTR	CH2-CZ3	2.52	1.42	1.37

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	FTR	CZ3-CE3-CD2	-2.83	116.58	118.80
1	A	168	FTR	CE3-CD2-CE2	2.18	121.24	118.26
1	A	168	FTR	CH2-CZ2-CE2	-2.16	118.11	120.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	13P	B	6001	-	9,9,9	3.75	2 (22%)	10,12,12	1.06	0
2	13P	A	5001	-	9,9,9	4.26	4 (44%)	10,12,12	1.26	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	13P	B	6001	-	-	3/7/8/8	-
2	13P	A	5001	-	-	3/7/8/8	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5001	13P	O1-C1	11.56	1.51	1.43
2	B	6001	13P	O1-C1	10.04	1.50	1.43
2	B	6001	13P	P-O1	4.04	1.73	1.60
2	A	5001	13P	P-O1	3.67	1.72	1.60
2	A	5001	13P	P-O1P	2.35	1.58	1.50
2	A	5001	13P	C1-C2	2.23	1.54	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5001	13P	O3-C3-C2	2.60	121.12	112.43

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	6001	13P	O2-C2-C3-O3
2	A	5001	13P	O1-C1-C2-O2
2	B	6001	13P	O1-C1-C2-O2
2	A	5001	13P	O2-C2-C3-O3
2	B	6001	13P	O1-C1-C2-C3
2	A	5001	13P	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	246/247 (99%)	-0.51	4 (1%) 72 72	4, 9, 20, 47	0
1	B	246/247 (99%)	-0.57	0 100 100	4, 7, 16, 28	0
All	All	492/494 (99%)	-0.54	4 (0%) 86 86	4, 8, 18, 47	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	7.5
1	A	34	GLU	3.2
1	A	101	TYR	2.7
1	A	35	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	FTR	A	168	15/16	0.99	0.05	6,7,12,39	0
1	FTR	B	168	15/16	0.99	0.05	4,6,9,36	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	13P	A	5001	10/10	0.97	0.09	9,14,21,32	0
2	13P	B	6001	10/10	0.97	0.09	7,14,24,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.