



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 12:24 PM BST

PDB ID : 4NFU
Title : Structure of the central plant immunity signaling node EDS1 in complex with its interaction partner SAG101
Authors : Wagner, S.; Stüttmann, J.; Rietz, S.; Guerois, R.; Niefind, K.; Parker, J.E.
Deposited on : 2013-11-01
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	636	<div> <div>8%</div> <div>91%</div> <div>7%</div> <div>.</div> </div>
2	B	540	<div> <div>9%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9638 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EDS1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	619	Total	C	N	O	S	0	2	0
			5024	3192	865	946	21			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	expression tag	UNP Q9XF23
A	-11	GLY	-	expression tag	UNP Q9XF23
A	-10	SER	-	expression tag	UNP Q9XF23
A	-9	SER	-	expression tag	UNP Q9XF23
A	-8	HIS	-	expression tag	UNP Q9XF23
A	-7	HIS	-	expression tag	UNP Q9XF23
A	-6	HIS	-	expression tag	UNP Q9XF23
A	-5	HIS	-	expression tag	UNP Q9XF23
A	-4	HIS	-	expression tag	UNP Q9XF23
A	-3	HIS	-	expression tag	UNP Q9XF23
A	-2	SER	-	expression tag	UNP Q9XF23
A	-1	GLN	-	expression tag	UNP Q9XF23
A	0	ASP	-	expression tag	UNP Q9XF23
A	1	PRO	-	expression tag	UNP Q9XF23

- Molecule 2 is a protein called Senescence-associated carboxylesterase 101.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	524	Total	C	N	O	S	0	2	0
			4273	2738	715	796	24			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	SER	-	expression tag	UNP Q4F883
B	-1	GLN	-	expression tag	UNP Q4F883

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	0	ASP	-	expression tag	UNP Q4F883
B	1	PRO	-	expression tag	UNP Q4F883

- Molecule 3 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 3 1	0	0
4	A	1	Total C O 4 3 1	0	0
4	B	1	Total C O 4 3 1	0	0

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C N O S 15 8 2 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	169	Total O 169 169	0	0
6	B	133	Total O 133 133	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EDS1



• Molecule 2: Senescence-associated carboxylesterase 101



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	112.59Å 113.64Å 125.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.28 – 2.21 34.28 – 2.21	Depositor EDS
% Data completeness (in resolution range)	97.3 (34.28-2.21) 97.3 (34.28-2.21)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.22Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.176 , 0.196 0.179 , 0.195	Depositor DCC
R_{free} test set	1568 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	44.5	Xtriage
Anisotropy	0.621	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9638	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, EPE, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/5142	0.38	0/6944
2	B	0.22	0/4374	0.37	0/5893
All	All	0.22	0/9516	0.37	0/12837

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5024	0	4941	24	0
2	B	4273	0	4282	27	0
3	A	12	0	12	0	0
4	A	8	0	16	0	0
4	B	4	0	8	1	0
5	B	15	0	17	1	0
6	A	169	0	0	3	0
6	B	133	0	0	2	0
All	All	9638	0	9276	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:392:GLU:OE2	6:B:825:HOH:O	2.03	0.76
1:A:72:PRO:O	1:A:75:ARG:NH1	2.28	0.65
2:B:347:GLU:OE1	2:B:350:ARG:NH1	2.29	0.65
1:A:280:ARG:NH2	1:A:347:ASP:OD2	2.29	0.64
2:B:0:ASP:O	2:B:269:ARG:NH2	2.25	0.61
1:A:150:GLU:OE2	6:A:955:HOH:O	2.16	0.60
1:A:382:GLU:HG3	1:A:386:LYS:HE3	1.83	0.60
1:A:13:LEU:HD11	1:A:118:VAL:HG21	1.85	0.58
1:A:4:GLU:HG2	1:A:10:ASN:HA	1.90	0.54
1:A:46:SER:OG	1:A:51:ASP:OD2	2.20	0.53
1:A:481:ASP:OD1	2:B:428:ARG:NH2	2.36	0.53
1:A:111:VAL:HG13	1:A:149:LEU:HD21	1.89	0.52
1:A:261:PHE:CE1	2:B:139:PRO:HB2	2.45	0.51
2:B:135:LYS:HD3	2:B:167:LEU:HD21	1.93	0.51
2:B:228:VAL:HG11	2:B:271:ILE:HD11	1.94	0.49
1:A:100:ARG:NE	6:A:944:HOH:O	2.45	0.49
1:A:403:LYS:NZ	1:A:591:GLU:OE1	2.39	0.49
1:A:93:LEU:HD22	1:A:131:LEU:HB3	1.95	0.49
2:B:179:LEU:HG	2:B:208:ILE:HG21	1.96	0.48
2:B:77:ARG:NH2	2:B:245:ASP:O	2.34	0.47
2:B:292:MET:C	2:B:294:PHE:H	2.17	0.47
1:A:278:GLU:HG3	1:A:279:LYS:HG2	1.95	0.47
2:B:274:ASP:OD1	2:B:274:ASP:N	2.48	0.47
1:A:248:THR:HG21	2:B:8:LYS:HE2	1.97	0.47
2:B:300[B]:MET:HG3	2:B:351:PHE:CG	2.50	0.46
2:B:422:ILE:HG12	2:B:426:LYS:HE3	1.97	0.46
2:B:190:GLU:OE2	2:B:233[B]:HIS:NE2	2.42	0.46
2:B:142:ILE:HG22	2:B:152:ALA:HB1	1.98	0.46
2:B:405:GLY:N	4:B:602:IPA:H12	2.31	0.45
2:B:401:TYR:CE2	2:B:406:ARG:HG2	2.52	0.45
1:A:207:VAL:HG21	1:A:228:PHE:CG	2.51	0.45
2:B:4:SER:O	2:B:8:LYS:HG2	2.17	0.45
2:B:362:ARG:NH1	6:B:764:HOH:O	2.49	0.45
1:A:185:ARG:NH2	1:A:349:GLY:HA3	2.32	0.44
2:B:296:LYS:O	2:B:300[B]:MET:HG2	2.17	0.44
1:A:28:GLU:O	1:A:30:TYR:N	2.44	0.44
1:A:54:ASP:HA	1:A:55:PRO:HD3	1.88	0.44
1:A:50:LYS:HG3	1:A:55:PRO:HG3	1.99	0.44
1:A:334:HIS:O	1:A:338:GLU:HG2	2.18	0.44
2:B:356:VAL:HG21	2:B:409:HIS:HB2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:481:GLY:HA2	5:B:601:EPE:H51	2.01	0.43
2:B:303:SER:O	2:B:307:ILE:HG12	2.19	0.43
2:B:181:GLY:HA3	2:B:185:LEU:HD23	2.00	0.42
2:B:319:ILE:HA	2:B:500:GLU:HB2	2.01	0.42
2:B:24:SER:O	2:B:28:LYS:HG2	2.20	0.42
1:A:93:LEU:O	1:A:96:VAL:HG12	2.19	0.41
1:A:205:PRO:O	6:A:906:HOH:O	2.22	0.41
1:A:3:PHE:HD1	1:A:332:PHE:CE1	2.38	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	619/636 (97%)	604 (98%)	15 (2%)	0	100	100
2	B	522/540 (97%)	504 (97%)	17 (3%)	1 (0%)	47	54
All	All	1141/1176 (97%)	1108 (97%)	32 (3%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	293	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	546/560 (98%)	543 (100%)	3 (0%)	88	94
2	B	486/499 (97%)	483 (99%)	3 (1%)	86	92
All	All	1032/1059 (98%)	1026 (99%)	6 (1%)	86	92

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	123	SER
1	A	333	ASN
1	A	406	TYR
2	B	211	ASP
2	B	337	ASP
2	B	375	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	IPA	B	602	-	3,3,3	0.73	0	3,3,3	0.30	0
4	IPA	A	703	-	3,3,3	0.54	0	3,3,3	0.30	0
5	EPE	B	601	-	15,15,15	0.82	1 (6%)	18,20,20	1.71	5 (27%)
4	IPA	A	702	-	3,3,3	0.53	0	3,3,3	0.29	0
3	BGC	A	701	-	12,12,12	0.41	0	17,17,17	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	B	601	-	-	8/9/19/19	0/1/1/1
3	BGC	A	701	-	-	1/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	601	EPE	C10-S	2.76	1.81	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	601	EPE	C5-N4-C3	3.79	117.37	108.83
5	B	601	EPE	C7-N4-C5	3.14	119.27	111.23
5	B	601	EPE	C7-N4-C3	2.91	118.67	111.23
5	B	601	EPE	O1S-S-C10	2.29	109.67	106.92
5	B	601	EPE	O3S-S-C10	2.23	109.37	105.77

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	601	EPE	C8-C7-N4-C5
5	B	601	EPE	C9-C10-S-O1S
5	B	601	EPE	C9-C10-S-O2S
5	B	601	EPE	C9-C10-S-O3S
3	A	701	BGC	O5-C5-C6-O6
5	B	601	EPE	C10-C9-N1-C6
5	B	601	EPE	N4-C7-C8-O8
5	B	601	EPE	C10-C9-N1-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	601	EPE	S-C10-C9-N1

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	IPA	1	0
5	B	601	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	619/636 (97%)	0.20	54 (8%) 10 8	30, 57, 113, 154	0
2	B	524/540 (97%)	0.31	48 (9%) 9 7	34, 57, 122, 168	0
All	All	1143/1176 (97%)	0.25	102 (8%) 9 8	30, 57, 116, 168	0

All (102) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	620	ILE	8.9
2	B	52	ASP	8.2
1	A	216	SER	7.7
2	B	-2	SER	7.0
2	B	0	ASP	5.9
1	A	148	TYR	5.9
2	B	288	LEU	5.7
1	A	530	LEU	5.2
2	B	516	PHE	5.1
2	B	421	SER	5.1
2	B	422	ILE	5.1
2	B	-1	GLN	5.0
2	B	424	ILE	5.0
2	B	289	ARG	4.9
2	B	35	PRO	4.8
1	A	219	GLN	4.7
2	B	266	ALA	4.6
2	B	292	MET	4.5
2	B	419	MET	4.4
2	B	53	SER	4.4
1	A	291	LEU	4.3
1	A	304	GLN	4.3
2	B	263	LEU	4.3
2	B	290	PHE	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	462	VAL	4.3
2	B	294	PHE	4.2
1	A	619	GLU	4.2
1	A	302	ASP	4.1
2	B	431	LYS	4.1
1	A	527	GLY	4.0
1	A	160	LEU	4.0
1	A	528	LEU	3.9
2	B	265	LEU	3.8
1	A	190	PRO	3.7
1	A	223	GLN	3.7
2	B	293	MET	3.7
2	B	291	ASP	3.6
2	B	429	CYS	3.5
2	B	34	ASN	3.4
2	B	515	GLY	3.2
1	A	605	ASN	3.2
1	A	218	VAL	3.2
2	B	420	GLU	3.2
1	A	329	MET	3.2
1	A	339	ASN	3.1
1	A	192	ILE	3.1
1	A	290	ILE	3.1
2	B	78	SER	3.1
2	B	277	LYS	3.1
1	A	609	HIS	3.0
1	A	217	SER	3.0
1	A	294	LEU	3.0
1	A	68	ARG	3.0
1	A	189	VAL	2.9
1	A	194	LEU	2.9
2	B	423	LEU	2.9
1	A	303	GLU	2.9
1	A	158	ALA	2.9
2	B	125	TYR	2.9
1	A	532	GLU	2.8
2	B	1	PRO	2.7
1	A	270	ALA	2.7
1	A	269	PRO	2.7
2	B	287	ASN	2.6
2	B	77	ARG	2.6
1	A	273	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	430	LYS	2.6
1	A	222	GLU	2.5
2	B	428	ARG	2.5
1	A	531	GLU	2.5
2	B	167	LEU	2.5
1	A	271	GLY	2.4
2	B	154	LEU	2.4
1	A	311	PHE	2.4
1	A	188	ILE	2.4
2	B	537	GLN	2.4
1	A	159	PRO	2.3
1	A	561	GLU	2.3
1	A	557	GLY	2.3
1	A	224	ARG	2.3
1	A	130	ILE	2.3
2	B	470	GLU	2.3
2	B	267	ASP	2.3
1	A	161	VAL	2.2
1	A	180	VAL	2.2
1	A	220	GLU	2.2
1	A	377	GLN	2.2
2	B	466	ASP	2.2
1	A	529	GLN	2.2
2	B	135	LYS	2.1
1	A	3	PHE	2.1
1	A	236	THR	2.1
1	A	191	ARG	2.1
2	B	463	GLY	2.1
1	A	272	THR	2.1
2	B	384	MET	2.1
1	A	181	ASN	2.1
2	B	284	GLU	2.1
1	A	306	TRP	2.1
1	A	155	THR	2.0
2	B	427	GLU	2.0
2	B	274	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	IPA	B	602	4/4	0.49	0.29	79,82,84,84	0
5	EPE	B	601	15/15	0.78	0.38	63,102,183,184	0
4	IPA	A	703	4/4	0.82	0.24	89,92,93,95	0
4	IPA	A	702	4/4	0.83	0.20	65,68,73,82	0
3	BGC	A	701	12/12	0.86	0.17	77,98,110,112	0

6.5 Other polymers [i](#)

There are no such residues in this entry.