



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 04:19 am BST

PDB ID : 6NHC  
Title : Structure of human neuronal nitric oxide synthase R354A/G357D mutant heme domain in complex with (S)-6-(2,3-difluoro-5-(2-(1-methylazetidin-2-yl)ethyl)phenethyl)-4-methylpyridin-2-amine  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2018-12-21  
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	421	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>17%</div> <div>..</div> </div> </div>



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total 3400	C 2177	N 577	O 623	S 23	0	4	0
1	B	414	Total 3386	C 2168	N 575	O 622	S 21	0	3	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	ALA	ARG	engineered mutation	UNP P29475
A	357	ASP	GLY	engineered mutation	UNP P29475
B	354	ALA	ARG	engineered mutation	UNP P29475
B	357	ASP	GLY	engineered mutation	UNP P29475

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	B	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O		
			17	9	5	3		
3	B	1	Total	C	N	O		
			17	9	5	3		

- Molecule 4 is 6-[2-(2,3-difluoro-5-{2-[(2S)-1-methylazetidin-2-yl]ethyl}phenyl)ethyl]-4-methylpyridin-2-amine (three-letter code: KNV) (formula:  $C_{20}H_{25}F_2N_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	0	0
			25	20	2	3		
4	B	1	Total	C	F	N	0	0
			25	20	2	3		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	144	Total	O	0	0
			144	144		
7	B	133	Total	O	0	0
			133	133		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.18Å 122.35Å 164.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.74 – 2.16 49.74 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.74-2.16) 99.2 (49.74-2.16)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 2.16Å)	Xtriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, $R_{free}$	0.185 , 0.237 0.184 , 0.236	Depositor DCC
$R_{free}$ test set	2788 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.4	Xtriage
Anisotropy	0.856	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 56.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KNV, GOL, ZN, H4B, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3509	0.53	0/4762
1	B	0.38	0/3491	0.52	0/4735
All	All	0.38	0/7000	0.53	0/9497

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3400	0	3313	23	0
1	B	3386	0	3295	40	0
2	A	43	0	30	2	0
2	B	43	0	30	2	0
3	A	17	0	15	0	0
3	B	17	0	15	1	0
4	A	25	0	0	1	0
4	B	25	0	0	1	0
5	A	12	0	16	1	0
5	B	6	0	8	0	0
6	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	144	0	0	4	0
7	B	133	0	0	2	0
All	All	7252	0	6722	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:LEU:HD22	1:A:536:PRO:HB2	1.60	0.84
1:B:528:LEU:HD22	1:B:536:PRO:HB2	1.63	0.78
1:A:392:GLU:OE2	7:A:901:HOH:O	2.08	0.72
1:B:445:ASN:ND2	7:B:901:HOH:O	2.26	0.68
1:B:341:MET:HE2	3:B:803:H4B:H9	1.80	0.62
2:B:802:HEM:HBC2	2:B:802:HEM:HMC2	1.79	0.62
2:A:801:HEM:HMC2	2:A:801:HEM:HBC2	1.83	0.60
1:B:506:PHE:HA	1:B:509:ILE:HD12	1.83	0.60
1:B:664:ILE:O	1:B:668[B]:GLU:HG2	2.03	0.59
2:A:801:HEM:HHC	2:A:801:HEM:HBB2	1.85	0.59
1:A:328:GLU:O	1:A:704:ARG:NH1	2.37	0.57
1:B:327:LEU:HB2	1:B:704:ARG:HG2	1.86	0.57
1:B:482:SER:HA	1:B:574:ASN:HB3	1.87	0.56
1:B:419:ARG:HD3	1:B:683:TRP:CD2	2.39	0.56
1:A:441:HIS:HD2	7:A:1005:HOH:O	1.88	0.56
1:A:664:ILE:HG13	1:A:694:PRO:HB2	1.86	0.56
1:A:689:SER:HB3	1:A:692:ILE:HD11	1.88	0.56
2:B:802:HEM:HHC	2:B:802:HEM:HBB2	1.87	0.55
1:A:483:GLN:NE2	4:A:803:KNV:F12	2.30	0.54
1:B:625:LYS:HD3	1:B:626:THR:N	2.22	0.54
1:B:414:TRP:CE3	1:B:426:TRP:HA	2.46	0.51
1:B:547:LEU:HD21	1:B:651:VAL:HG22	1.94	0.50
1:B:704:ARG:NH1	1:B:710:GLU:OE2	2.44	0.50
1:A:350:PRO:C	1:A:352:ASP:H	2.16	0.49
1:A:480:TRP:HB2	1:A:528:LEU:HB3	1.94	0.49
1:B:664:ILE:HG13	1:B:694:PRO:HB2	1.94	0.49
1:B:480:TRP:HB2	1:B:528:LEU:HB3	1.94	0.48
1:A:593:TYR:CD1	1:A:598:ILE:HD11	2.49	0.48
1:B:625:LYS:HD3	1:B:626:THR:H	1.78	0.48
1:B:577:LEU:HD21	1:B:579:ILE:HD11	1.95	0.48
1:B:428:LYS:HA	1:B:428:LYS:HE2	1.95	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:592:TRP:O	4:B:804:KNV:N02	2.48	0.47
1:A:691:SER:HA	1:A:696:PHE:CG	2.50	0.47
1:A:429:LEU:O	5:A:805:GOL:H12	2.15	0.46
1:B:600:VAL:O	1:B:604:CYS:HB2	2.16	0.46
1:B:625:LYS:HD2	1:B:627:SER:H	1.80	0.46
1:B:501:PRO:HA	1:B:504:VAL:HG23	1.97	0.45
1:B:360:PHE:N	1:B:361:PRO:HD2	2.31	0.45
1:A:414:TRP:CE3	1:A:426:TRP:HA	2.51	0.45
1:B:625:LYS:HD2	1:B:627:SER:OG	2.16	0.45
1:A:485:ILE:HD13	1:A:546:VAL:HG13	1.99	0.44
1:A:661:GLU:HG3	7:A:1029:HOH:O	2.18	0.44
1:B:470:ARG:HG3	1:B:476:ASP:OD1	2.17	0.44
1:B:327:LEU:HB2	1:B:704:ARG:CG	2.48	0.44
1:B:489:GLY:HA3	1:B:500:ASP:O	2.18	0.44
1:B:376:ARG:NH1	7:B:907:HOH:O	2.50	0.43
1:A:594:MET:HA	1:A:654:VAL:O	2.18	0.43
1:B:356:LYS:HG2	1:B:397:SER:HA	2.01	0.43
1:B:596:THR:HA	1:B:639:ASN:HD21	1.83	0.43
1:B:594:MET:HA	1:B:654:VAL:O	2.19	0.42
1:A:483:GLN:HB2	1:A:486:ARG:HG2	2.02	0.42
1:B:323:LEU:HB2	1:B:324:LYS:HE3	2.00	0.42
1:B:490:TYR:CE1	1:B:519:ARG:HA	2.54	0.42
1:B:512:GLN:HG2	1:B:512:GLN:O	2.19	0.42
1:B:681:TRP:CZ2	1:B:685:VAL:HG21	2.54	0.42
1:A:340:ILE:HD13	1:B:699:GLU:HB3	2.02	0.42
1:A:327:LEU:HD12	1:A:704:ARG:NE	2.35	0.41
1:A:445:ASN:ND2	7:A:911:HOH:O	2.54	0.41
1:A:633:GLN:HG3	1:B:636:VAL:HG11	2.02	0.41
1:A:547:LEU:HD21	1:A:651:VAL:HG22	2.01	0.41
1:B:593:TYR:CD1	1:B:598:ILE:HD11	2.56	0.41
1:B:625:LYS:CD	1:B:627:SER:H	2.33	0.41
1:B:691:SER:HA	1:B:696:PHE:CG	2.56	0.40
1:A:517:PRO:HA	1:A:518:PRO:HD3	1.95	0.40
1:B:478:ARG:HD3	1:B:585:SER:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/421 (99%)	405 (97%)	11 (3%)	0	100	100
1	B	413/421 (98%)	400 (97%)	13 (3%)	0	100	100
All	All	829/842 (98%)	805 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/376 (100%)	360 (96%)	16 (4%)	29	27
1	B	373/376 (99%)	356 (95%)	17 (5%)	27	23
All	All	749/752 (100%)	716 (96%)	33 (4%)	28	25

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	302	CYS
1	A	304	ARG
1	A	327	LEU
1	A	341	MET
1	A	351	GLU
1	A	372	SER
1	A	380	LYS
1	A	386	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	394	ASP
1	A	528	LEU
1	A	552	ARG
1	A	578	GLU
1	A	613	GLU
1	A	625	LYS
1	A	632	ASP
1	A	713	PRO
1	B	304	ARG
1	B	324	LYS
1	B	353	VAL
1	B	372	SER
1	B	373	SER
1	B	394	ASP
1	B	474	LYS
1	B	508	GLU
1	B	519	ARG
1	B	528	LEU
1	B	552	ARG
1	B	560	LYS
1	B	617	LYS
1	B	650	LYS
1	B	665	LYS
1	B	720	VAL
1	B	722	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	483	GLN
1	B	639	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	B	805	-	5,5,5	0.41	0	5,5,5	0.28	0
4	KNV	A	803	-	25,27,27	0.35	0	28,38,38	1.72	7 (25%)
4	KNV	B	804	-	25,27,27	0.41	0	28,38,38	1.93	7 (25%)
2	HEM	A	801	1	27,50,50	1.90	4 (14%)	17,82,82	2.13	4 (23%)
5	GOL	A	805	-	5,5,5	0.43	0	5,5,5	0.32	0
3	H4B	B	803	-	16,18,18	0.79	0	11,26,26	2.66	5 (45%)
2	HEM	B	802	1	27,50,50	1.83	4 (14%)	17,82,82	2.40	3 (17%)
3	H4B	A	802	-	16,18,18	1.06	0	11,26,26	2.50	4 (36%)
5	GOL	A	804	-	5,5,5	0.39	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	805	-	-	3/4/4/4	-
4	KNV	A	803	-	-	5/10/19/19	0/3/3/3
4	KNV	B	804	-	-	7/10/19/19	0/3/3/3
2	HEM	A	801	1	-	0/6/54/54	-
5	GOL	A	805	-	-	4/4/4/4	-
3	H4B	B	803	-	-	0/8/17/17	0/2/2/2
2	HEM	B	802	1	-	0/6/54/54	-
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	804	-	-	3/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C3B-C2B	-5.21	1.33	1.40
2	B	802	HEM	C3B-C2B	-4.27	1.34	1.40
2	A	801	HEM	C3C-C2C	-3.87	1.35	1.40
2	A	801	HEM	C3B-CAB	3.86	1.55	1.47
2	B	802	HEM	C3C-C2C	-3.82	1.35	1.40
2	B	802	HEM	C3B-CAB	3.43	1.54	1.47
2	A	801	HEM	C3C-CAC	3.31	1.54	1.47
2	B	802	HEM	C3C-CAC	3.24	1.54	1.47

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	802	HEM	CBA-CAA-C2A	-8.51	96.80	112.49
3	A	802	H4B	C4-C4A-C8A	5.76	119.69	114.57
2	A	801	HEM	CBA-CAA-C2A	-5.61	102.14	112.49
3	B	803	H4B	C4-C4A-C8A	5.02	119.03	114.57
4	B	804	KNV	C02-N01-C06	4.67	121.64	118.10
4	A	803	KNV	C02-N01-C06	4.57	121.56	118.10
2	A	801	HEM	CBD-CAD-C3D	-4.48	104.22	112.48
4	B	804	KNV	C09-C11-C12	-4.31	117.27	120.73
4	B	804	KNV	C16-C11-C12	3.73	120.18	116.76
3	B	803	H4B	C4-C4A-N5	3.48	122.04	119.12
3	B	803	H4B	N3-C2-N1	-3.38	120.12	125.42
3	B	803	H4B	C4-N3-C2	3.25	121.10	115.93
2	A	801	HEM	CMC-C2C-C3C	3.23	130.71	124.68
3	A	802	H4B	N3-C2-N1	-3.22	120.37	125.42
4	B	804	KNV	C09-C08-C06	-3.13	105.98	112.99
4	A	803	KNV	C16-C11-C12	3.04	119.54	116.76
4	B	804	KNV	C08-C06-N01	2.98	120.38	115.95
3	A	802	H4B	C2-N1-C8A	2.89	121.02	114.54
4	A	803	KNV	C05-C06-N01	-2.79	119.94	122.90
3	B	803	H4B	C2-N1-C8A	2.79	120.79	114.54
4	A	803	KNV	C08-C06-N01	2.66	119.92	115.95
4	A	803	KNV	C09-C08-C06	-2.65	107.05	112.99
2	B	802	HEM	CBD-CAD-C3D	-2.65	107.60	112.48
4	A	803	KNV	C09-C11-C12	-2.37	118.82	120.73
4	B	804	KNV	C05-C06-N01	-2.37	120.39	122.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	H4B	C4-N3-C2	2.23	119.47	115.93
2	B	802	HEM	CMC-C2C-C3C	2.22	128.83	124.68
2	A	801	HEM	C1D-C2D-C3D	2.14	108.48	107.00
4	B	804	KNV	N02-C02-N01	2.09	119.80	116.49
4	A	803	KNV	N02-C02-N01	2.09	119.79	116.49

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	805	GOL	O1-C1-C2-C3
4	A	803	KNV	C08-C09-C11-C12
4	A	803	KNV	C17-C18-C22-C23
4	A	803	KNV	C17-C18-C22-N21
4	A	803	KNV	C06-C08-C09-C11
4	B	804	KNV	C08-C09-C11-C12
4	B	804	KNV	C17-C18-C22-C23
4	B	804	KNV	C17-C18-C22-N21
4	B	804	KNV	C06-C08-C09-C11
5	A	805	GOL	O1-C1-C2-C3
5	A	804	GOL	O1-C1-C2-C3
5	A	805	GOL	C1-C2-C3-O3
5	B	805	GOL	O1-C1-C2-O2
5	A	805	GOL	O1-C1-C2-O2
5	A	804	GOL	O1-C1-C2-O2
5	A	805	GOL	O2-C2-C3-O3
4	B	804	KNV	C08-C09-C11-C16
4	A	803	KNV	C08-C09-C11-C16
5	A	804	GOL	O2-C2-C3-O3
4	B	804	KNV	C14-C15-C17-C18
4	B	804	KNV	C16-C15-C17-C18
5	B	805	GOL	O2-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	803	KNV	1	0
4	B	804	KNV	1	0
2	A	801	HEM	2	0
5	A	805	GOL	1	0


*Continued on next page...*





*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	803	H4B	1	0
2	B	802	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	416/421 (98%)	0.15	6 (1%) 75 80	44, 62, 93, 130	0
1	B	414/421 (98%)	0.22	13 (3%) 49 58	45, 67, 112, 157	0
All	All	830/842 (98%)	0.18	19 (2%) 60 68	44, 64, 106, 157	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	353	VAL	7.6
1	B	351	GLU	6.6
1	A	302	CYS	5.1
1	B	326	THR	4.1
1	B	624	ARG	3.9
1	B	352	ASP	3.8
1	A	351	GLU	3.3
1	B	721	TRP	3.2
1	A	353	VAL	3.0
1	A	305	PHE	2.6
1	B	327	LEU	2.3
1	B	359	LEU	2.3
1	B	722	LYS	2.3
1	B	383	MET	2.1
1	A	673[A]	CYS	2.1
1	B	557	GLU	2.0
1	A	562	LEU	2.0
1	B	558	TRP	2.0
1	B	720	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	805	6/6	0.66	0.22	90,95,96,96	0
5	GOL	B	805	6/6	0.86	0.16	69,81,84,92	0
5	GOL	A	804	6/6	0.88	0.17	72,79,80,82	0
3	H4B	A	802	17/17	0.89	0.14	54,59,71,76	0
4	KNV	B	804	25/25	0.89	0.26	44,122,138,142	0
3	H4B	B	803	17/17	0.90	0.16	58,70,84,88	0
4	KNV	A	803	25/25	0.92	0.24	49,122,142,143	0
2	HEM	A	801	43/43	0.98	0.16	36,50,62,71	0
2	HEM	B	802	43/43	0.98	0.13	38,54,73,83	0
6	ZN	B	801	1/1	0.99	0.17	54,54,54,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around KNV B 804:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around KNV A 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around HEM A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers ⓘ

There are no such residues in this entry.