



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 07:51 am BST

PDB ID : 5NIA  
Title : Crystal structure of human LTA4H mutant D375N in open conformation (crystal form I)  
Authors : Stsiapanava, A.  
Deposited on : 2017-03-23  
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	617	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10239 atoms, of which 4931 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leukotriene A-4 hydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	609	Total	C	H	N	O	S	0	14	0
			9878	3182	4931	821	923	21			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP P09960
A	-5	HIS	-	expression tag	UNP P09960
A	-4	HIS	-	expression tag	UNP P09960
A	-3	HIS	-	expression tag	UNP P09960
A	-2	HIS	-	expression tag	UNP P09960
A	-1	HIS	-	expression tag	UNP P09960
A	0	HIS	-	expression tag	UNP P09960
A	375	ASN	ASP	engineered mutation	UNP P09960

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	351	Total	O	1	9
			360	360		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leukotriene A-4 hydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.00 Å   153.00 Å   74.88 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	44.17 – 1.76 44.17 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.17-1.76) 92.9 (44.17-1.76)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.56 (at 1.76 Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575)	Depositor
R, $R_{free}$	0.180 , 0.207 0.180 , 0.207	Depositor DCC
$R_{free}$ test set	3271 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10239	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.75	1/5111 (0.0%)	0.74	2/6949 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	25	CYS	CB-SG	-5.98	1.72	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	A	277	PHE	CB-CG-CD2	-5.03	117.28	120.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4947	4931	4916	53	0
2	A	1	0	0	0	0
3	A	360	0	0	5	0
All	All	5308	4931	4916	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:381[A]:VAL:N	1:A:382[A]:PRO:CD	2.48	0.77
1:A:352:SER:OG	1:A:381[A]:VAL:CG2	2.33	0.76
1:A:352:SER:OG	1:A:381[A]:VAL:HG21	1.92	0.69
1:A:27:VAL:HG22	1:A:34:LEU:CD2	2.24	0.68
1:A:380:SER:CB	1:A:382[A]:PRO:HD2	2.25	0.66
1:A:380:SER:OG	1:A:382[A]:PRO:HD2	1.97	0.64
1:A:261:LEU:HD13	1:A:265:PHE:CE2	2.33	0.63
1:A:27:VAL:HG22	1:A:34:LEU:HD21	1.84	0.59
1:A:585:VAL:HA	1:A:607:LEU:HD21	1.84	0.59
1:A:513:PRO:HG2	1:A:516:HIS:CE1	2.39	0.57
1:A:381[A]:VAL:O	1:A:383[A]:TYR:N	2.39	0.55
1:A:35[A]:THR:HG22	1:A:107:GLU:HG3	1.90	0.54
1:A:134[B]:GLN:OE1	3:A:1001:HOH:O	2.18	0.53
1:A:307:THR:O	1:A:418:SER:HA	2.08	0.52
1:A:116:GLN:HG3	1:A:132:PHE:CE1	2.45	0.52
1:A:381[A]:VAL:N	1:A:382[A]:PRO:HD3	2.26	0.51
1:A:352:SER:HG	1:A:381[A]:VAL:HG21	1.76	0.50
1:A:513:PRO:HB2	1:A:516:HIS:CE1	2.46	0.49
1:A:135:CYS:HA	1:A:139:HIS:HB2	1.94	0.49
1:A:114:ALA:HB2	1:A:139:HIS:HB3	1.94	0.49
1:A:261:LEU:HD13	1:A:265:PHE:CD2	2.49	0.47
1:A:513:PRO:HB2	1:A:516:HIS:ND1	2.28	0.47
1:A:604:GLY:O	1:A:608:LYS:N	2.46	0.47
1:A:148:ASP:HA	1:A:199:CYS:SG	2.55	0.47
1:A:35[A]:THR:HG23	3:A:1199:HOH:O	2.14	0.47
1:A:165:GLU:HG2	1:A:166:LEU:HG	1.96	0.47
1:A:380:SER:OG	1:A:382[A]:PRO:CD	2.63	0.45
1:A:381[B]:VAL:N	1:A:382[B]:PRO:HD2	2.31	0.45
1:A:66:ILE:HG21	1:A:100:ILE:HD11	1.99	0.45
1:A:513:PRO:CG	1:A:516:HIS:CE1	3.00	0.44
1:A:561:GLN:OE1	1:A:563:ARG:N	2.47	0.44
1:A:266:PRO:HG2	1:A:267:TYR:CZ	2.53	0.43
1:A:353:VAL:O	1:A:357:GLY:N	2.47	0.43
1:A:558:ALA:O	1:A:567:THR:HG23	2.18	0.43
1:A:205:VAL:HG21	1:A:259:LEU:CD2	2.48	0.43
1:A:24:ARG:NH2	3:A:1028:HOH:O	2.50	0.43
1:A:266:PRO:HA	1:A:562:GLY:O	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:PHE:CE2	1:A:318[A]:GLU:HG3	2.53	0.43
1:A:261:LEU:HD11	1:A:277:PHE:HB3	2.00	0.43
1:A:472:SER:HA	1:A:502:PHE:HE1	1.84	0.42
1:A:27:VAL:HG22	1:A:34:LEU:HD23	1.97	0.42
1:A:315:TRP:CG	1:A:365[A]:LEU:HG	2.54	0.42
1:A:447:TRP:CZ3	1:A:454:PRO:HB3	2.54	0.42
1:A:396[A]:LEU:HG	1:A:438:VAL:CG1	2.49	0.42
1:A:352:SER:OG	1:A:381[A]:VAL:HG22	2.14	0.42
1:A:381[A]:VAL:N	1:A:382[A]:PRO:HD2	2.31	0.41
1:A:23:LEU:O	1:A:158:ALA:HA	2.20	0.41
1:A:194:LYS:HE2	3:A:1324:HOH:O	2.19	0.41
1:A:176:GLY:N	3:A:1026:HOH:O	2.49	0.41
1:A:494:LEU:HA	1:A:498:GLN:OE1	2.20	0.41
1:A:374:PRO:HA	1:A:377:ALA:HB3	2.03	0.40
1:A:392:TYR:CZ	1:A:396[B]:LEU:HD11	2.57	0.40
1:A:511:PRO:HB3	1:A:544:GLN:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	621/617 (101%)	596 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	555/549 (101%)	551 (99%)	4 (1%)	84 75

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	LEU
1	A	383[A]	TYR
1	A	383[B]	TYR
1	A	516	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	380:SER	C	381[B]:VAL	N	3.12

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	609/617 (98%)	0.55	64 (10%) 6 8	21, 40, 97, 251	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	585	VAL	9.8
1	A	476	ILE	7.7
1	A	541	LEU	6.2
1	A	3	ILE	6.1
1	A	571	PHE	6.1
1	A	589	GLN	5.9
1	A	514	LEU	5.4
1	A	609	VAL	5.0
1	A	481	ASP	4.7
1	A	507	LEU	4.7
1	A	607	LEU	4.6
1	A	608	LYS	4.5
1	A	581	HIS	4.3
1	A	377	ALA	4.3
1	A	577	PHE	4.2
1	A	551	ILE	4.1
1	A	267	TYR	4.0
1	A	509	ARG	3.9
1	A	92	ILE	3.9
1	A	548	GLU	3.9
1	A	383[A]	TYR	3.9
1	A	588	TYR	3.9
1	A	543	ILE	3.8
1	A	362	PHE	3.6
1	A	586	ARG	3.4
1	A	516	HIS	3.2
1	A	512	LEU	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	90	LEU	3.1
1	A	356	PHE	3.0
1	A	547	TRP	2.9
1	A	578	ASP	2.9
1	A	488	ALA	2.9
1	A	4	VAL	2.8
1	A	483	LEU	2.8
1	A	69	GLN	2.8
1	A	587	THR	2.7
1	A	574	LEU	2.7
1	A	354	LYS	2.7
1	A	381[A]	VAL	2.7
1	A	582	ASP	2.6
1	A	567	THR	2.6
1	A	555	LEU	2.6
1	A	552	PRO	2.6
1	A	513	PRO	2.6
1	A	526	PHE	2.5
1	A	584	ALA	2.5
1	A	583	GLN	2.5
1	A	473	GLN	2.5
1	A	94	LEU	2.5
1	A	576	ALA	2.4
1	A	378	TYR	2.4
1	A	601	MET	2.4
1	A	580	SER	2.4
1	A	232	TYR	2.4
1	A	1	PRO	2.3
1	A	602	LEU	2.3
1	A	517	ILE	2.3
1	A	477	THR	2.2
1	A	568	ARG	2.2
1	A	472	SER	2.2
1	A	603	VAL	2.1
1	A	590	GLU	2.1
1	A	600	ALA	2.1
1	A	382[A]	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	900	1/1	1.00	0.10	31,31,31,31	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.