



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 16, 2021 – 10:36 PM EDT

PDB ID : 1NLW
Title : Crystal structure of Mad-Max recognizing DNA
Authors : Nair, S.K.; Burley, S.K.
Deposited on : 2003-01-07
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	F	18	
1	G	18	
1	H	18	
1	J	18	
2	A	80	
2	D	80	
3	B	76	
3	E	76	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4212 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*AP*GP*TP*AP*GP*CP*AP*CP*GP*TP*GP*CP*TP*AP*CP*TP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	18	Total	C	N	O	P	0	0	0
			366	175	68	106	17			
1	G	18	Total	C	N	O	P	0	0	0
			366	175	68	106	17			
1	H	18	Total	C	N	O	P	0	0	0
			366	175	68	106	17			
1	J	18	Total	C	N	O	P	0	0	0
			366	175	68	106	17			

- Molecule 2 is a protein called MAD PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	79	Total	C	N	O	S	0	0	0
			634	388	133	112	1			
2	D	77	Total	C	N	O	S	0	0	0
			609	376	122	110	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	SER	CYS	engineered mutation	UNP Q05195
A	33	ASP	GLU	variant	UNP Q05195
A	56	SER	CYS	engineered mutation	UNP Q05195
D	520	SER	CYS	engineered mutation	UNP Q05195
D	533	ASP	GLU	variant	UNP Q05195
D	556	SER	CYS	engineered mutation	UNP Q05195

- Molecule 3 is a protein called MAX PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	76	Total 637	C 387	N 131	O 118	S 1	0	0	0
3	E	74	Total 617	C 375	N 125	O 116	S 1	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	26	Total 26	O 26	0	0
4	G	38	Total 38	O 38	0	0
4	H	23	Total 23	O 23	0	0
4	J	21	Total 21	O 21	0	0
4	A	36	Total 36	O 36	0	0
4	B	38	Total 38	O 38	0	0
4	D	41	Total 41	O 41	0	0
4	E	28	Total 28	O 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D(*GP*AP*GP*TP*AP*GP*CP*AP*CP*GP*TP*GP*CP*TP*AP*CP*TP*C)-3'

Chain F:  17% 83%



- Molecule 1: 5'-D(*GP*AP*GP*TP*AP*GP*CP*AP*CP*GP*TP*GP*CP*TP*AP*CP*TP*C)-3'

Chain G:  44% 56%




- Molecule 1: 5'-D(*GP*AP*GP*TP*AP*GP*CP*AP*CP*GP*TP*GP*CP*TP*AP*CP*TP*C)-3'

Chain H:  50% 50%




- Molecule 1: 5'-D(*GP*AP*GP*TP*AP*GP*CP*AP*CP*GP*TP*GP*CP*TP*AP*CP*TP*C)-3'

Chain J:  61% 39%



- Molecule 2: MAD PROTEIN

Chain A:  65% 30% ..



- Molecule 2: MAD PROTEIN



- Molecule 3: MAX PROTEIN



- Molecule 3: MAX PROTEIN



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.35Å 56.02Å 65.60Å 88.82° 79.08° 67.11°	Depositor
Resolution (Å)	20.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.00)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.264 , 0.324	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4212	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	F	0.41	0/410	0.69	0/631
1	G	0.42	0/410	0.74	0/631
1	H	0.35	0/410	0.69	0/631
1	J	0.34	0/410	0.68	0/631
2	A	0.34	0/640	0.55	1/852 (0.1%)
2	D	0.34	0/615	0.51	0/821
3	B	0.46	0/646	0.52	0/862
3	E	0.35	0/626	0.49	0/837
All	All	0.38	0/4167	0.60	1/5896 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	32	PRO	N-CA-CB	5.69	110.13	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	366	0	204	19	0
1	G	366	0	204	13	0
1	H	366	0	204	12	0
1	J	366	0	204	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	634	0	652	39	0
2	D	609	0	633	40	0
3	B	637	0	632	36	0
3	E	617	0	606	32	0
4	A	36	0	0	1	0
4	B	38	0	0	5	0
4	D	41	0	0	3	0
4	E	28	0	0	1	0
4	F	26	0	0	0	0
4	G	38	0	0	0	0
4	H	23	0	0	0	0
4	J	21	0	0	0	0
All	All	4212	0	3339	176	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 176 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:109:DA:H2''	1:F:110:DC:H5''	1.49	0.94
1:F:105:DT:H2''	1:F:106:DA:C8	2.06	0.90
2:D:566:GLN:HE22	2:D:569:ARG:HH21	1.21	0.88
2:D:578:LEU:HD21	3:E:778:VAL:HG22	1.56	0.86
2:D:527:LEU:HD22	3:E:754:ARG:HH22	1.42	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	77/80 (96%)	69 (90%)	6 (8%)	2 (3%)	5	2
2	D	75/80 (94%)	71 (95%)	3 (4%)	1 (1%)	12	6
3	B	74/76 (97%)	72 (97%)	1 (1%)	1 (1%)	11	5
3	E	72/76 (95%)	70 (97%)	2 (3%)	0	100	100
All	All	298/312 (96%)	282 (95%)	12 (4%)	4 (1%)	12	6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	231	SER
2	A	30	LEU
2	A	28	VAL
2	D	529	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	67/75 (89%)	66 (98%)	1 (2%)	65	69
2	D	66/75 (88%)	65 (98%)	1 (2%)	65	69
3	B	68/69 (99%)	65 (96%)	3 (4%)	28	25
3	E	66/69 (96%)	64 (97%)	2 (3%)	41	41
All	All	267/288 (93%)	260 (97%)	7 (3%)	46	48

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	269	ARG
2	D	530	LEU
3	E	766	ASP
3	E	725	LEU
3	B	245	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14

such sidechains are listed below:

Mol	Chain	Res	Type
2	D	562	HIS
2	D	566	GLN
3	E	717	HIS
3	E	707	HIS
3	E	708	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.