



wwPDB X-ray Structure Validation Summary Report ⓘ

May 31, 2020 – 11:47 am BST

PDB ID : 6NLF
Title : 1.45 Å resolution structure of apo BfrB from *Pseudomonas aeruginosa*
Authors : Lovell, S.; Punchi-Hewage, A.; Battaile, K.P.; Yao, H.; Nammalwar, B.;
Gnanasekaran, K.K.; Bunce, R.A.; Reitz, A.B.; Rivera, M.
Deposited on : 2019-01-08
Resolution : 1.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	
1	B	158	
1	C	158	
1	D	158	
1	E	158	
1	F	158	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	158	<div><div></div><div>96%</div><div></div><div>..</div></div>
1	H	158	<div><div></div><div>96%</div><div></div><div>..</div></div>
1	I	158	<div><div></div><div>97%</div><div></div><div>..</div></div>
1	J	158	<div><div></div><div>94%</div><div>5%</div><div>.</div></div>
1	K	158	<div><div></div><div>93%</div><div>6%</div><div>.</div></div>
1	L	158	<div><div></div><div>94%</div><div></div><div>..</div></div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 18267 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	7	0
			1324	836	224	256	8			
1	B	156	Total	C	N	O	S	0	6	0
			1307	826	219	254	8			
1	C	156	Total	C	N	O	S	0	6	0
			1312	828	221	255	8			
1	D	156	Total	C	N	O	S	0	6	0
			1315	831	222	254	8			
1	E	156	Total	C	N	O	S	0	6	0
			1307	826	219	254	8			
1	F	156	Total	C	N	O	S	0	7	0
			1316	831	220	257	8			
1	G	156	Total	C	N	O	S	0	8	0
			1328	838	225	257	8			
1	H	156	Total	C	N	O	S	0	6	0
			1315	830	222	255	8			
1	I	156	Total	C	N	O	S	0	6	0
			1318	832	223	255	8			
1	J	156	Total	C	N	O	S	0	7	0
			1327	838	225	256	8			
1	K	156	Total	C	N	O	S	0	7	0
			1324	837	224	255	8			
1	L	156	Total	C	N	O	S	0	7	0
			1323	836	223	256	8			

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		
2	A	1	Total	K	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	K	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	203	Total	O	0	0
			203	203		
3	B	197	Total	O	0	0
			197	197		
3	C	211	Total	O	0	0
			211	211		
3	D	203	Total	O	0	0
			203	203		
3	E	213	Total	O	0	0
			213	213		
3	F	201	Total	O	0	0
			201	201		
3	G	184	Total	O	0	0
			184	184		
3	H	192	Total	O	0	0
			192	192		
3	I	214	Total	O	0	0
			214	214		
3	J	223	Total	O	0	0
			223	223		
3	K	191	Total	O	0	0
			191	191		
3	L	216	Total	O	0	0
			216	216		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ferroxidase

Chain A:  95% ..



- Molecule 1: Ferroxidase

Chain B:  96% ..



- Molecule 1: Ferroxidase

Chain C:  94% ..



- Molecule 1: Ferroxidase

Chain D:  96% ..




- Molecule 1: Ferroxidase

Chain E:  95% ..



- Molecule 1: Ferroxidase

Chain F:  96% ..




- Molecule 1: Ferroxidase

Chain G:  96% ..



- Molecule 1: Ferroxidase

Chain H:  96% ..



- Molecule 1: Ferroxidase

Chain I:  97% ..



- Molecule 1: Ferroxidase

Chain J:  94% 5% .



- Molecule 1: Ferroxidase

Chain K:  93% 6% .



- Molecule 1: Ferroxidase

Chain L:  94% ..



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	129.73Å 197.77Å 204.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.18 – 1.45 47.90 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.18-1.45) 99.9 (47.90-1.45)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.45Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.147 , 0.165 0.160 , 0.176	Depositor DCC
R_{free} test set	22925 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.206	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	18267	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1354	0.55	1/1823 (0.1%)
1	B	0.40	0/1334	0.56	1/1799 (0.1%)
1	C	0.40	0/1339	0.54	0/1805
1	D	0.42	0/1342	0.55	1/1808 (0.1%)
1	E	0.42	0/1337	0.55	0/1803
1	F	0.39	0/1346	0.54	0/1815
1	G	0.42	0/1359	0.55	0/1832
1	H	0.40	0/1342	0.54	0/1808
1	I	0.43	0/1345	0.56	0/1811
1	J	0.44	0/1357	0.56	0/1826
1	K	0.40	0/1355	0.54	0/1825
1	L	0.42	0/1353	0.56	0/1822
All	All	0.41	0/16163	0.55	3/21777 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	90	ASP	CB-CG-OD1	5.92	123.63	118.30
1	A	90	ASP	CB-CG-OD1	5.15	122.94	118.30
1	D	90	ASP	CB-CG-OD1	5.14	122.93	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1324	0	1298	3	0
1	B	1307	0	1261	2	0
1	C	1312	0	1272	7	0
1	D	1315	0	1280	2	0
1	E	1307	0	1266	3	0
1	F	1316	0	1271	2	0
1	G	1328	0	1286	4	0
1	H	1315	0	1281	2	0
1	I	1318	0	1290	1	0
1	J	1327	0	1307	9	0
1	K	1324	0	1292	5	0
1	L	1323	0	1293	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	203	0	0	2	1
3	B	197	0	0	1	1
3	C	211	0	0	4	1
3	D	203	0	0	2	0
3	E	213	0	0	0	0
3	F	201	0	0	1	1
3	G	184	0	0	2	0
3	H	192	0	0	0	1
3	I	214	0	0	0	1
3	J	223	0	0	6	0
3	K	191	0	0	1	0
3	L	216	0	0	2	4
All	All	18267	0	15397	45	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84[B]:GLN:NE2	3:A:301:HOH:O	2.19	0.75
1:J:50[B]:ASP:OD2	3:J:201:HOH:O	2.05	0.74
1:L:132:ASP:OD1	3:L:201:HOH:O	2.07	0.73
1:J:84[B]:GLN:NE2	3:J:203:HOH:O	2.22	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:34:ASP:OD1	3:K:201:HOH:O	2.11	0.69

The worst 5 of 6 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:367:HOH:O	3:C:367:HOH:O[3_454]	1.74	0.46
3:H:235:HOH:O	3:I:203:HOH:O[3_554]	1.80	0.40
3:F:368:HOH:O	3:L:339:HOH:O[7_444]	1.99	0.21
3:B:409:HOH:O	3:L:388:HOH:O[3_554]	2.04	0.16
3:L:204:HOH:O	3:L:308:HOH:O[3_554]	2.11	0.09

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/158 (102%)	160 (99%)	1 (1%)	0	100	100
1	B	160/158 (101%)	157 (98%)	3 (2%)	0	100	100
1	C	160/158 (101%)	157 (98%)	3 (2%)	0	100	100
1	D	160/158 (101%)	157 (98%)	3 (2%)	0	100	100
1	E	160/158 (101%)	158 (99%)	2 (1%)	0	100	100
1	F	161/158 (102%)	158 (98%)	3 (2%)	0	100	100
1	G	162/158 (102%)	160 (99%)	2 (1%)	0	100	100
1	H	160/158 (101%)	157 (98%)	3 (2%)	0	100	100
1	I	160/158 (101%)	158 (99%)	2 (1%)	0	100	100
1	J	161/158 (102%)	159 (99%)	2 (1%)	0	100	100
1	K	161/158 (102%)	158 (98%)	3 (2%)	0	100	100
1	L	161/158 (102%)	158 (98%)	3 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1927/1896 (102%)	1897 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/144 (101%)	145 (99%)	1 (1%)	84	65
1	B	141/144 (98%)	140 (99%)	1 (1%)	84	65
1	C	143/144 (99%)	142 (99%)	1 (1%)	84	65
1	D	143/144 (99%)	142 (99%)	1 (1%)	84	65
1	E	142/144 (99%)	142 (100%)	0	100	100
1	F	143/144 (99%)	142 (99%)	1 (1%)	84	65
1	G	145/144 (101%)	144 (99%)	1 (1%)	84	65
1	H	144/144 (100%)	143 (99%)	1 (1%)	84	65
1	I	145/144 (101%)	144 (99%)	1 (1%)	84	65
1	J	147/144 (102%)	147 (100%)	0	100	100
1	K	145/144 (101%)	144 (99%)	1 (1%)	84	65
1	L	145/144 (101%)	145 (100%)	0	100	100
All	All	1729/1728 (100%)	1720 (100%)	9 (0%)	88	75

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	121	LYS
1	K	121	LYS
1	H	121	LYS
1	C	84	GLN
1	G	121	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	156/158 (98%)	-0.30	0 100 100	11, 14, 23, 40	0
1	B	156/158 (98%)	-0.29	0 100 100	11, 14, 23, 38	0
1	C	156/158 (98%)	-0.30	0 100 100	11, 14, 24, 42	0
1	D	156/158 (98%)	-0.23	0 100 100	11, 14, 24, 41	0
1	E	156/158 (98%)	-0.30	0 100 100	10, 14, 22, 40	0
1	F	156/158 (98%)	-0.32	0 100 100	10, 14, 23, 39	0
1	G	156/158 (98%)	-0.30	0 100 100	10, 14, 22, 39	0
1	H	156/158 (98%)	-0.32	0 100 100	11, 14, 23, 39	0
1	I	156/158 (98%)	-0.28	0 100 100	10, 13, 22, 42	0
1	J	156/158 (98%)	-0.28	0 100 100	9, 13, 22, 38	0
1	K	156/158 (98%)	-0.29	0 100 100	10, 14, 24, 42	0
1	L	156/158 (98%)	-0.29	0 100 100	11, 14, 21, 39	0
All	All	1872/1896 (98%)	-0.29	0 100 100	9, 14, 23, 42	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	K	B	201	1/1	1.00	0.07	10,10,10,10	0
2	K	D	201	1/1	1.00	0.08	11,11,11,11	0
2	K	A	201	1/1	1.00	0.08	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.