



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 08:36 pm BST

PDB ID : 3NMW  
Title : Crystal structure of armadillo repeats domain of APC  
Authors : Zhang, Z.; Chen, L.; Gao, L.; Lin, K.; Wu, G.  
Deposited on : 2010-06-22  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 83%, yellow 8%, orange 6%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>83%</span> <span>8%</span> <span>6%</span> </div> </div>
1	B	354	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 82%, yellow 10%, orange 6%, grey 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>82%</span> <span>10%</span> <span>6%</span> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APC variant protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	Se	1	0	0
			2568	1599	464	479	12	14			
1	B	331	Total	C	N	O	S	Se	0	0	0
			2568	1599	464	479	12	14			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	398	MSE	-	expression tag	UNP Q4LE70
A	399	GLY	-	expression tag	UNP Q4LE70
A	400	HIS	-	expression tag	UNP Q4LE70
A	401	HIS	-	expression tag	UNP Q4LE70
A	402	HIS	-	expression tag	UNP Q4LE70
A	403	HIS	-	expression tag	UNP Q4LE70
A	404	HIS	-	expression tag	UNP Q4LE70
A	405	HIS	-	expression tag	UNP Q4LE70
A	406	MSE	-	expression tag	UNP Q4LE70
B	398	MSE	-	expression tag	UNP Q4LE70
B	399	GLY	-	expression tag	UNP Q4LE70
B	400	HIS	-	expression tag	UNP Q4LE70
B	401	HIS	-	expression tag	UNP Q4LE70
B	402	HIS	-	expression tag	UNP Q4LE70
B	403	HIS	-	expression tag	UNP Q4LE70
B	404	HIS	-	expression tag	UNP Q4LE70
B	405	HIS	-	expression tag	UNP Q4LE70
B	406	MSE	-	expression tag	UNP Q4LE70

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total	O	0	0
			25	25		
3	B	655	Total	O	0	0
			655	655		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: APC variant protein



- Molecule 1: APC variant protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.39Å 52.68Å 63.45Å 90.31° 89.95° 95.16°	Depositor
Resolution (Å)	50.00 – 1.60 32.92 – 1.60	Depositor EDS
% Data completeness (in resolution range)	94.9 (50.00-1.60) 93.7 (32.92-1.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.168 , 0.201 0.177 , 0.209	Depositor DCC
$R_{free}$ test set	4173 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 33.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.417 for -h,-k,l 0.015 for -k,-h,l 0.015 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5826	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.63	1/2594 (0.0%)	0.71	0/3481
1	B	0.63	0/2594	0.71	3/3481 (0.1%)
All	All	0.63	1/5188 (0.0%)	0.71	3/6962 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	553	TRP	CZ3-CH2	7.22	1.51	1.40

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	497	LEU	CA-CB-CG	8.13	133.99	115.30
1	B	497	LEU	CB-CG-CD1	-5.91	100.95	111.00
1	B	549	ARG	NE-CZ-NH2	-5.56	117.52	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	640	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2568	0	2588	36	3
1	B	2568	0	2588	39	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	25	0	0	5	0
3	B	655	0	0	10	5
All	All	5826	0	5176	74	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:698:LEU:HD23	1:A:701:MSE:HE1	1.37	1.07
1:A:698:LEU:HA	1:A:701:MSE:HE2	1.43	1.00
1:A:666:LEU:CD1	1:A:701:MSE:HE3	1.95	0.96
1:A:666:LEU:HD11	1:A:701:MSE:HE3	1.48	0.96
1:A:490:ASN:H	1:A:490:ASN:HD22	1.20	0.89
1:A:698:LEU:HD23	1:A:701:MSE:CE	2.02	0.87
1:A:691:ASN:ND2	3:A:755:HOH:O	1.82	0.87
1:B:553:TRP:NE1	1:B:554:ARG:HD2	1.91	0.84
1:B:690:ARG:O	1:B:690:ARG:CG	2.25	0.84
1:B:490:ASN:H	1:B:490:ASN:HD22	1.25	0.82
1:B:690:ARG:O	1:B:690:ARG:HG3	1.81	0.81
1:A:690:ARG:O	1:A:690:ARG:CG	2.28	0.80
1:B:445:GLN:OE1	3:B:388:HOH:O	1.99	0.80
1:B:660:ASN:HA	3:B:3:HOH:O	1.83	0.79
1:A:403:HIS:N	3:B:984:HOH:O	2.16	0.77
1:A:738:LYS:O	3:A:771:HOH:O	2.05	0.75
1:A:690:ARG:O	1:A:690:ARG:HG3	1.88	0.73
1:B:696:GLU:HG3	3:B:993:HOH:O	1.89	0.71
1:A:666:LEU:HD12	1:A:701:MSE:HE3	1.73	0.70
1:B:460:GLU:HG2	1:B:463:ARG:HH11	1.59	0.66
1:A:447:CYS:HB2	1:A:493:TYR:CZ	2.34	0.63

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:698:LEU:HA	1:B:701:MSE:HE3	1.80	0.63
1:B:460:GLU:HG2	1:B:463:ARG:NH1	2.15	0.62
1:B:467:ASN:HD21	1:B:515:ASN:HD21	1.48	0.61
1:B:467:ASN:ND2	1:B:515:ASN:HD21	2.00	0.59
1:A:698:LEU:HA	1:A:701:MSE:CE	2.27	0.58
1:A:460:GLU:HG2	1:A:463:ARG:NH2	2.19	0.57
1:B:698:LEU:HA	1:B:701:MSE:CE	2.34	0.57
1:B:553:TRP:CE2	1:B:554:ARG:HD2	2.40	0.57
1:B:447:CYS:HB2	1:B:493:TYR:CZ	2.41	0.56
1:A:708:LYS:HE3	3:B:897:HOH:O	2.05	0.56
1:B:603:LYS:HE3	1:B:641:ASN:O	2.06	0.56
1:B:498:ARG:HH11	1:B:532:GLN:HE22	1.54	0.55
1:B:727:ARG:NH2	3:B:366:HOH:O	2.32	0.55
1:A:490:ASN:N	1:A:490:ASN:HD22	1.93	0.54
1:B:490:ASN:N	1:B:490:ASN:HD22	2.01	0.53
1:B:698:LEU:HD23	1:B:701:MSE:CE	2.40	0.52
1:B:460:GLU:CG	1:B:463:ARG:NH1	2.73	0.51
1:A:444:HIS:HD2	3:A:763:HOH:O	1.94	0.50
1:B:553:TRP:NE1	1:B:554:ARG:CD	2.71	0.50
1:A:715:HIS:HD2	1:A:718:ILE:H	1.59	0.50
1:B:498:ARG:HH11	1:B:532:GLN:NE2	2.10	0.49
1:A:626:THR:OG1	1:A:626:THR:O	2.28	0.49
1:B:626:THR:OG1	1:B:626:THR:O	2.31	0.49
1:A:690:ARG:HG2	1:A:690:ARG:O	2.11	0.48
1:A:460:GLU:HG2	1:A:463:ARG:HH21	1.79	0.48
1:B:715:HIS:HD2	1:B:718:ILE:H	1.62	0.48
1:A:696:GLU:HB3	3:B:846:HOH:O	2.12	0.48
1:B:696:GLU:CG	3:B:993:HOH:O	2.57	0.48
1:B:407:LEU:O	1:B:411:GLU:HG3	2.15	0.47
1:A:447:CYS:HB3	1:A:448:PRO:HD3	1.96	0.46
1:B:690:ARG:O	1:B:690:ARG:HG2	2.13	0.46
1:A:691:ASN:CG	3:A:755:HOH:O	2.37	0.46
1:A:557:VAL:HG13	3:B:932:HOH:O	2.16	0.46
1:B:464:HIS:O	1:B:468:GLU:HG2	2.18	0.44
1:A:490:ASN:N	1:A:490:ASN:ND2	2.64	0.44
1:A:532:GLN:HB2	1:A:544:ILE:HG13	1.99	0.44
1:B:554:ARG:HG2	3:B:280:HOH:O	2.17	0.44
1:A:407:LEU:O	1:A:411:GLU:HG3	2.17	0.43
1:B:490:ASN:H	1:B:490:ASN:ND2	2.05	0.43
1:B:698:LEU:HD23	1:B:701:MSE:HE1	2.01	0.43
1:B:413:ILE:HD12	1:B:478:LEU:HD22	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:418:GLU:O	1:B:422:GLU:HG3	2.17	0.43
1:B:714:LYS:HD3	1:B:714:LYS:HA	1.88	0.43
1:A:715:HIS:CD2	1:A:718:ILE:H	2.36	0.43
1:A:460:GLU:CG	1:A:463:ARG:NH2	2.82	0.42
1:B:555:ALA:O	1:B:560:LYS:HE2	2.19	0.42
1:B:693:LYS:HE3	1:B:696:GLU:OE1	2.20	0.41
1:B:553:TRP:HE1	1:B:554:ARG:HH21	1.66	0.41
1:A:464:HIS:O	1:A:468:GLU:HG2	2.21	0.41
1:A:407:LEU:HD21	1:B:716:LYS:HD2	2.03	0.41
1:A:701:MSE:HB2	1:A:701:MSE:HE3	1.92	0.40
1:A:444:HIS:CD2	3:A:763:HOH:O	2.73	0.40
1:A:529:LEU:O	1:A:544:ILE:HG21	2.22	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:623:ARG:NH1	3:B:975:HOH:O[1_456]	1.83	0.37
1:A:623:ARG:NH1	3:B:832:HOH:O[1_456]	1.95	0.25
3:B:806:HOH:O	3:B:992:HOH:O[1_556]	1.98	0.22
3:B:305:HOH:O	3:B:849:HOH:O[1_565]	2.06	0.14
1:A:623:ARG:CD	3:B:975:HOH:O[1_456]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/354 (92%)	321 (98%)	6 (2%)	0	100	100
1	B	327/354 (92%)	324 (99%)	3 (1%)	0	100	100
All	All	654/708 (92%)	645 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/283 (99%)	266 (95%)	15 (5%)	22	5
1	B	281/283 (99%)	271 (96%)	10 (4%)	35	12
All	All	562/566 (99%)	537 (96%)	25 (4%)	28	8

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	407	LEU
1	A	428	GLU
1	A	447	CYS
1	A	460	GLU
1	A	469	LEU
1	A	472	LEU
1	A	490	ASN
1	A	536	GLU
1	A	557	VAL
1	A	623	ARG
1	A	626	THR
1	A	628	THR
1	A	640	ARG
1	A	663	GLN
1	A	690	ARG
1	B	407	LEU
1	B	428	GLU
1	B	460	GLU
1	B	472	LEU
1	B	490	ASN
1	B	497	LEU
1	B	536	GLU
1	B	626	THR
1	B	628	THR
1	B	690	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	412	GLN
1	A	444	HIS
1	A	490	ASN
1	A	542	GLN
1	A	663	GLN
1	A	695	GLN
1	A	709	ASN
1	A	715	HIS
1	B	467	ASN
1	B	490	ASN
1	B	532	GLN
1	B	659	ASN
1	B	695	GLN
1	B	709	ASN
1	B	715	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	2	-	4,4,4	0.16	0	6,6,6	0.40	0
2	SO4	B	1	-	4,4,4	0.11	0	6,6,6	0.29	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	317/354 (89%)	-0.40	3 (0%) 84 84	11, 17, 37, 55	16 (5%)
1	B	317/354 (89%)	-0.36	4 (1%) 77 77	10, 17, 36, 54	25 (7%)
All	All	634/708 (89%)	-0.38	7 (1%) 80 80	10, 17, 36, 55	41 (6%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	625	GLN	6.5
1	A	625	GLN	3.7
1	B	626	THR	3.1
1	B	403	HIS	2.9
1	A	403	HIS	2.8
1	A	404	HIS	2.6
1	B	435	LYS	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	2	5/5	0.93	0.09	31,41,43,44	5
2	SO4	B	1	5/5	0.95	0.09	27,34,39,43	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.