



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 06:10 am BST

PDB ID : 1NPB  
Title : Crystal structure of the fosfomycin resistance protein from transposon Tn2921  
Authors : Pakhomova, S.; Rife, C.L.; Armstrong, R.N.; Newcomer, M.E.  
Deposited on : 2003-01-17  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	141	86% 11% ..
1	B	141	79% 16% ..
1	C	141	84% 13% ..
1	D	141	79% 13% . .
1	E	141	87% 11% .
1	F	141	77% 16% . .

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called fosfomycin-resistance protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	1	0
			1118	715	189	210	4			
1	B	139	Total	C	N	O	S	0	0	0
			1100	705	183	207	5			
1	C	138	Total	C	N	O	S	0	0	0
			1101	705	184	208	4			
1	D	137	Total	C	N	O	S	0	0	0
			1093	702	183	203	5			
1	E	138	Total	C	N	O	S	0	1	0
			1097	703	184	205	5			
1	F	138	Total	C	N	O	S	0	0	0
			1101	705	184	208	4			

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total 6	C 3	O 3	0	0
3	D	1	Total 6	C 3	O 3	0	0
3	E	1	Total 6	C 3	O 3	0	0
3	F	1	Total 6	C 3	O 3	0	0
3	F	1	Total 6	C 3	O 3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total 54	O 54	0	0
4	B	67	Total 67	O 67	0	0
4	C	74	Total 74	O 74	0	0
4	D	52	Total 52	O 52	0	0
4	E	42	Total 42	O 42	0	0
4	F	59	Total 59	O 59	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: fosfomycin-resistance protein

Chain A: 




- Molecule 1: fosfomycin-resistance protein

Chain B: 





- Molecule 1: fosfomycin-resistance protein

Chain C: 



- Molecule 1: fosfomycin-resistance protein

Chain D: 




- Molecule 1: fosfomycin-resistance protein

Chain E: 



- Molecule 1: fosfomycin-resistance protein



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	208.60 Å   208.60 Å   136.36 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	12.00 – 2.50	Depositor
% Data completeness (in resolution range)	93.8 (12.00-2.50)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.183 , 0.230	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7050	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	0/1153	0.88	1/1570 (0.1%)
1	B	0.83	0/1131	0.89	1/1542 (0.1%)
1	C	0.83	0/1132	0.91	2/1542 (0.1%)
1	D	0.78	0/1124	0.86	2/1531 (0.1%)
1	E	0.79	1/1133 (0.1%)	0.85	2/1544 (0.1%)
1	F	0.80	0/1132	0.89	0/1542
All	All	0.81	1/6805 (0.0%)	0.88	8/9271 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	138	SER	CA-CB	5.17	1.60	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ASP	CB-CG-OD2	6.73	124.36	118.30
1	C	52	ASP	CB-CG-OD2	6.69	124.32	118.30
1	D	52	ASP	CB-CG-OD2	6.15	123.83	118.30
1	C	108	ASP	CB-CG-OD2	5.85	123.57	118.30
1	B	44	ASP	CB-CG-OD2	5.63	123.37	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1118	0	1061	6	0
1	B	1100	0	1037	12	0
1	C	1101	0	1048	7	0
1	D	1093	0	1043	12	0
1	E	1097	0	1042	2	0
1	F	1101	0	1048	16	0
2	A	10	0	0	1	0
2	B	5	0	0	0	0
2	C	10	0	0	1	0
2	D	10	0	0	0	0
2	E	10	0	0	0	0
2	F	5	0	0	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	D	6	0	8	0	0
3	E	6	0	8	1	0
3	F	12	0	16	2	0
4	A	54	0	0	2	0
4	B	67	0	0	3	0
4	C	74	0	0	1	0
4	D	52	0	0	1	0
4	E	42	0	0	2	0
4	F	59	0	0	1	0
All	All	7050	0	6335	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 54 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:45:LEU:HD22	1:D:5:LEU:HB2	1.58	0.86
1:F:31:HIS:CE1	1:F:41:THR:HB	2.11	0.85
1:F:31:HIS:HE1	1:F:41:THR:HB	1.46	0.80
1:D:1:MET:HA	4:D:550:HOH:O	1.83	0.78
1:C:17:LYS:HE2	4:C:569:HOH:O	1.85	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	132 (95%)	5 (4%)	2 (1%)	11	20
1	B	137/141 (97%)	132 (96%)	4 (3%)	1 (1%)	22	39
1	C	136/141 (96%)	133 (98%)	2 (2%)	1 (1%)	22	39
1	D	135/141 (96%)	131 (97%)	3 (2%)	1 (1%)	22	39
1	E	137/141 (97%)	132 (96%)	4 (3%)	1 (1%)	22	39
1	F	136/141 (96%)	132 (97%)	3 (2%)	1 (1%)	22	39
All	All	820/846 (97%)	792 (97%)	21 (3%)	7 (1%)	17	31

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	98	GLU
1	F	98	GLU
1	A	139	ASP
1	B	138	SER
1	A	138	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/119 (98%)	105 (90%)	12 (10%)	7	14
1	B	114/119 (96%)	101 (89%)	13 (11%)	5	11
1	C	116/119 (98%)	106 (91%)	10 (9%)	10	20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	114/119 (96%)	101 (89%)	13 (11%)	5	11
1	E	115/119 (97%)	107 (93%)	8 (7%)	15	29
1	F	116/119 (98%)	98 (84%)	18 (16%)	2	4
All	All	692/714 (97%)	618 (89%)	74 (11%)	6	13

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	135	VAL
1	D	80	LEU
1	F	96	LYS
1	C	138	SER
1	D	17	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	35	ASN
1	C	56	GLN
1	F	31	HIS
1	B	35	ASN
1	E	94	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	F	506	-	5,5,5	0.39	0	5,5,5	0.97	0
2	SO4	F	408	-	4,4,4	0.38	0	6,6,6	0.58	0
3	GOL	D	504	-	5,5,5	0.55	0	5,5,5	0.83	0
2	SO4	D	407	-	4,4,4	0.20	0	6,6,6	0.41	0
2	SO4	E	410	-	4,4,4	0.43	0	6,6,6	1.37	1 (16%)
3	GOL	A	501	-	5,5,5	0.41	0	5,5,5	0.49	0
2	SO4	A	403	-	4,4,4	0.30	0	6,6,6	0.43	0
2	SO4	E	409	-	4,4,4	0.21	0	6,6,6	0.18	0
2	SO4	A	402	-	4,4,4	0.34	0	6,6,6	0.42	0
2	SO4	C	406	-	4,4,4	0.19	0	6,6,6	0.71	0
2	SO4	D	404	-	4,4,4	0.46	0	6,6,6	0.59	0
3	GOL	F	507	-	5,5,5	0.46	0	5,5,5	0.66	0
2	SO4	B	401	-	4,4,4	0.36	0	6,6,6	0.40	0
2	SO4	C	405	-	4,4,4	0.41	0	6,6,6	0.33	0
3	GOL	C	503	-	5,5,5	0.59	0	5,5,5	0.85	0
3	GOL	E	505	-	5,5,5	0.39	0	5,5,5	0.80	0
3	GOL	B	502	-	5,5,5	0.42	0	5,5,5	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	F	506	-	-	4/4/4/4	-
3	GOL	B	502	-	-	0/4/4/4	-
3	GOL	A	501	-	-	2/4/4/4	-
3	GOL	D	504	-	-	4/4/4/4	-
3	GOL	F	507	-	-	4/4/4/4	-
3	GOL	E	505	-	-	3/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	503	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	410	SO4	O3-S-O2	2.61	122.95	109.31

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	506	GOL	O1-C1-C2-C3
3	F	506	GOL	C1-C2-C3-O3
3	F	506	GOL	O2-C2-C3-O3
3	D	504	GOL	O1-C1-C2-C3
3	D	504	GOL	C1-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	403	SO4	1	0
3	F	507	GOL	2	0
2	C	405	SO4	1	0
3	E	505	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.