



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2020 – 11:00 pm BST

PDB ID : 2NPP  
Title : Structure of the Protein Phosphatase 2A Holoenzyme  
Authors : Xu, Y.; Chen, Y.; Xing, Y.; Chao, Y.; Shi, Y.  
Deposited on : 2006-10-28  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	589	
1	D	589	
2	B	449	
2	E	449	
3	C	309	
3	F	309	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	X	7	 71% 29%
4	Y	7	 43% 57%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein Phosphatase 2, regulatory subunit A (PR 65), alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	0	0	0
			4535	2881	764	863	27			
1	D	582	Total	C	N	O	S	0	0	0
			4535	2881	764	863	27			

- Molecule 2 is a protein called Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit gamma isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	388	Total	C	N	O	S	0	0	0
			3130	2043	513	560	14			
2	E	388	Total	C	N	O	S	0	0	0
			3130	2043	513	560	14			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	433	VAL	-	CLONING ARTIFACT	UNP Q13362
B	434	LEU	-	CLONING ARTIFACT	UNP Q13362
B	435	LYS	-	CLONING ARTIFACT	UNP Q13362
B	436	LYS	-	CLONING ARTIFACT	UNP Q13362
B	437	ARG	-	CLONING ARTIFACT	UNP Q13362
B	438	ILE	-	CLONING ARTIFACT	UNP Q13362
B	439	THR	-	CLONING ARTIFACT	UNP Q13362
E	433	VAL	-	CLONING ARTIFACT	UNP Q13362
E	434	LEU	-	CLONING ARTIFACT	UNP Q13362
E	435	LYS	-	CLONING ARTIFACT	UNP Q13362
E	436	LYS	-	CLONING ARTIFACT	UNP Q13362
E	437	ARG	-	CLONING ARTIFACT	UNP Q13362
E	438	ILE	-	CLONING ARTIFACT	UNP Q13362
E	439	THR	-	CLONING ARTIFACT	UNP Q13362

- Molecule 3 is a protein called Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	308	Total	C	N	O	S	0	0	0
			2497	1579	431	472	15			
3	F	308	Total	C	N	O	S	0	0	0
			2497	1579	431	472	15			

- Molecule 4 is a protein called microcystin LR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	X	7	Total	C	N	O	0	0	0
			69	49	10	10			
4	Y	7	Total	C	N	O	0	0	0
			69	49	10	10			


- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	2	Total	Mn	0	0
			2	2		
5	F	2	Total	Mn	0	0
			2	2		


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein Phosphatase 2, regulatory subunit A (PR 65), alpha isoform




- Molecule 1: Protein Phosphatase 2, regulatory subunit A (PR 65), alpha isoform








- Molecule 3: Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform



- Molecule 3: Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform







● Molecule 4: microcystin LR



● Molecule 4: microcystin LR



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.26Å 159.05Å 269.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 3.30 49.33 – 3.31	Depositor EDS
% Data completeness (in resolution range)	98.4 (100.00-3.30) 99.3 (49.33-3.31)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 3.33Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.255 , 0.299 0.225 , 0.267	Depositor DCC
$R_{free}$ test set	3549 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	93.9	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 72.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	20466	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACB, DAL, DAM, MN, 1ZN, FGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/4609	0.64	0/6257
1	D	0.45	0/4609	0.66	0/6257
2	B	0.42	0/3212	0.66	2/4359 (0.0%)
2	E	0.42	0/3212	0.66	3/4359 (0.1%)
3	C	0.40	0/2559	0.65	1/3469 (0.0%)
3	F	0.41	0/2559	0.66	0/3469
4	X	0.43	0/17	0.78	0/19
4	Y	0.48	0/17	0.69	0/19
All	All	0.42	0/20794	0.65	6/28208 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	162	ASP	N-CA-C	6.21	127.77	111.00
2	B	162	ASP	N-CA-C	6.08	127.41	111.00
2	B	49	PRO	N-CA-CB	5.29	109.65	103.30
3	C	128	GLY	N-CA-C	5.16	126.01	113.10
2	E	49	PRO	N-CA-CB	5.16	109.50	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4535	0	4637	292	0
1	D	4535	0	4637	282	0
2	B	3130	0	3050	258	0
2	E	3130	0	3050	253	0
3	C	2497	0	2393	171	0
3	F	2497	0	2393	161	0
4	X	69	0	68	3	0
4	Y	69	0	68	6	0
5	C	2	0	0	0	0
5	F	2	0	0	0	0
All	All	20466	0	20296	1397	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 1397 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:301:THR:HG22	3:C:302:ARG:H	1.27	0.99
3:F:174:ILE:HD13	3:F:180:ILE:HG12	1.48	0.96
3:F:301:THR:HG22	3:F:302:ARG:H	1.30	0.95
2:E:273:GLU:HA	2:E:313:ILE:HD11	1.46	0.93
2:B:322:ALA:HB2	2:B:360:MET:HE1	1.51	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	580/589 (98%)	469 (81%)	94 (16%)	17 (3%)	4	24
1	D	580/589 (98%)	481 (83%)	85 (15%)	14 (2%)	6	28

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	386/449 (86%)	273 (71%)	78 (20%)	35 (9%)	1	4
2	E	386/449 (86%)	277 (72%)	72 (19%)	37 (10%)	0	4
3	C	306/309 (99%)	245 (80%)	44 (14%)	17 (6%)	2	11
3	F	306/309 (99%)	250 (82%)	42 (14%)	14 (5%)	2	15
4	X	1/7 (14%)	1 (100%)	0	0	100	100
4	Y	1/7 (14%)	1 (100%)	0	0	100	100
All	All	2546/2708 (94%)	1997 (78%)	415 (16%)	134 (5%)	2	12

5 of 134 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	25	VAL
1	A	391	VAL
2	B	43	PHE
2	B	45	PHE
2	B	55	TRP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	509/512 (99%)	474 (93%)	35 (7%)	15	44
1	D	509/512 (99%)	480 (94%)	29 (6%)	20	51
2	B	331/414 (80%)	309 (93%)	22 (7%)	16	46
2	E	331/414 (80%)	306 (92%)	25 (8%)	13	39
3	C	273/274 (100%)	260 (95%)	13 (5%)	25	56
3	F	273/274 (100%)	260 (95%)	13 (5%)	25	56
4	X	2/2 (100%)	2 (100%)	0	100	100
4	Y	2/2 (100%)	2 (100%)	0	100	100
All	All	2230/2404 (93%)	2093 (94%)	137 (6%)	18	48

5 of 137 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	119	GLU
1	D	122	SER
3	F	43	SER
3	C	211	ILE
3	C	294	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	44	ASN
1	D	200	ASN
2	E	392	GLN
1	D	22	ASN
1	D	230	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DAM	Y	7	3,4	4,5,6	1.63	1 (25%)	3,5,7	3.99	3 (100%)
4	1ZN	X	5	4	23,23,24	1.09	2 (8%)	24,29,31	0.85	0
4	FGA	X	6	4	6,6,9	1.19	0	5,6,11	0.47	0
4	ACB	X	3	-	5,8,9	3.65	3 (60%)	4,10,12	0.76	0
4	ACB	Y	3	-	5,8,9	3.07	2 (40%)	4,10,12	0.88	0
4	DAM	X	7	3,4	4,5,6	1.90	1 (25%)	3,5,7	4.10	3 (100%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FGA	Y	6	4	6,6,9	1.27	1 (16%)	5,6,11	0.62	0
4	1ZN	Y	5	4	23,23,24	1.16	2 (8%)	24,29,31	0.82	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DAM	Y	7	3,4	-	0/0/4/6	-
4	1ZN	X	5	4	-	1/22/25/27	0/1/1/1
4	FGA	X	6	4	-	1/3/4/9	-
4	ACB	X	3	-	-	1/5/10/12	-
4	ACB	Y	3	-	-	1/5/10/12	-
4	DAM	X	7	3,4	-	0/0/4/6	-
4	FGA	Y	6	4	-	1/3/4/9	-
4	1ZN	Y	5	4	-	1/22/25/27	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	X	3	ACB	CA-N	6.53	1.60	1.47
4	Y	3	ACB	CA-N	6.15	1.59	1.47
4	X	3	ACB	CB-CA	4.22	1.59	1.55
4	X	7	DAM	C-CA	3.27	1.50	1.45
4	Y	7	DAM	C-CA	2.67	1.49	1.45

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	X	7	DAM	O-C-CA	-5.31	118.41	125.22
4	Y	7	DAM	O-C-CA	-4.87	118.97	125.22
4	Y	7	DAM	CM-N-CA	-4.45	116.68	123.45
4	X	7	DAM	CM-N-CA	-4.11	117.20	123.45
4	X	7	DAM	CB-CA-N	-2.29	120.35	125.91

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	X	3	ACB	CA-CB-CG-OD1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	Y	3	ACB	CA-CB-CG-OD1
4	X	6	FGA	CA-CB-CG-CD
4	X	5	1ZN	C10-C2-C3-C4
4	Y	5	1ZN	C10-C2-C3-C4

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Y	7	DAM	3	0
4	X	5	1ZN	1	0
4	X	7	DAM	2	0
4	Y	6	FGA	1	0
4	Y	5	1ZN	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:



Mol	Chain	Number of breaks
4	X	2
4	Y	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	X	6:FGA	C	7:DAM	N	5.35
1	Y	6:FGA	C	7:DAM	N	5.26
1	X	3:ACB	C	4:ARG	N	3.49
1	Y	3:ACB	C	4:ARG	N	3.45

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	582/589 (98%)	-0.11	18 (3%) 49 48	25, 65, 112, 147	0
1	D	582/589 (98%)	-0.03	9 (1%) 73 72	22, 53, 109, 145	0
2	B	388/449 (86%)	0.12	22 (5%) 23 23	31, 81, 140, 155	0
2	E	388/449 (86%)	-0.02	12 (3%) 49 48	28, 75, 136, 155	0
3	C	308/309 (99%)	0.04	6 (1%) 66 65	26, 68, 110, 168	0
3	F	308/309 (99%)	0.03	7 (2%) 60 59	27, 64, 109, 169	0
4	X	2/7 (28%)	0.46	0 100 100	91, 91, 91, 103	0
4	Y	2/7 (28%)	-0.12	0 100 100	76, 76, 76, 96	0
All	All	2560/2708 (94%)	-0.01	74 (2%) 51 50	22, 65, 127, 169	0

The worst 5 of 74 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	73	ASN	5.3
2	B	130	PHE	4.9
2	B	71	THR	4.3
3	F	170	LEU	4.2
2	B	77	ILE	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	FGA	X	6	7/10	0.79	0.24	81,88,93,97	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FGA	Y	6	7/10	0.85	0.24	77,81,87,93	0
4	1ZN	X	5	23/24	0.92	0.47	64,70,89,93	0
4	ACB	X	3	9/10	0.93	0.25	93,97,99,101	0
4	1ZN	Y	5	23/24	0.93	0.32	64,74,82,84	0
4	DAL	X	1	5/6	0.94	0.20	77,77,80,80	0
4	DAM	Y	7	6/7	0.95	0.16	72,77,79,81	0
4	ACB	Y	3	9/10	0.95	0.25	80,85,92,93	0
4	DAM	X	7	6/7	0.96	0.22	76,80,81,82	0
4	DAL	Y	1	5/6	0.97	0.14	72,72,73,74	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MN	C	502	1/1	0.94	0.16	65,65,65,65	0
5	MN	F	502	1/1	0.97	0.20	56,56,56,56	0
5	MN	C	501	1/1	0.97	0.18	46,46,46,46	0
5	MN	F	501	1/1	0.98	0.15	35,35,35,35	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.