



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 08:19 pm BST

PDB ID : 3NPL
Title : Structure of Ru(bpy)2(A-Phen)(K97C) P450 BM3 heme domain, a ruthenium modified P450 BM3 mutant
Authors : Ener, M.; Lee, Y.-T.; Goodin, D.B.; Winkler, J.R.; Gray, H.B.; Cheruzel, L.
Deposited on : 2010-06-28
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	470	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div>••</div> </div> </div>
1	B	470	<div> <div>3%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>•</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional P-450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	4	0
			3690	2357	628	689	16			
1	B	456	Total	C	N	O	S	0	2	0
			3676	2348	626	686	16			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	ALA	CYS	ENGINEERED MUTATION	UNP P14779
A	97	CYS	LYS	ENGINEERED MUTATION	UNP P14779
A	156	SER	CYS	ENGINEERED MUTATION	UNP P14779
A	464	HIS	-	EXPRESSION TAG	UNP P14779
A	465	HIS	-	EXPRESSION TAG	UNP P14779
A	466	HIS	-	EXPRESSION TAG	UNP P14779
A	467	HIS	-	EXPRESSION TAG	UNP P14779
A	468	HIS	-	EXPRESSION TAG	UNP P14779
A	469	HIS	-	EXPRESSION TAG	UNP P14779
B	62	ALA	CYS	ENGINEERED MUTATION	UNP P14779
B	97	CYS	LYS	ENGINEERED MUTATION	UNP P14779
B	156	SER	CYS	ENGINEERED MUTATION	UNP P14779
B	464	HIS	-	EXPRESSION TAG	UNP P14779
B	465	HIS	-	EXPRESSION TAG	UNP P14779
B	466	HIS	-	EXPRESSION TAG	UNP P14779
B	467	HIS	-	EXPRESSION TAG	UNP P14779
B	468	HIS	-	EXPRESSION TAG	UNP P14779
B	469	HIS	-	EXPRESSION TAG	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is bis(2,2'-bipyridine-kappa 2 N 1 ,N 1') [2-iodo-N-(1,10-phenanthrolin-

5-yl-kappa 2 N 1 ,N 10)acetamide]ruthenium(2+) (three-letter code: RU8) (formula: C₃₄H₂₆IN₇ORu).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	Ru	0	0
			43	34	7	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	227	Total	O	0	0
			227	227		
5	B	209	Total	O	0	0
			209	209		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional P-450/NADPH-P450 reductase



- Molecule 1: Bifunctional P-450/NADPH-P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	117.08Å 117.08Å 273.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.40 10.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.7 (10.00-2.40) 98.7 (10.00-2.40)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.69 (at 2.41Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.201 , 0.238 0.200 , 0.238	Depositor DCC
R_{free} test set	3677 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7936	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, RU8, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/3788	0.64	1/5122 (0.0%)
1	B	0.55	0/3768	0.64	1/5096 (0.0%)
All	All	0.56	0/7556	0.64	2/10218 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ASP	CB-CG-OD1	5.44	123.20	118.30
1	A	148	LEU	CA-CB-CG	5.09	127.02	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	46	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3690	0	3664	24	0
1	B	3676	0	3646	21	0
2	A	43	0	30	0	0
2	B	43	0	30	1	0
3	A	5	0	0	0	0
4	B	43	0	24	0	0
5	A	227	0	0	1	0
5	B	209	0	0	2	0
All	All	7936	0	7394	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:THR:HG22	1:B:24:LYS:H	1.55	0.71
1:B:49:THR:HG21	1:B:354:MET:HG2	1.72	0.71
1:A:102:ILE:HG22	1:A:103:LEU:HD13	1.73	0.69
1:B:217:ASP:OD1	1:B:255:ARG:NH1	2.28	0.64
1:B:148:LEU:HD11	1:B:413:VAL:HG21	1.80	0.63
1:B:330:ALA:HB1	1:B:354:MET:CE	2.28	0.63
1:B:330:ALA:HB1	1:B:354:MET:HE3	1.83	0.61
1:B:53:SER:HB3	1:B:359:GLN:HB3	1.81	0.60
1:B:110:GLN:HG3	5:B:807:HOH:O	2.07	0.54
1:B:261:PHE:O	1:B:265:GLY:HA3	2.10	0.52
1:B:391:LYS:NZ	1:B:395:ASN:OD1	2.42	0.52
1:A:192:ASN:N	1:A:193:PRO:HD3	2.26	0.51
1:B:49:THR:HG21	1:B:354:MET:CG	2.40	0.49
1:A:192:ASN:N	1:A:192:ASN:OD1	2.45	0.49
1:B:216:VAL:HG21	1:B:259:ILE:HG13	1.95	0.48
1:A:81:PHE:HB3	1:A:209:ILE:HG12	1.96	0.47
1:A:97:CYS:HB2	5:A:606:HOH:O	2.14	0.47
1:A:60:GLU:OE2	1:A:342:GLY:HA2	2.15	0.47
1:A:377:GLU:OE2	1:A:380:GLU:HG3	2.15	0.47
1:B:76:LYS:HE3	5:B:790:HOH:O	2.15	0.46
1:B:141:VAL:HB	1:B:142:PRO:HD3	1.97	0.46
1:A:21:ASN:HA	1:A:189:GLN:HG2	1.98	0.46
1:A:266:HIS:CG	1:A:267:GLU:N	2.84	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:330:ALA:HB1	1:B:354:MET:HE1	1.99	0.45
1:A:134:ASN:HB3	1:B:222:ASP:OD2	2.17	0.44
1:A:228:GLU:HG3	1:A:229:GLN:N	2.33	0.44
1:A:272:LEU:HD13	1:A:322:LEU:HG	1.98	0.44
1:A:300:ASP:HB3	1:A:301:PRO:CD	2.48	0.43
1:A:458:ILE:HA	1:A:459:PRO:HD2	1.83	0.43
1:A:158:PHE:CD1	1:A:234:LEU:HB2	2.53	0.42
1:B:81:PHE:HB3	1:B:209:ILE:HG12	2.00	0.42
1:A:301:PRO:HB2	1:A:456:GLY:HA3	2.01	0.42
1:A:7:GLN:HG3	1:A:41:LYS:O	2.19	0.42
1:B:112:MET:HE1	1:B:405:PHE:HA	2.02	0.41
1:A:253:ASN:O	1:A:257:GLN:HG2	2.21	0.41
1:A:58:ILE:HD12	1:A:360:LEU:HD13	2.03	0.41
1:A:20:LEU:HB2	1:A:189:GLN:NE2	2.36	0.41
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.51	0.41
1:A:216:VAL:HG21	1:A:259:ILE:HG13	2.01	0.41
1:A:49:THR:HG23	1:A:352[A]:GLU:CD	2.41	0.41
1:B:278:TYR:HA	1:B:444:PHE:CZ	2.55	0.41
1:A:385:ILE:HA	1:A:386:PRO:HD3	1.99	0.40
1:A:148:LEU:HD11	1:A:413:VAL:HG21	2.02	0.40
1:B:434:LYS:HB2	1:B:442:GLU:HB2	2.03	0.40
1:B:39:ILE:HA	1:B:51:TYR:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/470 (97%)	447 (98%)	10 (2%)	1 (0%)	47	62
1	B	456/470 (97%)	444 (97%)	12 (3%)	0	100	100
All	All	914/940 (97%)	891 (98%)	22 (2%)	1 (0%)	51	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	196	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/411 (98%)	394 (98%)	7 (2%)	60	78
1	B	399/411 (97%)	393 (98%)	6 (2%)	65	80
All	All	800/822 (97%)	787 (98%)	13 (2%)	62	79

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ASP
1	A	49	THR
1	A	97	CYS
1	A	103	LEU
1	A	182	ASP
1	A	188	LEU
1	A	380	GLU
1	B	31	LYS
1	B	94	LYS
1	B	228	GLU
1	B	267	GLU
1	B	381	ASN
1	B	383	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	500	1	27,50,50	2.12	9 (33%)	17,82,82	1.81	3 (17%)
3	SO4	A	550	-	4,4,4	0.17	0	6,6,6	0.18	0
4	RU8	B	501	1	46,52,53	4.41	32 (69%)	43,87,88	3.00	25 (58%)
2	HEM	B	500	1	27,50,50	2.13	7 (25%)	17,82,82	1.62	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	0/6/54/54	-
4	RU8	B	501	1	-	1/4/116/118	0/10/10/10
2	HEM	B	500	1	-	0/6/54/54	-

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	501	RU8	C32-N33	-11.06	1.27	1.49
4	B	501	RU8	C35-N34	-10.04	1.29	1.49
4	B	501	RU8	C45-N41	-9.87	1.29	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	501	RU8	C23-N22	-9.70	1.30	1.49
4	B	501	RU8	C29-C28	-6.13	1.40	1.53
4	B	501	RU8	C38-C39	-6.12	1.40	1.53
4	B	501	RU8	C26-C27	-5.77	1.40	1.53
4	B	501	RU8	C42-C40	-5.67	1.41	1.53
2	A	500	HEM	C3D-C2D	5.31	1.53	1.37
4	B	501	RU8	C30-C29	-5.27	1.39	1.53
4	B	501	RU8	C03-C05	5.27	1.42	1.36
2	B	500	HEM	C3D-C2D	5.17	1.53	1.37
4	B	501	RU8	C37-C38	-4.96	1.40	1.53
4	B	501	RU8	C43-C42	-4.76	1.40	1.53
4	B	501	RU8	O48-C47	4.74	1.33	1.23
4	B	501	RU8	C25-C26	-4.71	1.40	1.53
2	A	500	HEM	C3C-C2C	-4.34	1.34	1.40
4	B	501	RU8	C49-C47	-4.06	1.42	1.50
2	B	500	HEM	C3B-C2B	-3.90	1.35	1.40
4	B	501	RU8	C14-C11	3.87	1.52	1.43
2	A	500	HEM	C3B-CAB	3.79	1.55	1.47
2	A	500	HEM	C3B-C2B	-3.68	1.35	1.40
2	B	500	HEM	C3C-C2C	-3.62	1.35	1.40
2	B	500	HEM	C3B-CAB	3.60	1.55	1.47
4	B	501	RU8	C05-N46	-3.59	1.32	1.41
2	B	500	HEM	C3C-CAC	3.49	1.55	1.47
4	B	501	RU8	C15-C17	-3.48	1.30	1.38
4	B	501	RU8	C35-C36	-3.26	1.39	1.51
2	A	500	HEM	C3C-CAC	3.22	1.54	1.47
4	B	501	RU8	C40-C39	-3.21	1.43	1.52
4	B	501	RU8	C32-C31	-3.21	1.39	1.51
4	B	501	RU8	C23-C24	-3.02	1.40	1.51
4	B	501	RU8	C37-C36	-2.99	1.39	1.51
4	B	501	RU8	C31-C30	-2.98	1.39	1.51
4	B	501	RU8	C28-C27	-2.91	1.44	1.52
2	B	500	HEM	CAA-C2A	2.84	1.56	1.52
4	B	501	RU8	C45-C44	-2.83	1.41	1.51
4	B	501	RU8	C44-C43	-2.83	1.40	1.51
4	B	501	RU8	C14-N13	-2.71	1.31	1.37
4	B	501	RU8	C25-C24	-2.65	1.41	1.51
4	B	501	RU8	C08-N10	-2.38	1.29	1.33
2	A	500	HEM	CAA-C2A	2.34	1.55	1.52
4	B	501	RU8	C47-N46	-2.24	1.31	1.36
4	B	501	RU8	C11-N10	-2.22	1.32	1.37
2	A	500	HEM	C1D-ND	2.10	1.40	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	CMB-C2B	2.06	1.56	1.51
2	B	500	HEM	C1D-ND	2.04	1.40	1.36
2	A	500	HEM	CMC-C2C	2.01	1.56	1.51

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	RU8	C05-C21-C14	5.29	121.27	118.25
2	A	500	HEM	CBD-CAD-C3D	-4.98	103.29	112.48
4	B	501	RU8	C44-C45-N41	4.84	120.88	113.00
4	B	501	RU8	C35-C36-C37	4.62	119.26	111.44
4	B	501	RU8	C43-C42-C40	4.57	119.45	111.22
4	B	501	RU8	C32-C31-C30	4.48	119.02	111.44
4	B	501	RU8	C03-C05-C21	-4.31	117.49	120.50
4	B	501	RU8	C45-C44-C43	4.23	118.61	111.44
4	B	501	RU8	C24-C23-N22	4.08	119.65	113.00
4	B	501	RU8	C25-C26-C27	4.07	118.56	111.22
4	B	501	RU8	C23-C24-C25	3.94	118.11	111.44
4	B	501	RU8	C31-C32-N33	3.93	119.41	113.00
4	B	501	RU8	C36-C37-C38	3.79	119.14	111.42
4	B	501	RU8	C37-C38-C39	3.75	117.98	111.22
4	B	501	RU8	C24-C25-C26	3.75	119.06	111.42
4	B	501	RU8	C30-C29-C28	3.72	117.93	111.22
4	B	501	RU8	C21-C05-N46	3.70	124.60	118.36
4	B	501	RU8	C36-C35-N34	3.62	118.90	113.00
4	B	501	RU8	C31-C30-C29	3.56	118.67	111.42
2	B	500	HEM	CBD-CAD-C3D	-3.42	106.17	112.48
4	B	501	RU8	C49-C47-N46	3.40	119.97	114.98
4	B	501	RU8	C44-C43-C42	3.12	117.77	111.42
2	B	500	HEM	CMA-C3A-C4A	-2.86	124.07	128.46
2	B	500	HEM	C1D-C2D-C3D	-2.57	105.21	107.00
4	B	501	RU8	C14-C11-N10	2.50	119.94	116.56
2	A	500	HEM	C1D-C2D-C3D	-2.47	105.28	107.00
4	B	501	RU8	C02-C11-C14	-2.43	116.76	119.86
4	B	501	RU8	O48-C47-C49	-2.32	117.75	122.06
2	A	500	HEM	C4C-C3C-C2C	2.32	108.52	106.90
4	B	501	RU8	C15-N13-C14	2.20	122.33	117.51
4	B	501	RU8	C08-N10-C11	2.13	122.18	117.51
2	B	500	HEM	CMC-C2C-C3C	2.13	128.66	124.68

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	501	RU8	C03-C05-N46-C47

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	500	HEM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand RU8 B 501




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/470 (97%)	-0.43	16 (3%) 44 43	22, 33, 60, 82	0
1	B	456/470 (97%)	-0.42	12 (2%) 56 54	24, 35, 55, 68	0
All	All	912/940 (97%)	-0.43	28 (3%) 49 47	22, 35, 58, 82	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	383	SER	5.8
1	A	460	SER	4.9
1	A	229	GLN	3.8
1	B	460	SER	3.7
1	A	227	GLY	3.7
1	A	193	PRO	3.4
1	A	196	PRO	3.4
1	A	228	GLU	3.3
1	A	191	ALA	3.2
1	B	382	PRO	3.1
1	A	285	HIS	3.1
1	B	229	GLN	3.0
1	B	381	ASN	2.7
1	A	203	ARG	2.6
1	A	46	GLY	2.5
1	B	228	GLU	2.5
1	A	194	ASP	2.5
1	A	110	GLN	2.5
1	B	227	GLY	2.4
1	A	381	ASN	2.3
1	B	343	GLY	2.3
1	A	383	SER	2.3
1	B	384	ALA	2.2
1	A	198	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	192	ASN	2.2
1	B	285	HIS	2.1
1	B	46	GLY	2.0
1	B	459	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	RU8	B	501	43/44	0.95	0.17	40,44,46,47	0
3	SO4	A	550	5/5	0.98	0.17	57,57,58,59	0
2	HEM	A	500	43/43	0.98	0.10	18,23,26,28	0
2	HEM	B	500	43/43	0.98	0.09	18,22,27,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around RU8 B 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.