



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 11:32 PM BST

PDB ID : 5NQ9
Title : Crystal structure of laccases from Pycnoporus sanguineus, izoform II, monoclinic
Authors : Orlikowska, M.; de J.Rostro-Alanis, M.; Bujacz, A.; Hernandez-Luna, C.; Rubio, R.; Parra, R.; Bujacz, G.
Deposited on : 2017-04-19
Resolution : 2.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> % 83% 11% .. </div> </div>
1	C	518	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> % 85% 10% .. </div> </div>
2	B	5	<div style="width: 100%; height: 10px; background-color: yellow; position: relative;"> 40% 60% </div>
3	D	2	<div style="width: 100%; height: 10px; background-color: yellow; position: relative;"> 100% </div>
3	F	2	<div style="width: 100%; height: 10px; background-color: green; position: relative;"> 100% </div>
4	E	3	<div style="width: 100%; height: 10px; background-color: yellow; position: relative;"> 100% </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	G	3	 67% 33%

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 7992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Laccase, izoform II.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	497	Total	C	N	O	S	0	0	0
			3803	2422	642	729	10			
1	C	497	Total	C	N	O	S	0	0	0
			3803	2422	642	729	10			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	G	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	6	Total	Cu	0	0
			6	6		
5	C	6	Total	Cu	0	0
			6	6		

- Molecule 6 is PEROXIDE ION (three-letter code: PER) (formula: O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O	0	0
			2	2		
6	C	1	Total	O	0	0
			2	2		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			3	1	2		
8	A	1	Total	C	O	0	0
			3	1	2		
8	A	1	Total	C	O	0	0
			3	1	2		
8	A	1	Total	C	O	0	0
			3	1	2		
8	A	1	Total	C	O	0	0
			3	1	2		
8	A	1	Total	C	O	0	0
			3	1	2		
8	C	1	Total	C	O	0	0
			3	1	2		
8	C	1	Total	C	O	0	0
			3	1	2		
8	C	1	Total	C	O	0	0
			3	1	2		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	38	Total	O	0	0
			38	38		
10	C	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Laccase, izoform II



- Molecule 1: Laccase, izoform II



- Molecule 2: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

UAG1
UAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

UAG1
UAG2

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

UAG1
UAG2
BMA3

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  67%  33%

UAG1
UAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.46 Å 61.96 Å 103.31 Å 90.00° 109.82° 90.00°	Depositor
Resolution (Å)	48.65 – 2.72 48.65 – 2.72	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.65-2.72) 99.4 (48.65-2.72)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.73 Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.170 , 0.246 0.176 , 0.249	Depositor DCC
R_{free} test set	1462 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	55.7	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7992	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, FMT, PER, PEG, CU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/3924	0.83	0/5394
1	C	0.78	0/3924	0.82	0/5394
All	All	0.78	0/7848	0.83	0/10788

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3803	0	3601	27	0
1	C	3803	0	3601	21	0
2	B	61	0	52	0	0
3	D	28	0	25	0	0
3	F	28	0	25	0	0
4	E	39	0	34	0	0
4	G	39	0	34	0	0
5	A	6	0	0	0	0
5	C	6	0	0	0	0
6	A	2	0	0	0	0
6	C	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	28	0	40	0	0
7	C	21	0	30	0	0
8	A	21	0	7	0	0
8	C	12	0	4	0	0
9	C	14	0	13	0	0
10	A	38	0	0	0	0
10	C	41	0	0	0	0
All	All	7992	0	7466	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:THR:HG22	1:A:136:GLN:OE1	1.76	0.84
1:C:389:GLU:N	1:C:389:GLU:OE2	2.19	0.72
1:A:198:SER:HG	1:A:201:THR:HG1	1.36	0.71
1:A:243:GLU:HB3	1:A:266:SER:HB2	1.81	0.63
1:A:227:ASP:HB3	1:A:228:PRO:CD	2.30	0.62
1:C:171:ASP:HB2	1:C:187:SER:HB3	1.83	0.60
1:A:81:THR:HG22	1:A:109:ALA:HA	1.83	0.60
1:A:243:GLU:OE2	1:A:246:SER:HA	2.02	0.59
1:C:227:ASP:HB3	1:C:228:PRO:CD	2.33	0.58
1:A:54:ALA:HB1	1:A:129:TYR:OH	2.06	0.56
1:C:83:SER:OG	1:C:99:GLY:O	2.26	0.53
1:A:358:PHE:HB3	1:A:482:PHE:CD1	2.44	0.53
1:C:471:HIS:ND1	1:C:472:CYS:O	2.31	0.53
1:A:216:ARG:HG2	1:A:270:ASP:HA	1.92	0.52
1:C:489:ASP:CG	1:C:492:ASP:HB2	2.33	0.50
1:C:85:HIS:CE1	1:C:421:HIS:CE1	3.00	0.49
1:A:26:VAL:HG22	1:A:66:GLN:HB2	1.95	0.48
1:C:171:ASP:OD2	1:C:228:PRO:HD2	2.14	0.48
1:C:361:ASN:O	1:C:362:ASN:HB2	2.14	0.48
1:A:78:MET:O	1:A:79:LEU:HB2	2.14	0.47
1:A:171:ASP:HB2	1:A:187:SER:HB3	1.97	0.47
1:C:52:THR:HA	1:C:53:PRO:C	2.35	0.47
1:A:54:ALA:HB1	1:A:55:PRO:HD2	1.96	0.46
1:C:227:ASP:HB3	1:C:228:PRO:HD3	1.98	0.46
1:C:353:PHE:HA	1:C:357:ASN:O	2.16	0.45
1:C:82:THR:OG1	1:C:83:SER:N	2.48	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:ASP:OD2	1:A:226:CYS:HB2	2.17	0.44
1:A:494:THR:HG23	1:A:495:ALA:N	2.32	0.44
1:C:54:ALA:HB1	1:C:129:TYR:OH	2.18	0.44
1:C:67:LEU:HD13	1:C:86:TRP:CZ3	2.53	0.44
1:A:368:PRO:HB3	1:A:387:VAL:HG23	2.00	0.44
1:C:87:HIS:CE1	1:C:262:ALA:HB1	2.53	0.43
1:A:259:ILE:HG23	1:A:259:ILE:O	2.19	0.43
1:A:46:VAL:HG11	1:A:141:LEU:HB2	2.00	0.43
1:A:58:SER:HA	1:A:146:VAL:O	2.20	0.42
1:C:184:PRO:HG2	1:C:476:PHE:CZ	2.55	0.42
1:C:467:PRO:HA	1:C:487:ALA:HA	2.02	0.41
1:A:261:ALA:O	1:A:262:ALA:HB3	2.20	0.41
1:C:105:GLN:HG3	1:C:106:CYS:O	2.20	0.41
1:A:477:HIS:O	1:A:478:LEU:C	2.56	0.41
1:A:490:THR:N	1:A:491:PRO:CD	2.83	0.41
1:A:154:GLN:NE2	1:A:249:THR:O	2.54	0.41
1:C:31:LEU:CD2	1:C:48:VAL:HG22	2.51	0.41
1:A:57:ILE:O	1:A:145:PHE:HA	2.20	0.41
1:A:429:ARG:HD2	1:A:436:TYR:CE1	2.55	0.41
1:A:303:LEU:HD23	1:A:304:GLY:N	2.36	0.41
1:A:82:THR:CG2	1:A:136:GLN:OE1	2.60	0.40
1:C:397:SER:O	1:C:399:SER:OG	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	495/518 (96%)	462 (93%)	31 (6%)	2 (0%)	34	58
1	C	495/518 (96%)	458 (92%)	35 (7%)	2 (0%)	34	58
All	All	990/1036 (96%)	920 (93%)	66 (7%)	4 (0%)	34	58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	ASP
1	C	227	ASP
1	A	79	LEU
1	C	79	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/434 (96%)	394 (94%)	23 (6%)	21	44
1	C	417/434 (96%)	397 (95%)	20 (5%)	25	51
All	All	834/868 (96%)	791 (95%)	43 (5%)	23	47

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	LEU
1	A	82	THR
1	A	83	SER
1	A	113	SER
1	A	137	TYR
1	A	145	PHE
1	A	163	ASP
1	A	202	THR
1	A	233	SER
1	A	246	SER
1	A	289	THR
1	A	297	SER
1	A	316	THR
1	A	319	LYS
1	A	369	SER
1	A	378	SER
1	A	397	SER
1	A	399	SER
1	A	469	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	475	ASP
1	A	482	PHE
1	A	490	THR
1	A	501	GLN
1	C	23	ILE
1	C	28	ASP
1	C	94	THR
1	C	151	ASN
1	C	179	LEU
1	C	209	ILE
1	C	242	ILE
1	C	285	SER
1	C	289	THR
1	C	315	THR
1	C	319	LYS
1	C	369	SER
1	C	391	SER
1	C	397	SER
1	C	408	THR
1	C	415	PRO
1	C	462	THR
1	C	469	PHE
1	C	490	THR
1	C	492	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.31	0	17,19,21	1.89	3 (17%)
2	NAG	B	2	2	14,14,15	0.39	0	17,19,21	1.23	3 (17%)
2	MAN	B	3	2	11,11,12	0.30	0	15,15,17	1.43	2 (13%)
2	MAN	B	4	2	11,11,12	0.27	0	15,15,17	0.84	0
2	MAN	B	5	2	11,11,12	0.30	0	15,15,17	0.62	0
3	NAG	D	1	1,3	14,14,15	0.31	0	17,19,21	1.09	1 (5%)
3	NAG	D	2	3	14,14,15	0.31	0	17,19,21	0.82	1 (5%)
4	NAG	E	1	1,4	14,14,15	0.38	0	17,19,21	0.69	1 (5%)
4	NAG	E	2	4	14,14,15	0.41	0	17,19,21	0.95	1 (5%)
4	BMA	E	3	4	11,11,12	0.36	0	15,15,17	1.00	1 (6%)
3	NAG	F	1	1,3	14,14,15	0.28	0	17,19,21	0.81	0
3	NAG	F	2	3	14,14,15	0.34	0	17,19,21	0.75	0
4	NAG	G	1	1,4	14,14,15	0.31	0	17,19,21	0.63	0
4	NAG	G	2	4	14,14,15	0.30	0	17,19,21	0.77	0
4	BMA	G	3	4	11,11,12	0.23	0	15,15,17	1.15	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	B	2	2	-	1/6/23/26	0/1/1/1
2	MAN	B	3	2	-	1/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	1/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	E	2	4	-	2/6/23/26	0/1/1/1
4	BMA	E	3	4	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	6.09	120.45	112.19
2	B	3	MAN	C1-C2-C3	3.58	114.07	109.67
2	B	3	MAN	C1-O5-C5	3.22	116.55	112.19
4	E	3	BMA	C1-C2-C3	2.94	113.28	109.67
2	B	1	NAG	O5-C1-C2	2.65	115.47	111.29
2	B	2	NAG	C1-C2-N2	-2.61	106.04	110.49
3	D	1	NAG	C1-O5-C5	2.60	115.72	112.19
4	G	3	BMA	C1-O5-C5	2.49	115.57	112.19
4	G	3	BMA	O5-C1-C2	-2.48	106.95	110.77
2	B	1	NAG	C3-C4-C5	2.45	114.61	110.24
3	D	2	NAG	O5-C1-C2	-2.37	107.55	111.29
4	E	2	NAG	C4-C3-C2	2.34	114.44	111.02
4	E	1	NAG	C1-C2-N2	-2.30	106.56	110.49
2	B	2	NAG	C4-C3-C2	2.29	114.38	111.02
2	B	2	NAG	O5-C5-C6	2.04	110.41	107.20

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	2	NAG	O5-C5-C6-O6
3	F	1	NAG	O5-C5-C6-O6
4	E	3	BMA	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
4	E	3	BMA	O5-C5-C6-O6
3	F	1	NAG	C4-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6
4	E	2	NAG	O5-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6
4	G	2	NAG	O5-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	G	3	BMA	C4-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
4	E	2	NAG	C4-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6
2	B	3	MAN	O5-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6
4	G	3	BMA	O5-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6
4	E	1	NAG	C4-C5-C6-O6



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry

Of 33 ligands modelled in this entry, 12 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	PEG	A	619	-	6,6,6	0.64	0	5,5,5	0.45	0
7	PEG	C	612	-	6,6,6	0.68	0	5,5,5	0.64	0
6	PER	C	601	5	0,1,1	0.00	-	-		
8	FMT	A	624	-	0,2,2	0.00	-	0,1,1	0.00	-
9	NAG	C	606	1	14,14,15	0.60	0	17,19,21	1.00	1 (5%)
8	FMT	A	620	-	0,2,2	0.00	-	0,1,1	0.00	-
7	PEG	C	613	-	6,6,6	0.51	0	5,5,5	0.19	0
7	PEG	A	617	-	6,6,6	0.60	0	5,5,5	0.38	0
8	FMT	C	617	-	0,2,2	0.00	-	0,1,1	0.00	-
7	PEG	A	618	-	6,6,6	0.57	0	5,5,5	0.49	0
8	FMT	A	622	-	0,2,2	0.00	-	0,1,1	0.00	-
8	FMT	A	625	-	0,2,2	0.00	-	0,1,1	0.00	-
8	FMT	A	623	-	0,2,2	0.00	-	0,1,1	0.00	-
8	FMT	A	621	-	0,2,2	0.00	-	0,1,1	0.00	-
7	PEG	A	616	-	6,6,6	0.65	0	5,5,5	0.64	0
8	FMT	C	615	-	0,2,2	0.00	-	0,1,1	0.00	-
8	FMT	C	616	-	0,2,2	0.00	-	0,1,1	0.00	-
8	FMT	A	626	-	0,2,2	0.00	-	0,1,1	0.00	-
8	FMT	C	618	-	0,2,2	0.00	-	0,1,1	0.00	-
7	PEG	C	614	-	6,6,6	0.53	0	5,5,5	0.42	0
6	PER	A	605	5	0,1,1	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	C	606	1	-	4/6/23/26	0/1/1/1
7	PEG	C	612	-	-	3/4/4/4	-
7	PEG	C	613	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PEG	A	617	-	-	1/4/4/4	-
7	PEG	C	614	-	-	1/4/4/4	-
7	PEG	A	619	-	-	3/4/4/4	-
7	PEG	A	618	-	-	3/4/4/4	-
7	PEG	A	616	-	-	1/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	C	606	NAG	C8-C7-N2	2.37	120.11	116.10

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	C	606	NAG	O5-C5-C6-O6
7	C	612	PEG	C4-C3-O2-C2
9	C	606	NAG	C8-C7-N2-C2
9	C	606	NAG	O7-C7-N2-C2
9	C	606	NAG	C4-C5-C6-O6
7	A	617	PEG	O1-C1-C2-O2
7	A	618	PEG	O2-C3-C4-O4
7	C	612	PEG	O1-C1-C2-O2
7	A	616	PEG	O1-C1-C2-O2
7	A	619	PEG	C4-C3-O2-C2
7	C	614	PEG	C1-C2-O2-C3
7	A	618	PEG	C1-C2-O2-C3
7	A	618	PEG	C4-C3-O2-C2
7	A	619	PEG	O1-C1-C2-O2
7	C	612	PEG	O2-C3-C4-O4
7	A	619	PEG	O2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	497/518 (95%)	-0.16	4 (0%) 86 87	34, 51, 72, 90	0
1	C	497/518 (95%)	-0.04	6 (1%) 79 80	38, 55, 75, 103	0
All	All	994/1036 (95%)	-0.10	10 (1%) 82 83	34, 53, 74, 103	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	513	LEU	2.8
1	A	518	LEU	2.6
1	C	384	GLN	2.5
1	C	338	PRO	2.5
1	C	434	SER	2.5
1	C	151	ASN	2.4
1	C	435	GLU	2.4
1	A	324	VAL	2.3
1	A	514	ASP	2.3
1	C	509	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MAN	B	3	11/12	0.72	0.29	82,95,103,104	0
3	NAG	F	2	14/15	0.82	0.30	80,92,111,119	0
2	NAG	B	2	14/15	0.83	0.27	81,100,107,111	0
2	MAN	B	4	11/12	0.83	0.22	84,96,101,103	0
3	NAG	D	2	14/15	0.87	0.16	73,90,99,101	0
2	MAN	B	5	11/12	0.90	0.38	90,95,101,102	0
2	NAG	B	1	14/15	0.90	0.17	81,88,92,93	0
4	BMA	E	3	11/12	0.91	0.15	73,79,84,91	0
4	BMA	G	3	11/12	0.92	0.17	81,92,99,103	0
3	NAG	D	1	14/15	0.92	0.16	64,71,79,88	0
4	NAG	G	2	14/15	0.96	0.11	46,52,61,66	0
4	NAG	E	2	14/15	0.96	0.21	46,63,70,72	0
3	NAG	F	1	14/15	0.96	0.14	58,63,69,74	0
4	NAG	E	1	14/15	0.97	0.14	44,50,55,58	0
4	NAG	G	1	14/15	0.97	0.14	41,48,49,49	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	FMT	C	618	3/3	0.46	0.34	67,67,71,72	0
5	CU	C	619	1/1	0.61	0.14	116,116,116,116	0
8	FMT	A	623	3/3	0.74	0.18	61,61,66,66	0
8	FMT	C	617	3/3	0.75	0.30	71,71,73,74	0
8	FMT	A	625	3/3	0.77	0.21	64,64,71,75	0
7	PEG	A	617	7/7	0.79	0.23	63,72,74,75	0
9	NAG	C	606	14/15	0.79	0.22	74,84,100,102	0
5	CU	A	627	1/1	0.79	0.10	109,109,109,109	0
7	PEG	C	612	7/7	0.82	0.27	61,66,67,69	0
8	FMT	C	615	3/3	0.84	0.20	71,71,73,76	0
8	FMT	A	626	3/3	0.85	0.18	68,68,71,73	0
5	CU	C	620	1/1	0.86	0.09	113,113,113,113	0
8	FMT	A	620	3/3	0.88	0.16	59,59,65,66	0
7	PEG	C	614	7/7	0.88	0.30	63,71,76,79	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	PEG	C	613	7/7	0.88	0.24	76,77,81,81	0
8	FMT	A	621	3/3	0.88	0.14	57,57,60,63	0
7	PEG	A	618	7/7	0.88	0.17	54,57,59,59	0
7	PEG	A	619	7/7	0.89	0.22	61,63,65,66	0
5	CU	A	628	1/1	0.89	0.05	111,111,111,111	0
7	PEG	A	616	7/7	0.90	0.17	58,62,65,66	0
8	FMT	A	624	3/3	0.91	0.16	74,74,82,82	0
8	FMT	A	622	3/3	0.94	0.12	52,52,54,58	0
8	FMT	C	616	3/3	0.94	0.15	72,72,74,76	0
5	CU	A	603	1/1	0.98	0.12	46,46,46,46	0
6	PER	C	601	2/2	0.99	0.14	62,62,62,71	0
5	CU	C	605	1/1	0.99	0.14	47,47,47,47	0
5	CU	A	601	1/1	0.99	0.16	52,52,52,52	0
5	CU	C	602	1/1	0.99	0.17	46,46,46,46	0
5	CU	A	604	1/1	0.99	0.14	49,49,49,49	0
6	PER	A	605	2/2	0.99	0.25	69,69,69,76	0
5	CU	C	604	1/1	0.99	0.14	52,52,52,52	0
5	CU	C	603	1/1	1.00	0.14	58,58,58,58	0
5	CU	A	602	1/1	1.00	0.12	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.