



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 08:52 am BST

PDB ID : 3NST
Title : Crystal Structure of Salicylate 1,2-dioxygenase G106A mutant from Pseudomonas salicylatoxidans
Authors : Ferraroni, M.; Briganti, F.; Matera, I.
Deposited on : 2010-07-02
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>14%</div> <div>• • 5%</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3136 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gentisate 1,2-dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	3	0
			2754	1753	483	504	14			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	106	ALA	GLY	ENGINEERED MUTATION	UNP Q67FT0
A	163	MET	HIS	CONFLICT	UNP Q67FT0

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	1
			8	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	343	Total	O	0	0
			343	343		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gentisate 1,2-dioxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	73.72Å 87.05Å 167.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.92 – 2.40 38.64 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.8 (83.92-2.40) 97.8 (38.64-2.40)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.152 , 0.219 0.152 , 0.217	Depositor DCC
R_{free} test set	1088 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.562	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3136	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	1/2842 (0.0%)	0.90	5/3866 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	GLU	CG-CD	5.23	1.59	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	98	TYR	CB-CG-CD2	-5.97	117.42	121.00
1	A	247	LEU	CA-CB-CG	5.71	128.43	115.30
1	A	98	TYR	CB-CG-CD1	5.21	124.12	121.00
1	A	321	LEU	CA-CB-CG	5.04	126.89	115.30
1	A	257	VAL	CB-CA-C	-5.03	101.84	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2754	0	2664	61	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	38	0	47	7	0
4	A	343	0	0	7	0
All	All	3136	0	2711	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:THR:HG22	1:A:168:GLN:H	1.18	1.05
3:A:374:GOL:H11	4:A:699:HOH:O	1.58	1.04
1:A:365:GLU:HB2	4:A:697:HOH:O	1.68	0.93
1:A:312:VAL:HG22	1:A:321:LEU:CD2	2.02	0.89
1:A:148:ARG:HH21	3:A:373:GOL:H32	1.43	0.83
1:A:312:VAL:HG22	1:A:321:LEU:HD21	1.59	0.83
1:A:261:HIS:NE2	1:A:284:HIS:HD2	1.79	0.81
1:A:271:THR:CG2	1:A:273:GLY:H	1.95	0.79
1:A:170:MET:HE3	1:A:172:TRP:CE3	2.19	0.76
1:A:217:ARG:HH22	1:A:224:ASN:HD22	1.31	0.76
1:A:236:GLU:HG2	4:A:631:HOH:O	1.87	0.74
1:A:205:SER:OG	1:A:208:GLU:HG3	1.89	0.73
1:A:245:GLN:HE22	1:A:263:ALA:H	1.34	0.73
1:A:271:THR:HG22	1:A:273:GLY:H	1.54	0.73
1:A:312:VAL:HG22	1:A:321:LEU:HD22	1.73	0.71
1:A:20:THR:HG23	1:A:22:GLU:OE2	1.96	0.65
1:A:120:ARG:HG2	1:A:159:PHE:CE2	2.32	0.65
1:A:338:GLN:HE21	3:A:374:GOL:H31	1.63	0.63
1:A:338:GLN:HE21	3:A:374:GOL:C3	2.16	0.57
1:A:291:GLU:HG2	1:A:338:GLN:HG3	1.86	0.57
1:A:271:THR:HG23	1:A:273:GLY:H	1.70	0.57
1:A:69[B]:ARG:HD3	1:A:109:TYR:OH	2.05	0.56
1:A:120:ARG:HG2	1:A:159:PHE:CD2	2.42	0.55
1:A:170:MET:CE	1:A:172:TRP:CE3	2.90	0.53
1:A:163:MET:O	1:A:163:MET:HG2	2.09	0.53
1:A:366:GLY:HA2	4:A:713:HOH:O	2.07	0.53
1:A:247:LEU:O	1:A:251:GLU:HG3	2.08	0.52
1:A:163:MET:HE2	1:A:199:TYR:CD1	2.44	0.52
1:A:22:GLU:HG2	4:A:695:HOH:O	2.08	0.52
1:A:166:THR:HG22	1:A:168:GLN:N	2.04	0.52
1:A:217:ARG:HH22	1:A:224:ASN:ND2	2.04	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ARG:HH11	1:A:113:ARG:CG	2.24	0.51
1:A:229:PRO:HD2	1:A:269:PRO:HD2	1.93	0.51
1:A:176:LEU:HD21	3:A:375[A]:GOL:H31	1.93	0.51
1:A:69[B]:ARG:NH1	1:A:109:TYR:OH	2.44	0.50
1:A:170:MET:CE	1:A:172:TRP:CZ3	2.95	0.50
1:A:140:ASN:ND2	1:A:159:PHE:H	2.10	0.50
1:A:82:GLU:H	1:A:82:GLU:CD	2.18	0.47
1:A:170:MET:HE2	1:A:172:TRP:CZ3	2.49	0.47
1:A:243:THR:HG22	1:A:247:LEU:HD22	1.96	0.46
1:A:367:GLN:HE21	1:A:367:GLN:N	2.13	0.46
1:A:146:MET:SD	1:A:170:MET:HE1	2.55	0.46
1:A:240:ARG:HG3	4:A:606:HOH:O	2.15	0.45
1:A:291:GLU:HG2	1:A:338:GLN:CG	2.47	0.45
1:A:148:ARG:NH2	3:A:373:GOL:H32	2.23	0.45
1:A:140:ASN:HD21	1:A:159:PHE:H	1.62	0.45
1:A:119:HIS:C	1:A:119:HIS:CD2	2.88	0.45
1:A:367:GLN:H	1:A:367:GLN:NE2	2.15	0.44
1:A:293:ALA:HB1	1:A:365:GLU:HG2	1.98	0.44
1:A:104:TRP:C	1:A:104:TRP:CD1	2.91	0.44
1:A:181:SER:HA	1:A:186:VAL:HG22	2.01	0.43
1:A:61:TRP:HB2	1:A:324:GLY:C	2.38	0.43
1:A:69[A]:ARG:NH1	4:A:524:HOH:O	2.43	0.42
1:A:286:LEU:N	1:A:286:LEU:HD12	2.35	0.42
1:A:295:ARG:O	1:A:334:PRO:HA	2.19	0.42
1:A:98:TYR:HH	1:A:182:GLN:CD	2.17	0.42
1:A:148:ARG:HH21	3:A:373:GOL:C3	2.22	0.41
1:A:20:THR:CG2	1:A:22:GLU:OE2	2.67	0.41
1:A:367:GLN:H	1:A:367:GLN:HE21	1.66	0.41
1:A:170:MET:HE2	1:A:172:TRP:HZ3	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	348/368 (95%)	335 (96%)	10 (3%)	3 (1%)	17 25

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	GLY
1	A	255	ALA
1	A	366	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	289/307 (94%)	271 (94%)	18 (6%)	18 29

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	THR
1	A	40	THR
1	A	60	LYS
1	A	104	TRP
1	A	113	ARG
1	A	120	ARG
1	A	163	MET
1	A	203	ASN
1	A	247	LEU
1	A	257	VAL
1	A	271	THR
1	A	312	VAL
1	A	321	LEU
1	A	345	LEU
1	A	348	PHE
1	A	361	ARG
1	A	365	GLU
1	A	367	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	140	ASN
1	A	183	GLN
1	A	224	ASN
1	A	245	GLN
1	A	284	HIS
1	A	315	ASN
1	A	338	GLN
1	A	367	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	373	-	5,5,5	0.44	0	5,5,5	0.41	0
3	GOL	A	370	-	5,5,5	0.40	0	5,5,5	0.91	0
3	GOL	A	374	-	5,5,5	0.33	0	5,5,5	1.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	375[A]	2	5,5,5	0.50	0	5,5,5	1.62	2 (40%)
3	GOL	A	371	-	5,5,5	0.62	0	5,5,5	1.57	1 (20%)
3	GOL	A	372	-	5,5,5	0.32	0	5,5,5	0.69	0
3	GOL	A	375[B]	2	5,5,5	0.38	0	5,5,5	1.32	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	373	-	-	4/4/4/4	-
3	GOL	A	370	-	-	1/4/4/4	-
3	GOL	A	374	-	-	4/4/4/4	-
3	GOL	A	375[A]	2	-	2/4/4/4	-
3	GOL	A	371	-	-	2/4/4/4	-
3	GOL	A	372	-	-	0/4/4/4	-
3	GOL	A	375[B]	2	-	2/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	375[A]	GOL	O2-C2-C1	2.38	119.60	109.12
3	A	375[B]	GOL	O2-C2-C1	2.38	119.60	109.12
3	A	371	GOL	O2-C2-C3	2.27	119.13	109.12
3	A	375[A]	GOL	C3-C2-C1	2.10	119.85	111.70

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	373	GOL	O1-C1-C2-O2
3	A	373	GOL	O1-C1-C2-C3
3	A	373	GOL	C1-C2-C3-O3
3	A	374	GOL	C1-C2-C3-O3
3	A	375[A]	GOL	O1-C1-C2-C3
3	A	370	GOL	C1-C2-C3-O3
3	A	374	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	371	GOL	C1-C2-C3-O3
3	A	375[B]	GOL	O1-C1-C2-C3
3	A	374	GOL	O1-C1-C2-O2
3	A	374	GOL	O2-C2-C3-O3
3	A	375[A]	GOL	O1-C1-C2-O2
3	A	375[B]	GOL	O1-C1-C2-O2
3	A	373	GOL	O2-C2-C3-O3
3	A	371	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	373	GOL	3	0
3	A	374	GOL	3	0
3	A	375[A]	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/368 (94%)	-0.50	6 (1%) 70 68	20, 32, 51, 73	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	366	GLY	4.8
1	A	199	TYR	3.0
1	A	367	GLN	2.9
1	A	42	LEU	2.7
1	A	173	ILE	2.2
1	A	167[A]	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	373	6/6	0.80	0.20	70,72,75,77	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	370	6/6	0.84	0.21	76,81,82,84	0
3	GOL	A	372	6/6	0.90	0.17	43,59,60,60	0
3	GOL	A	374	6/6	0.92	0.16	63,64,67,67	0
3	GOL	A	371	6/6	0.96	0.12	26,33,34,35	0
3	GOL	A	375[A]	6/6	0.97	0.26	24,37,40,41	2
3	GOL	A	375[B]	6/6	0.97	0.26	27,37,40,41	2
2	FE2	A	369	1/1	1.00	0.11	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.