



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 04:48 am BST

PDB ID : 3NT0
Title : C500S (T1D) Mutant of CueO soaked in and bound to Cu(I)
Authors : Roberts, S.A.; Montfort, W.R.; Singh, S.K.
Deposited on : 2010-07-02
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue copper oxidase cueO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3699	2347	644	681	27	0	13	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	500	SER	CYS	ENGINEERED MUTATION	UNP P36649
A	517	LEU	-	EXPRESSION TAG	UNP P36649
A	518	GLN	-	EXPRESSION TAG	UNP P36649
A	519	GLY	-	EXPRESSION TAG	UNP P36649
A	520	ASP	-	EXPRESSION TAG	UNP P36649
A	521	HIS	-	EXPRESSION TAG	UNP P36649
A	522	GLY	-	EXPRESSION TAG	UNP P36649
A	523	LEU	-	EXPRESSION TAG	UNP P36649
A	524	SER	-	EXPRESSION TAG	UNP P36649
A	525	ALA	-	EXPRESSION TAG	UNP P36649
A	526	TRP	-	EXPRESSION TAG	UNP P36649
A	527	SER	-	EXPRESSION TAG	UNP P36649
A	528	HIS	-	EXPRESSION TAG	UNP P36649
A	529	PRO	-	EXPRESSION TAG	UNP P36649
A	530	GLN	-	EXPRESSION TAG	UNP P36649
A	531	PHE	-	EXPRESSION TAG	UNP P36649
A	532	GLU	-	EXPRESSION TAG	UNP P36649
A	533	LYS	-	EXPRESSION TAG	UNP P36649

- Molecule 2 is COPPER (I) ION (three-letter code: CU1) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	9	Total	Cu	0	0
			9	9		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	386	Total	O	0	0
			386	386		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Blue copper oxidase cueO



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.86Å 92.51Å 54.33Å 90.00° 102.47° 90.00°	Depositor
Resolution (Å)	20.50 – 1.80 20.47 – 1.80	Depositor EDS
% Data completeness (in resolution range)	93.4 (20.50-1.80) 93.5 (20.47-1.80)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.207 , 0.261 0.211 , 0.263	Depositor DCC
R_{free} test set	2159 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.399	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.49 , 84.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4098	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU1, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/3796	0.74	3/5154 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	507	ASP	CB-CG-OD1	5.46	123.22	118.30
1	A	280	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	A	47	ARG	NE-CZ-NH2	-5.26	117.67	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3699	0	3675	34	0
2	A	9	0	0	0	0
3	A	4	0	3	0	0
4	A	386	0	0	6	0
All	All	4098	0	3678	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:TRP:HB3	1:A:105[B]:LEU:HD13	1.63	0.81
1:A:337:LEU:HD13	4:A:840:HOH:O	1.84	0.77
1:A:358:MET:HA	1:A:358:MET:HE3	1.73	0.68
1:A:358:MET:HA	1:A:358:MET:CE	2.24	0.66
1:A:307[A]:ILE:HD13	1:A:307[A]:ILE:N	2.11	0.63
1:A:202:VAL:CG1	1:A:366[B]:MET:SD	2.88	0.62
1:A:520:ASP:C	1:A:521:HIS:HD1	2.03	0.61
1:A:319:ILE:N	1:A:319:ILE:HD12	2.17	0.60
1:A:307[A]:ILE:HD11	1:A:375:ALA:HA	1.84	0.60
1:A:520:ASP:O	1:A:521:HIS:ND1	2.32	0.58
1:A:440:MET:HE3	1:A:440:MET:HA	1.90	0.54
1:A:48:ILE:HD12	1:A:86[B]:VAL:HG11	1.89	0.53
1:A:358:MET:HE1	1:A:361:MET:HE2	1.89	0.53
1:A:358:MET:HE1	1:A:361:MET:CE	2.39	0.52
1:A:202:VAL:HG13	1:A:366[B]:MET:SD	2.50	0.52
1:A:487:ASN:ND2	4:A:834:HOH:O	2.40	0.52
1:A:364[B]:MET:O	1:A:368:MET:HG2	2.12	0.50
1:A:359:LEU:HD22	1:A:359:LEU:O	2.13	0.48
1:A:199:GLN:HG2	1:A:199:GLN:O	2.15	0.47
1:A:201:ASP:OD1	1:A:204:THR:HG23	2.16	0.46
1:A:252:ARG:HH12	1:A:288:ASN:HD22	1.65	0.45
1:A:189:LYS:NZ	1:A:210:PHE:O	2.46	0.44
1:A:319:ILE:CD1	1:A:319:ILE:N	2.80	0.44
1:A:341:PRO:CG	1:A:483:LEU:HD21	2.47	0.43
1:A:203:MET:O	1:A:207:VAL:HG22	2.19	0.42
1:A:374:GLN:HG2	4:A:669:HOH:O	2.19	0.41
1:A:141:HIS:HB2	1:A:142:PRO:HD2	2.00	0.41
1:A:418:ASN:ND2	4:A:804:HOH:O	2.53	0.41
1:A:440:MET:HE3	4:A:792:HOH:O	2.21	0.41
1:A:118:ILE:HD11	4:A:549:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	478/505 (95%)	459 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/414 (96%)	379 (95%)	20 (5%)	24	10

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	HIS
1	A	181	VAL
1	A	199	GLN
1	A	228	ARG
1	A	266	GLU
1	A	280	ARG
1	A	307[A]	ILE
1	A	307[B]	ILE
1	A	327	SER
1	A	333[A]	THR
1	A	333[B]	THR
1	A	337	LEU
1	A	340	LEU
1	A	343	LEU
1	A	358	MET
1	A	359	LEU
1	A	361	MET
1	A	440	MET
1	A	470	LYS
1	A	517	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	GLN
1	A	186	GLN
1	A	288	ASN
1	A	374	GLN
1	A	408	ASN
1	A	418	ASN
1	A	428	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 9 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	610	2	1,3,3	1.85	0	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	468/505 (92%)	0.55	36 (7%) 13 10	9, 20, 33, 51	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	ALA	5.3
1	A	520	ASP	5.2
1	A	403	ASP	4.9
1	A	378	GLY	4.9
1	A	377	ALA	4.3
1	A	327	SER	4.2
1	A	326	ALA	4.2
1	A	341	PRO	4.0
1	A	517	LEU	4.0
1	A	406	HIS	3.7
1	A	342	SER	3.5
1	A	521	HIS	3.4
1	A	361	MET	3.4
1	A	30	GLU	3.3
1	A	171[A]	MET	3.2
1	A	519	GLY	3.1
1	A	344	GLU	3.1
1	A	343	LEU	3.1
1	A	376	MET	3.0
1	A	345	GLY	2.9
1	A	518	GLN	2.9
1	A	340	LEU	2.7
1	A	358	MET	2.7
1	A	207	VAL	2.6
1	A	165	ASP	2.6
1	A	105[A]	LEU	2.6
1	A	364[A]	MET	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	337	LEU	2.5
1	A	338	PRO	2.4
1	A	60	GLY	2.4
1	A	405	HIS	2.4
1	A	440	MET	2.3
1	A	307[A]	ILE	2.3
1	A	429	TYR	2.2
1	A	231	LEU	2.1
1	A	192	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU1	A	609	1/1	0.85	0.32	35,35,35,35	1
3	ACT	A	610	4/4	0.92	0.13	19,20,21,21	0
2	CU1	A	606	1/1	0.96	0.12	45,45,45,45	0
2	CU1	A	607	1/1	0.97	0.10	38,38,38,38	0
2	CU1	A	605	1/1	0.98	0.13	44,44,44,44	0
2	CU1	A	602	1/1	0.99	0.07	18,18,18,18	0
2	CU1	A	608	1/1	0.99	0.08	33,33,33,33	0
2	CU1	A	603	1/1	0.99	0.06	19,19,19,19	0
2	CU1	A	601	1/1	0.99	0.05	17,17,17,17	0
2	CU1	A	604	1/1	1.00	0.04	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.