



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:36 am BST

PDB ID : 4O0G
Title : Crystal structure of the human L-asparaginase protein T219V mutant
Authors : Nomme, J.; Lavie, A.
Deposited on : 2013-12-13
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	 71% 21% • 5%
1	B	309	 65% 25% • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoaspartyl peptidase/L-asparaginase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	3	0
			2120	1324	374	407	15			
1	B	293	Total	C	N	O	S	0	2	0
			2122	1320	376	412	14			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	EXPRESSION TAG	UNP Q7L266
A	219	VAL	THR	ENGINEERED MUTATION	UNP Q7L266
B	0	HIS	-	EXPRESSION TAG	UNP Q7L266
B	219	VAL	THR	ENGINEERED MUTATION	UNP Q7L266

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	69	Total	O	0	0
			69	69		
3	B	83	Total	O	0	0
			83	83		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isoaspartyl peptidase/L-asparaginase

Chain A: 



- Molecule 1: Isoaspartyl peptidase/L-asparaginase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	59.44Å 59.44Å 306.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.18 – 2.10 29.18 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.1 (29.18-2.10) 97.1 (29.18-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.165 , 0.236 0.170 , 0.215	Depositor DCC
R_{free} test set	1734 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.509	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 29.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.477 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.514 for H, K, L 0.486 for K, H, -L	Depositor
Outliers	0 of 35125 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4396	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2160	0.89	1/2918 (0.0%)
1	B	0.70	0/2158	0.87	1/2915 (0.0%)
All	All	0.69	0/4318	0.88	2/5833 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	ASP	CB-CG-OD2	5.81	123.53	118.30
1	B	293	LEU	CA-CB-CG	5.38	127.66	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2120	0	2116	46	0
1	B	2122	0	2118	51	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	69	0	0	4	0
3	B	83	0	0	5	0
All	All	4396	0	4234	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:HIS:O	3:A:530:HOH:O	2.09	0.71
1:B:69:LEU:HD21	1:B:140:VAL:HG13	1.73	0.71
1:A:143:ARG:O	1:A:147:ARG:HG3	1.92	0.69
1:A:221:HIS:HD1	1:A:224:SER:HG	1.35	0.69
1:A:202:CYS:HB3	3:B:542:HOH:O	1.92	0.68
1:A:245:GLU:O	1:A:249:ASP:OD1	2.13	0.67
1:B:241:GLY:O	3:B:539:HOH:O	2.14	0.65
1:B:2:ASN:O	3:B:545:HOH:O	2.15	0.64
1:A:248:ALA:O	1:A:252:LEU:HB2	1.99	0.61
1:B:49:GLU:OE1	1:B:106:ARG:NH1	2.33	0.61
1:B:5:VAL:HG12	1:B:288:ALA:HB3	1.81	0.61
1:B:293:LEU:HD22	1:B:307:LEU:CD2	2.30	0.61
1:B:295:PHE:O	1:B:302:THR:HA	2.00	0.60
1:A:117:LEU:HD23	1:B:196:ARG:HA	1.83	0.59
1:A:80:SER:OG	1:A:200:SER:O	2.18	0.58
1:B:49:GLU:O	1:B:53:VAL:HG23	2.04	0.56
1:A:248:ALA:HB2	1:A:265:LEU:HD11	1.87	0.56
1:A:216:VAL:HG22	1:A:267:VAL:HG23	1.86	0.56
1:A:180:VAL:HG22	1:A:211:ASN:HA	1.88	0.55
1:A:24:HIS:O	1:A:28:VAL:HG23	2.06	0.55
1:A:97:CYS:HB3	1:A:134:ILE:O	2.08	0.54
1:B:245:GLU:HG2	1:B:275:VAL:HG21	1.89	0.54
1:B:232:ARG:NH1	1:B:236[A]:PHE:CD1	2.76	0.54
1:A:39:ARG:C	1:A:41:GLY:H	2.11	0.53
1:B:205:ALA:CB	1:B:226:LEU:HG	2.39	0.52
1:B:256:LYS:O	1:B:260:LYS:HA	2.10	0.52
1:B:43:SER:N	1:B:46:ASP:OD2	2.38	0.52
1:A:97:CYS:HA	1:A:139:LEU:HD11	1.92	0.52
1:B:150:LYS:HA	1:B:153:HIS:HD2	1.77	0.50
1:A:221:HIS:CE1	1:A:224:SER:HG	2.27	0.50
1:A:224:SER:HA	1:A:227:LYS:HE2	1.94	0.50
1:B:19:ARG:O	1:B:20:LYS:C	2.48	0.50
1:B:233:LEU:HA	1:B:236[A]:PHE:CD2	2.46	0.50
1:B:276:ALA:O	1:B:277:LYS:HD2	2.11	0.50
1:B:56:GLU:OE1	1:B:102:ILE:HG22	2.10	0.50
1:A:216:VAL:CG2	1:A:267:VAL:HG23	2.41	0.50
1:A:175:ASP:OD2	1:A:179:ASN:HB2	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:THR:HB	1:B:219:VAL:HG12	1.94	0.49
1:A:218:THR:O	3:A:551:HOH:O	2.19	0.49
1:A:200:SER:N	1:A:201:PRO:CD	2.76	0.49
1:B:293:LEU:HD22	1:B:307:LEU:HD21	1.93	0.49
1:A:70:ASN:C	1:A:70:ASN:OD1	2.52	0.48
1:A:242:LYS:HD2	1:A:246:GLU:OE1	2.14	0.48
1:B:232:ARG:NH1	1:B:236[A]:PHE:CE1	2.82	0.47
1:B:264:GLY:HA2	1:B:278:TRP:HA	1.96	0.47
1:A:92:VAL:HB	1:A:95:VAL:CG1	2.45	0.47
1:B:216:VAL:HG22	1:B:267:VAL:HB	1.94	0.47
1:B:293:LEU:HD22	1:B:307:LEU:HD22	1.96	0.47
1:B:205:ALA:HB1	1:B:226:LEU:HG	1.97	0.47
1:B:199:ASP:HA	1:B:226:LEU:HD12	1.95	0.47
1:B:252:LEU:HD21	1:B:265:LEU:HG	1.98	0.46
1:A:16:SER:O	1:A:20:LYS:HB2	2.16	0.46
1:A:135:PRO:HB2	1:A:138:LYS:HD3	1.97	0.46
1:A:216:VAL:HG22	1:A:267:VAL:CG2	2.46	0.46
1:A:248:ALA:CB	1:A:265:LEU:HD11	2.44	0.46
1:A:174:LEU:HD12	1:A:179:ASN:O	2.15	0.46
1:B:244:VAL:HG22	3:B:509:HOH:O	2.16	0.45
1:A:118:THR:O	1:A:119:ASP:C	2.55	0.45
1:A:121:GLY:HA3	1:B:194:VAL:O	2.16	0.45
1:B:275:VAL:HG22	1:B:276:ALA:N	2.31	0.45
1:B:268:VAL:HA	1:B:273:ASP:O	2.17	0.45
1:B:75:VAL:HG12	1:B:77:MET:HG3	1.98	0.44
1:B:58:ASP:C	1:B:58:ASP:OD1	2.56	0.44
1:A:208:TYR:O	1:A:215:ALA:HA	2.17	0.44
1:A:267:VAL:CG1	1:A:275:VAL:HG12	2.47	0.44
1:A:114:HIS:CG	1:B:196:ARG:HG3	2.52	0.44
1:B:78:ASP:HB3	1:B:201:PRO:HG3	2.00	0.44
1:B:3:PRO:HB3	1:B:38:LEU:HD13	2.00	0.44
1:A:6:VAL:O	1:A:171:ALA:HA	2.17	0.43
1:B:45:VAL:HG13	1:B:81:ILE:HD11	2.01	0.43
1:B:8:HIS:CD2	1:B:266:ILE:CD1	3.01	0.43
1:A:223:GLU:O	1:A:226:LEU:HB2	2.19	0.42
1:B:92:VAL:HA	1:B:117:LEU:O	2.19	0.42
1:A:142:GLU:HA	1:A:145:LYS:HD2	2.00	0.42
1:A:36:GLY:O	1:A:39:ARG:HB2	2.19	0.42
1:A:136:GLY:O	1:A:140:VAL:HG23	2.19	0.42
1:B:8:HIS:CD2	1:B:266:ILE:HD13	2.54	0.42
1:A:213:ILE:HD11	3:A:511:HOH:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:SER:O	1:B:228:VAL:HG22	2.20	0.42
1:B:15:ILE:HG13	1:B:20:LYS:HG2	2.02	0.41
1:A:259:VAL:HB	3:A:526:HOH:O	2.19	0.41
1:B:244:VAL:CG1	1:B:269:SER:HB3	2.50	0.41
1:A:230:LEU:O	1:A:231:ALA:C	2.57	0.41
1:B:221:HIS:CE1	1:B:223:GLU:HB2	2.55	0.41
1:A:230:LEU:O	1:A:233:LEU:HB3	2.21	0.41
1:B:102:ILE:HA	1:B:102:ILE:HD12	1.96	0.41
1:B:107:LEU:O	1:B:111:LYS:HB2	2.21	0.41
1:A:182:TYR:CD1	1:A:182:TYR:C	2.93	0.41
1:A:200:SER:N	1:A:201:PRO:HD2	2.36	0.41
1:B:21:GLU:OE2	1:B:25:GLN:HG3	2.21	0.41
1:A:92:VAL:HB	1:A:95:VAL:HG11	2.03	0.41
1:B:264:GLY:HA3	1:B:283:MET:SD	2.61	0.41
1:B:182:TYR:CD1	1:B:182:TYR:C	2.94	0.40
1:B:245:GLU:HB2	3:B:572:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/309 (94%)	272 (94%)	16 (6%)	2 (1%)	22	18
1	B	289/309 (94%)	271 (94%)	15 (5%)	3 (1%)	15	11
All	All	579/618 (94%)	543 (94%)	31 (5%)	5 (1%)	17	12

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	SER
1	A	280	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	63	ALA
1	B	146	LYS
1	B	147	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/235 (92%)	201 (93%)	16 (7%)	13	10
1	B	218/235 (93%)	186 (85%)	32 (15%)	3	1
All	All	435/470 (93%)	387 (89%)	48 (11%)	6	3

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	18	ASP
1	A	40	GLU
1	A	71	THR
1	A	133	GLU
1	A	134	ILE
1	A	142	GLU
1	A	150	LYS
1	A	182	TYR
1	A	192	LYS
1	A	226	LEU
1	A	229	ASN
1	A	245	GLU
1	A	269	SER
1	A	277	LYS
1	A	305	THR
1	B	1	MET
1	B	4	ILE
1	B	8	HIS
1	B	17	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	21	GLU
1	B	72	ASN
1	B	111	LYS
1	B	116	PHE
1	B	137	GLU
1	B	149	GLU
1	B	150	LYS
1	B	182	TYR
1	B	184	THR
1	B	189	ILE
1	B	217	SER
1	B	226	LEU
1	B	229	ASN
1	B	240	GLN
1	B	244	VAL
1	B	245	GLU
1	B	260	LYS
1	B	267	VAL
1	B	269	SER
1	B	270	LYS
1	B	273	ASP
1	B	279	THR
1	B	282[A]	SER
1	B	282[B]	SER
1	B	290	ASP
1	B	292	LYS
1	B	293	LEU
1	B	307	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	124	GLN
1	B	153	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/309 (94%)	-0.69	0 100 100	19, 26, 42, 56	0
1	B	293/309 (94%)	-0.68	0 100 100	18, 27, 45, 53	0
All	All	586/618 (94%)	-0.69	0 100 100	18, 27, 43, 56	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NA	A	401	1/1	0.99	0.05	31,31,31,31	0
2	NA	B	401	1/1	1.00	0.05	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.