



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2021 – 11:24 PM EDT

PDB ID : 1O1N
Title : Deoxy hemoglobin (A:GLYGLYGLY-C:V1M,L29W; B,D:V1M)
Authors : Brucker, E.A.
Deposited on : 2002-11-17
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	 86% 13% .
2	B	146	 2% 86% 12% .
2	D	146	 2% 87% 10% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4818 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemoglobin Alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	0	0
			2145	1376	376	387	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	VAL	engineered mutation	UNP P69905
A	29	TRP	LEU	engineered mutation	UNP P69905
A	142	GLY	-	linker	UNP P69905
A	143	GLY	-	linker	UNP P69905
A	144	GLY	-	linker	UNP P69905
A	173	TRP	LEU	engineered mutation	UNP P69905


- Molecule 2 is a protein called Hemoglobin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1124	724	195	201	4			
2	D	146	Total	C	N	O	S	0	0	0
			1124	724	195	201	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	VAL	engineered mutation	UNP P68871
D	1	MET	VAL	engineered mutation	UNP P68871

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	136	Total	O	0	0
			136	136		
4	B	54	Total	O	0	0
			54	54		
4	D	63	Total	O	0	0
			63	63		

- Molecule 1: Hemoglobin Alpha chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.23Å 83.13Å 53.80Å 90.00° 98.96° 90.00°	Depositor
Resolution (Å)	8.00 – 1.80 25.95 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.4 (8.00-1.80) 92.0 (25.95-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.80Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.191 , 0.242 0.173 , 0.182	Depositor DCC
R_{free} test set	2346 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4818	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/2205	0.65	0/3000
2	B	0.29	0/1154	0.68	0/1566
2	D	0.27	0/1154	0.66	0/1566
All	All	0.29	0/4513	0.66	0/6132

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2145	0	2127	22	0
2	B	1124	0	1118	13	0
2	D	1124	0	1118	14	0
3	A	86	0	60	0	0
3	B	43	0	30	0	0
3	D	43	0	30	1	0
4	A	136	0	0	4	0
4	B	54	0	0	1	0
4	D	63	0	0	2	0
All	All	4818	0	4483	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:GLU:HG3	4:A:1212:HOH:O	1.85	0.74
1:A:51:GLY:O	1:A:56:LYS:HE2	1.89	0.73
1:A:238:ASP:OD1	1:A:240:VAL:HG13	1.92	0.70
2:B:100:PRO:HB2	2:B:104:ARG:HH21	1.59	0.68
2:D:139:ASN:HB3	4:D:1147:HOH:O	1.95	0.67
1:A:94:ASP:OD1	1:A:96:VAL:HG13	1.94	0.66
2:D:18:VAL:HG13	2:D:23:VAL:HG21	1.82	0.62
2:D:1:MET:HB3	2:D:78:LEU:O	2.00	0.61
1:A:7:LYS:O	1:A:11:LYS:HG3	2.00	0.61
2:D:101:GLU:HA	2:D:104:ARG:HH11	1.68	0.59
1:A:221:PRO:HD2	4:A:1154:HOH:O	2.03	0.59
2:D:66:LYS:HD2	3:D:147:HEM:HAA1	1.84	0.59
1:A:76:MET:HE1	1:A:132:VAL:HA	1.84	0.59
2:D:20:VAL:CG1	2:D:68:LEU:HB3	2.33	0.58
1:A:112:HIS:O	1:A:113:LEU:HD23	2.03	0.58
2:B:21:ASP:OD1	2:B:65:LYS:HG3	2.03	0.58
1:A:160:LYS:HE3	1:A:257:LEU:HD22	1.86	0.57
2:B:1:MET:HB3	2:B:78:LEU:O	2.05	0.57
1:A:256:HIS:O	1:A:257:LEU:HD23	2.05	0.56
1:A:257:LEU:HB3	1:A:260:GLU:HB2	1.88	0.55
1:A:3:SER:O	1:A:7:LYS:HG3	2.05	0.55
1:A:84:SER:HB3	1:A:139:LYS:HD2	1.88	0.55
1:A:146:LEU:O	1:A:151:LYS:HE3	2.08	0.54
2:D:20:VAL:HG13	2:D:68:LEU:HB3	1.89	0.54
1:A:204:LYS:HE3	1:A:208:ASP:OD2	2.10	0.52
2:B:24:GLY:HA2	2:B:68:LEU:HD22	1.90	0.51
2:B:23:VAL:HG12	2:B:68:LEU:HD21	1.93	0.50
2:D:3:LEU:HD12	2:D:8:LYS:HG3	1.93	0.50
2:B:23:VAL:HG12	2:B:68:LEU:CD2	2.43	0.49
2:D:101:GLU:HA	2:D:104:ARG:NH1	2.27	0.49
2:B:144:LYS:HE2	2:B:144:LYS:HB3	1.68	0.46
2:D:95:LYS:HA	2:D:95:LYS:HD2	1.79	0.45
2:D:21:ASP:OD1	2:D:65:LYS:HE3	2.16	0.45
1:A:236:ARG:NH1	4:A:1078:HOH:O	2.50	0.45
2:D:101:GLU:OE1	2:D:104:ARG:NH1	2.50	0.44
2:B:95:LYS:HD3	2:B:95:LYS:HA	1.69	0.44
2:B:39:GLN:HB2	4:B:1115:HOH:O	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ALA:HB1	1:A:63:ALA:HB1	1.99	0.43
1:A:66:LEU:O	1:A:70:VAL:HG23	2.17	0.43
2:B:24:GLY:CA	2:B:68:LEU:HD22	2.48	0.43
1:A:196:SER:O	1:A:200:LYS:HG3	2.18	0.42
2:B:38:THR:HG22	2:B:102:ASN:OD1	2.20	0.42
2:B:51:PRO:O	2:B:55:MET:HG2	2.19	0.42
1:A:258:PRO:HD2	4:A:1222:HOH:O	2.19	0.41
1:A:256:HIS:C	1:A:257:LEU:HD23	2.41	0.41
2:D:144:LYS:HD3	4:D:1194:HOH:O	2.21	0.41
2:D:124:PRO:HB2	2:D:125:PRO:HD3	2.02	0.41
2:B:124:PRO:HB2	2:B:125:PRO:HD3	2.01	0.40
1:A:112:HIS:C	1:A:113:LEU:HD23	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/285 (98%)	273 (98%)	5 (2%)	0	100	100
2	B	144/146 (99%)	143 (99%)	1 (1%)	0	100	100
2	D	144/146 (99%)	143 (99%)	1 (1%)	0	100	100
All	All	566/577 (98%)	559 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/226 (99%)	219 (98%)	5 (2%)	52	39
2	B	118/118 (100%)	113 (96%)	5 (4%)	30	15
2	D	118/118 (100%)	112 (95%)	6 (5%)	24	10
All	All	460/462 (100%)	444 (96%)	16 (4%)	36	21

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	SER
1	A	81	SER
1	A	84	SER
1	A	167	GLU
1	A	228	SER
2	B	1	MET
2	B	8	LYS
2	B	66	LYS
2	B	104	ARG
2	B	144	LYS
2	D	1	MET
2	D	8	LYS
2	D	59	LYS
2	D	66	LYS
2	D	68	LEU
2	D	95	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
1	A	216	HIS
2	B	63	HIS
2	B	117	HIS
2	D	63	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	HEM	A	287	1	27,50,50	1.40	6 (22%)	17,82,82	1.80	6 (35%)
3	HEM	D	147	2	27,50,50	1.38	6 (22%)	17,82,82	1.92	7 (41%)
3	HEM	B	147	2	27,50,50	1.38	6 (22%)	17,82,82	1.88	6 (35%)
3	HEM	A	286	1	27,50,50	1.39	6 (22%)	17,82,82	1.93	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	287	1	-	0/6/54/54	-
3	HEM	D	147	2	-	2/6/54/54	-
3	HEM	B	147	2	-	0/6/54/54	-
3	HEM	A	286	1	-	0/6/54/54	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	286	HEM	C3B-CAB	-2.65	1.42	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	147	HEM	C3C-CAC	-2.64	1.42	1.47
3	D	147	HEM	C3B-CAB	-2.64	1.42	1.47
3	D	147	HEM	C3C-C2C	-2.62	1.36	1.40
3	A	287	HEM	C3C-C2C	-2.61	1.36	1.40
3	A	287	HEM	C3B-CAB	-2.61	1.42	1.47
3	A	286	HEM	C3C-CAC	-2.59	1.42	1.47
3	B	147	HEM	CBB-CAB	2.54	1.46	1.29
3	A	286	HEM	C3B-C2B	-2.54	1.36	1.40
3	A	287	HEM	C3B-C2B	-2.51	1.36	1.40
3	A	287	HEM	CBC-CAC	2.47	1.45	1.29
3	D	147	HEM	C3B-C2B	-2.46	1.36	1.40
3	B	147	HEM	C3B-CAB	-2.46	1.42	1.47
3	A	286	HEM	CBC-CAC	2.44	1.45	1.29
3	A	287	HEM	CBB-CAB	2.44	1.45	1.29
3	B	147	HEM	C3C-C2C	-2.43	1.37	1.40
3	D	147	HEM	CBC-CAC	2.42	1.45	1.29
3	A	287	HEM	C3C-CAC	-2.42	1.42	1.47
3	A	286	HEM	CBB-CAB	2.42	1.45	1.29
3	A	286	HEM	C3C-C2C	-2.41	1.37	1.40
3	B	147	HEM	C3B-C2B	-2.41	1.37	1.40
3	D	147	HEM	C3C-CAC	-2.39	1.42	1.47
3	B	147	HEM	CBC-CAC	2.37	1.44	1.29
3	D	147	HEM	CBB-CAB	2.36	1.44	1.29

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	147	HEM	CBA-CAA-C2A	4.42	120.64	112.49
3	A	286	HEM	CBA-CAA-C2A	4.17	120.17	112.49
3	B	147	HEM	CBA-CAA-C2A	3.44	118.83	112.49
3	A	287	HEM	CBD-CAD-C3D	2.91	117.84	112.48
3	B	147	HEM	CBD-CAD-C3D	2.84	117.71	112.48
3	A	286	HEM	CBD-CAD-C3D	2.83	117.69	112.48
3	A	287	HEM	CBA-CAA-C2A	2.80	117.65	112.49
3	A	287	HEM	CMA-C3A-C4A	-2.43	124.73	128.46
3	A	286	HEM	CMA-C3A-C4A	-2.39	124.79	128.46
3	B	147	HEM	CMA-C3A-C4A	-2.39	124.80	128.46
3	D	147	HEM	CMA-C3A-C4A	-2.38	124.81	128.46
3	D	147	HEM	CMD-C2D-C1D	-2.21	125.07	128.46
3	B	147	HEM	CAD-CBD-CGD	2.21	116.37	112.67
3	B	147	HEM	CMD-C2D-C1D	-2.19	125.10	128.46
3	D	147	HEM	CMC-C2C-C3C	2.18	128.75	124.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	287	HEM	CMA-C3A-C2A	2.13	128.96	124.94
3	B	147	HEM	CMA-C3A-C2A	2.13	128.95	124.94
3	A	287	HEM	CMB-C2B-C3B	2.12	128.65	124.68
3	A	286	HEM	CMA-C3A-C2A	2.09	128.88	124.94
3	A	286	HEM	CMB-C2B-C3B	2.09	128.58	124.68
3	D	147	HEM	CMB-C2B-C3B	2.04	128.50	124.68
3	D	147	HEM	CAA-CBA-CGA	2.03	116.07	112.67
3	D	147	HEM	CMA-C3A-C2A	2.02	128.76	124.94
3	A	287	HEM	CMD-C2D-C1D	-2.00	125.39	128.46

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	147	HEM	C1A-C2A-CAA-CBA
3	D	147	HEM	C3A-C2A-CAA-CBA

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	147	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand HEM D 147


Bond lengths

Bond angles

Torsions

Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/285 (98%)	-0.18	1 (0%) 92 90	14, 24, 43, 56	0
2	B	146/146 (100%)	-0.06	2 (1%) 75 72	15, 25, 44, 73	0
2	D	146/146 (100%)	0.07	3 (2%) 63 59	15, 25, 43, 73	0
All	All	574/577 (99%)	-0.09	6 (1%) 82 80	14, 25, 44, 73	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	MET	6.0
2	B	2	HIS	3.6
2	D	47	ASP	3.2
2	D	2	HIS	2.8
2	D	1	MET	2.7
1	A	160	LYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	HEM	B	147	43/43	0.96	0.10	16,20,41,53	0
3	HEM	D	147	43/43	0.96	0.13	14,20,41,53	0
3	HEM	A	286	43/43	0.97	0.09	14,19,35,41	0
3	HEM	A	287	43/43	0.98	0.09	14,19,35,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM D 147:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 286:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.