



wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 01:19 am BST

PDB ID : 1O2D
Title : Crystal structure of Alcohol dehydrogenase, iron-containing (TM0920) from *Thermotoga maritima* at 1.30 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2003-02-27
Resolution : 1.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	371	 91% 5% . .
1	B	371	 92% 5% . .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6604 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase, iron-containing.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	359	Total	C	N	O	S	Se	0	12	0
			2805	1797	461	533	2	12			
1	B	360	Total	C	N	O	S	Se	0	10	0
			2791	1789	460	527	2	13			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	LEADER SEQUENCE	UNP Q9X022
A	-10	GLY	-	LEADER SEQUENCE	UNP Q9X022
A	-9	SER	-	LEADER SEQUENCE	UNP Q9X022
A	-8	ASP	-	LEADER SEQUENCE	UNP Q9X022
A	-7	LYS	-	LEADER SEQUENCE	UNP Q9X022
A	-6	ILE	-	LEADER SEQUENCE	UNP Q9X022
A	-5	HIS	-	LEADER SEQUENCE	UNP Q9X022
A	-4	HIS	-	LEADER SEQUENCE	UNP Q9X022
A	-3	HIS	-	LEADER SEQUENCE	UNP Q9X022
A	-2	HIS	-	LEADER SEQUENCE	UNP Q9X022
A	-1	HIS	-	LEADER SEQUENCE	UNP Q9X022
A	0	HIS	-	LEADER SEQUENCE	UNP Q9X022
A	1	VAL	-	LEADER SEQUENCE	UNP Q9X022
A	6	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	76	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	99	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	163	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	178	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	215	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	236	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	245	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	276	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	282	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
A	285	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	-11	MET	-	LEADER SEQUENCE	UNP Q9X022

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	GLY	-	LEADER SEQUENCE	UNP Q9X022
B	-9	SER	-	LEADER SEQUENCE	UNP Q9X022
B	-8	ASP	-	LEADER SEQUENCE	UNP Q9X022
B	-7	LYS	-	LEADER SEQUENCE	UNP Q9X022
B	-6	ILE	-	LEADER SEQUENCE	UNP Q9X022
B	-5	HIS	-	LEADER SEQUENCE	UNP Q9X022
B	-4	HIS	-	LEADER SEQUENCE	UNP Q9X022
B	-3	HIS	-	LEADER SEQUENCE	UNP Q9X022
B	-2	HIS	-	LEADER SEQUENCE	UNP Q9X022
B	-1	HIS	-	LEADER SEQUENCE	UNP Q9X022
B	0	HIS	-	LEADER SEQUENCE	UNP Q9X022
B	1	VAL	-	LEADER SEQUENCE	UNP Q9X022
B	6	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	76	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	99	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	163	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	178	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	215	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	236	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	245	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	276	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	282	MSE	MET	MODIFIED RESIDUE	UNP Q9X022
B	285	MSE	MET	MODIFIED RESIDUE	UNP Q9X022

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



-
- Chemical structure of TRS (2,3,4-trihydroxy-5-aminopentanoate) is shown. The structure is a five-carbon chain with a carboxylate group at C1 and an amino group at C5. The carbons are labeled C1, C2, C3, C4, and C5. The hydroxyl groups are labeled O1, O2, and O3. The amino group is labeled NH₃⁺N.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	1
			10	5	1	4		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	479	Total	O	0	0
			479	479		
5	B	413	Total	O	0	0
			413	413		

- Molecule 1: Alcohol dehydrogenase, iron-containing

MET	GLY	SER	ASP	LYS	ILE	HIS	HIS	HIS	HIS	HIS	HIS	V1	Y5	E18	D25	K51	D54	E55	F72	D73	M76	D100	D113	R158	H256	Y260	T264	H270	M276	T343	K348	N351	V353
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------

Chain B: 92% 5% •

MET	GLY	SER	ASP	LYS	ILE	HIS	HIS	HIS	HIS	HIS	H-1	E18	D25	K41	F72	M76	E196	R230	E231	K234	H256	T264	H270	G271	K272	M276	V281	M282	M285	K286	E287	L304	G358
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.05Å 85.28Å 72.08Å 90.00° 96.29° 90.00°	Depositor
Resolution (Å)	38.19 – 1.30 38.19 – 1.30	Depositor EDS
% Data completeness (in resolution range)	97.8 (38.19-1.30) 97.8 (38.19-1.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.1.955	Depositor
R, R_{free}	0.137 , 0.170 0.204 , 0.225	Depositor DCC
R_{free} test set	8371 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6604	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, NAP, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	1/2903 (0.0%)	0.76	3/3906 (0.1%)
1	B	0.58	0/2886	0.74	1/3887 (0.0%)
All	All	0.61	1/5789 (0.0%)	0.75	4/7793 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	276	MSE	SE-CE	-7.78	1.49	1.95

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	25	ASP	CB-CG-OD2	6.19	123.87	118.30
1	A	25	ASP	CB-CG-OD1	5.88	123.59	118.30
1	A	113	ASP	CB-CG-OD2	5.86	123.57	118.30
1	A	158	ARG	NE-CZ-NH2	-5.68	117.46	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2805	0	2819	14	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2791	0	2763	14	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	48	0	24	2	0
3	B	48	0	25	2	0
4	A	8	0	12	0	0
4	B	10	0	6	0	0
5	A	479	0	0	7	1
5	B	413	0	0	6	1
All	All	6604	0	5649	29	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:THR:HG22	5:A:3015:HOH:O	1.64	0.96
1:B:287:GLU:O	5:B:2833:HOH:O	1.92	0.86
1:A:351[B]:ASN:ND2	5:A:2920:HOH:O	2.15	0.77
1:A:351[B]:ASN:OD1	5:A:3041:HOH:O	2.11	0.68
1:A:256[B]:HIS:NE2	1:A:260:TYR:OH	2.27	0.66

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:-1:HIS:N	5:A:2941:HOH:O[2_647]	2.15	0.05
1:A:5:TYR:OH	5:B:3213:HOH:O[2_657]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/371 (100%)	361 (98%)	8 (2%)	0	100	100
1	B	368/371 (99%)	361 (98%)	7 (2%)	0	100	100
All	All	737/742 (99%)	722 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/308 (101%)	308 (99%)	3 (1%)	76	48
1	B	302/308 (98%)	299 (99%)	3 (1%)	76	48
All	All	613/616 (100%)	607 (99%)	6 (1%)	84	48

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256[B]	HIS
1	B	256[B]	HIS
1	B	25	ASP
1	A	256[A]	HIS
1	B	256[A]	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	-1	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	TRS	A	2801	-	7,7,7	0.26	0	9,9,9	0.80	0
4	TRS	B	2800[A]	-	7,7,7	0.21	0	9,9,9	0.35	0
4	TRS	B	2800[B]	-	7,7,7	0.29	0	9,9,9	0.59	0
3	NAP	A	1800	-	45,52,52	1.52	3 (6%)	56,80,80	1.36	7 (12%)
3	NAP	B	1801	-	45,52,52	1.68	4 (8%)	56,80,80	1.16	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	A	2801	-	-	0/9/9/9	-
4	TRS	B	2800[A]	-	-	0/9/9/9	-
4	TRS	B	2800[B]	-	-	1/9/9/9	-
3	NAP	A	1800	-	-	5/31/67/67	0/5/5/5
3	NAP	B	1801	-	-	5/31/67/67	0/5/5/5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1801	NAP	O7N-C7N	8.43	1.40	1.24
3	A	1800	NAP	O7N-C7N	8.14	1.39	1.24
3	B	1801	NAP	C2A-N3A	3.79	1.38	1.32
3	B	1801	NAP	C2N-N1N	3.36	1.39	1.35
3	A	1800	NAP	C2A-N3A	3.25	1.37	1.32

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1800	NAP	N3A-C2A-N1A	-5.53	120.04	128.68
3	B	1801	NAP	O7N-C7N-N7N	-3.65	117.39	122.58
3	A	1800	NAP	O7N-C7N-N7N	-3.54	117.55	122.58
3	B	1801	NAP	N3A-C2A-N1A	-3.46	123.28	128.68
3	A	1800	NAP	O7N-C7N-C3N	2.83	123.02	119.63

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1800	NAP	O4D-C1D-N1N-C2N
3	A	1800	NAP	O4D-C1D-N1N-C6N
3	A	1800	NAP	C2D-C1D-N1N-C2N
3	A	1800	NAP	C2D-C1D-N1N-C6N
3	B	1801	NAP	O4D-C1D-N1N-C2N


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1800	NAP	2	0
3	B	1801	NAP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates ⓘ


Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.