



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 02:44 am BST

PDB ID : 6O30  
Title : Lipid A transporter MsbA from Salmonella typhimurium  
Authors : Padayatti, P.S.; Zhang, Q.; Wilson, I.A.; Lee, S.C.; Stanfield, R.L.  
Deposited on : 2019-02-25  
Resolution : 4.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 4.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1051 (5.12-3.80)
Clashscore	141614	1119 (5.12-3.80)
Ramachandran outliers	138981	1065 (5.12-3.80)
Sidechain outliers	138945	1047 (5.12-3.80)
RSRZ outliers	127900	1099 (5.20-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -10px; left: 0;">%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>90%</span> <span>10%</span> </div> </div>
1	B	576	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -10px; left: 0;">2%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>89%</span> <span>11%</span> </div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lipid A export ATP-binding/permease protein MsbA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	575	Total	C	N	O	S	0	0	0
			4447	2823	770	827	27			
1	B	575	Total	C	N	O	S	0	0	0
			4447	2823	770	827	27			


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lipid A export ATP-binding/permease protein MsbA



- Molecule 1: Lipid A export ATP-binding/permease protein MsbA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.26Å 153.20Å 232.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.75 – 4.47 46.75 – 4.46	Depositor EDS
% Data completeness (in resolution range)	80.7 (46.75-4.47) 83.0 (46.75-4.46)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 4.45Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, $R_{free}$	0.307 , 0.340 0.311 , 0.350	Depositor DCC
$R_{free}$ test set	742 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	184.6	Xtriage
Anisotropy	0.730	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 169.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	8894	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	271.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/4514	0.49	0/6101
1	B	0.27	0/4514	0.49	0/6101
All	All	0.27	0/9028	0.49	0/12202

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4447	0	4568	21	0
1	B	4447	0	4568	23	0
All	All	8894	0	9136	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:474:ILE:HG22	1:A:478:GLY:HA2	1.82	0.60
1:B:49:LYS:NZ	1:B:53:ASP:OD2	2.36	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:474:ILE:HG22	1:B:478:GLY:HA2	1.87	0.57
1:A:550:VAL:HG11	1:A:571:VAL:HG23	1.87	0.56
1:B:37:ASN:OD1	1:B:78:ARG:HG2	2.07	0.54
1:B:175:ALA:HB3	1:B:176:PRO:HD3	1.90	0.54
1:B:350:THR:HG22	1:B:357:PRO:HA	1.91	0.53
1:B:550:VAL:HG11	1:B:571:VAL:HG23	1.91	0.53
1:A:412:LEU:HD22	1:A:416:ARG:CZ	2.40	0.52
1:B:281:ALA:O	1:B:285:THR:HG23	2.10	0.51
1:B:412:LEU:HD22	1:B:416:ARG:CZ	2.39	0.51
1:A:350:THR:HG22	1:A:357:PRO:HA	1.93	0.51
1:A:281:ALA:O	1:A:285:THR:HG23	2.11	0.50
1:B:67:MET:HB3	1:B:68:PRO:HD3	1.93	0.50
1:A:67:MET:HB3	1:A:68:PRO:HD3	1.95	0.49
1:A:218:LEU:HD23	1:A:223:GLN:HE21	1.77	0.49
1:B:63:VAL:HA	1:B:66:TRP:NE1	2.28	0.49
1:A:175:ALA:HB3	1:A:176:PRO:HD3	1.95	0.48
1:A:558:GLU:HB3	1:A:565:LEU:HD22	1.96	0.48
1:A:38:ALA:O	1:A:42:THR:HG23	2.14	0.48
1:A:37:ASN:OD1	1:A:78:ARG:HG2	2.14	0.48
1:A:238:ARG:O	1:A:242:MET:HG2	2.14	0.48
1:A:45:LEU:HD11	1:A:289:SER:HB3	1.95	0.48
1:B:38:ALA:O	1:B:42:THR:HG23	2.14	0.48
1:B:218:LEU:HD23	1:B:223:GLN:HE21	1.80	0.47
1:A:131:ASP:O	1:A:135:VAL:HG23	2.15	0.47
1:B:238:ARG:O	1:B:242:MET:HG2	2.15	0.46
1:A:49:LYS:NZ	1:A:53:ASP:OD2	2.45	0.46
1:B:280:THR:OG1	1:B:283:THR:HG23	2.15	0.46
1:A:63:VAL:HA	1:A:66:TRP:NE1	2.33	0.44
1:A:280:THR:OG1	1:A:283:THR:HG23	2.17	0.44
1:B:174:LEU:HD11	1:B:265:PHE:CD2	2.53	0.44
1:B:174:LEU:HD11	1:B:265:PHE:HD2	1.81	0.44
1:B:37:ASN:HD22	1:B:148:ARG:HE	1.65	0.44
1:B:558:GLU:HB3	1:B:565:LEU:HD22	1.99	0.44
1:B:469:GLY:O	1:B:472:THR:OG1	2.36	0.43
1:B:131:ASP:O	1:B:135:VAL:HG23	2.17	0.43
1:B:45:LEU:HD11	1:B:289:SER:HB3	2.00	0.43
1:A:14:ARG:HA	1:A:14:ARG:HD2	1.79	0.42
1:A:174:LEU:HD11	1:A:265:PHE:HD2	1.83	0.42
1:A:174:LEU:HD11	1:A:265:PHE:CD2	2.54	0.42
1:B:283:THR:O	1:B:287:VAL:HG23	2.21	0.41
1:B:93:SER:HB2	1:B:136:ALA:HB1	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:469:GLY:O	1:A:472:THR:OG1	2.38	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	573/576 (100%)	537 (94%)	30 (5%)	6 (1%)	15	54
1	B	573/576 (100%)	535 (93%)	32 (6%)	6 (1%)	15	54
All	All	1146/1152 (100%)	1072 (94%)	62 (5%)	12 (1%)	15	54

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	513	THR
1	B	513	THR
1	A	356	VAL
1	B	356	VAL
1	A	164	SER
1	B	164	SER
1	B	379	GLY
1	B	439	TYR
1	A	379	GLY
1	A	439	TYR
1	A	475	GLY
1	B	475	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	485/487 (100%)	470 (97%)	15 (3%)	40	63
1	B	485/487 (100%)	469 (97%)	16 (3%)	38	61
All	All	970/974 (100%)	939 (97%)	31 (3%)	39	62

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	MET
1	A	242	MET
1	A	252	ASP
1	A	260	SER
1	A	274	SER
1	A	341	ASP
1	A	371	THR
1	A	377	ARG
1	A	383	SER
1	A	394	ASP
1	A	436	ASN
1	A	477	ASN
1	A	513	THR
1	A	532	SER
1	A	546	ASP
1	B	44	MET
1	B	252	ASP
1	B	260	SER
1	B	274	SER
1	B	341	ASP
1	B	371	THR
1	B	377	ARG
1	B	382	LYS
1	B	383	SER
1	B	394	ASP
1	B	425	ASN
1	B	436	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	477	ASN
1	B	513	THR
1	B	532	SER
1	B	546	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	575/576 (99%)	-0.20	5 (0%) 84 77	168, 258, 319, 375	0
1	B	575/576 (99%)	-0.15	11 (1%) 66 58	206, 280, 327, 365	0
All	All	1150/1152 (99%)	-0.17	16 (1%) 75 66	168, 269, 324, 375	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	569	HIS	5.1
1	B	233	VAL	4.4
1	B	237	MET	3.9
1	B	457	ALA	2.9
1	A	513	THR	2.8
1	B	456	MET	2.7
1	B	108	MET	2.6
1	A	86	SER	2.6
1	B	450	ILE	2.3
1	B	234	SER	2.3
1	B	163	TYR	2.2
1	B	100	MET	2.1
1	B	488	ARG	2.1
1	B	119	GLN	2.1
1	A	437	ILE	2.1
1	A	477	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.