



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 08:38 AM BST

PDB ID : 4O5N  
Title : Crystal structure of A/Victoria/361/2011 (H3N2) influenza virus hemagglutinin  
Authors : Lee, P.S.; Wilson, I.A.  
Deposited on : 2013-12-19  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	<div> <div>7%</div> <div>96%</div> <div>..</div> </div>
2	B	176	<div> <div>2%</div> <div>94%</div> <div>...</div> </div>
3	C	2	<div> <div>100%</div> </div>
4	D	3	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	2	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	5	0
			2498	1570	441	475	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	expression tag	UNP R9U684
A	8	ASP	-	expression tag	UNP R9U684
A	9	PRO	-	expression tag	UNP R9U684
A	10	GLY	-	expression tag	UNP R9U684

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	173	Total	C	N	O	S	0	7	0
			1431	899	248	277	7			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	222	Total	O	0	0
			222	222		
8	B	204	Total	O	0	0
			204	204		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemagglutinin HA1 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.69Å 100.69Å 383.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.84 – 1.75 46.84 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.84-1.75) 99.8 (46.84-1.75)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.26 (at 1.75Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.165 , 0.185 0.166 , 0.186	Depositor DCC
$R_{free}$ test set	3824 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.2	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4533	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2570	0.55	0/3495
2	B	0.49	0/1476	0.67	2/1982 (0.1%)
All	All	0.43	0/4046	0.60	2/5477 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	54	ARG	NE-CZ-NH2	-6.65	116.98	120.30
2	B	54	ARG	NE-CZ-NH1	6.29	123.44	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2498	0	2463	4	0
2	B	1431	0	1393	8	0
3	C	28	0	25	0	0
4	D	39	0	34	0	0
5	A	56	0	52	0	0
6	A	12	0	16	0	0
6	B	36	0	48	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	7	0	10	3	0
8	A	222	0	0	2	0
8	B	204	0	0	2	0
All	All	4533	0	4041	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48[B]:ILE:HG13	8:A:718:HOH:O	1.96	0.64
1:A:48[B]:ILE:HD13	1:A:50[B]:GLU:HG2	1.81	0.61
2:B:20:GLY:O	7:B:201:PEG:H42	2.03	0.59
1:A:197:GLN:NE2	1:A:248:THR:O	2.37	0.58
2:B:134:GLY:O	6:B:205:GOL:H11	2.04	0.57
2:B:54:ARG:HD2	6:B:202:GOL:O1	2.06	0.56
2:B:135:ASN:HB3	6:B:205:GOL:H31	1.90	0.53
2:B:18:VAL:C	7:B:201:PEG:H41	2.32	0.50
6:B:205:GOL:H12	8:B:392:HOH:O	2.12	0.49
2:B:46:ASP:HB2	6:B:206:GOL:H32	2.01	0.43
1:A:48[A]:ILE:HG23	8:A:718:HOH:O	2.18	0.42
7:B:201:PEG:H21	8:B:403:HOH:O	2.21	0.41
2:B:54:ARG:NH1	6:B:202:GOL:H31	2.36	0.41
2:B:39:LYS:HG2	6:B:207:GOL:O2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/323 (99%)	311 (97%)	9 (3%)	0	100	100
2	B	178/176 (101%)	170 (96%)	8 (4%)	0	100	100
All	All	498/499 (100%)	481 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/285 (100%)	282 (99%)	3 (1%)	73	60
2	B	153/148 (103%)	153 (100%)	0	100	100
All	All	438/433 (101%)	435 (99%)	3 (1%)	84	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	246	ASN
1	A	325	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	246	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	C	1	1,3	14,14,15	0.55	0	17,19,21	0.58	0
3	NAG	C	2	3	14,14,15	0.54	0	17,19,21	0.68	0
4	NAG	D	1	1,4	14,14,15	0.51	0	17,19,21	0.84	1 (5%)
4	NAG	D	2	4	14,14,15	0.55	0	17,19,21	0.77	1 (5%)
4	BMA	D	3	4	11,11,12	0.61	0	15,15,17	0.93	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	C1-O5-C5	2.31	115.32	112.19
4	D	3	BMA	C1-C2-C3	2.30	112.50	109.67
4	D	3	BMA	O5-C5-C6	2.24	110.71	107.20
4	D	2	NAG	O5-C5-C6	2.16	110.60	107.20

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	3	BMA	O5-C5-C6-O6
4	D	3	BMA	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	GOL	B	207	-	5,5,5	0.38	0	5,5,5	0.60	0
5	NAG	A	408	1	14,14,15	0.62	0	17,19,21	1.10	1 (5%)
6	GOL	A	410	-	5,5,5	0.37	0	5,5,5	0.42	0
6	GOL	B	203	-	5,5,5	0.37	0	5,5,5	0.41	0
7	PEG	B	201	-	6,6,6	0.43	0	5,5,5	0.81	0
6	GOL	B	206	-	5,5,5	0.34	0	5,5,5	0.22	0
5	NAG	A	409	1	14,14,15	0.53	0	17,19,21	0.64	1 (5%)
6	GOL	B	205	-	5,5,5	0.37	0	5,5,5	0.66	0
6	GOL	B	202	-	5,5,5	0.33	0	5,5,5	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	401	1	14,14,15	0.48	0	17,19,21	1.09	1 (5%)
6	GOL	A	411	-	5,5,5	0.40	0	5,5,5	0.42	0
6	GOL	B	204	-	5,5,5	0.44	0	5,5,5	0.40	0
5	NAG	A	402	1	14,14,15	0.49	0	17,19,21	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	207	-	-	2/4/4/4	-
5	NAG	A	408	1	-	0/6/23/26	0/1/1/1
6	GOL	A	410	-	-	2/4/4/4	-
6	GOL	B	203	-	-	2/4/4/4	-
7	PEG	B	201	-	-	1/4/4/4	-
6	GOL	B	206	-	-	2/4/4/4	-
5	NAG	A	409	1	-	0/6/23/26	0/1/1/1
6	GOL	B	205	-	-	2/4/4/4	-
6	GOL	B	202	-	-	3/4/4/4	-
5	NAG	A	401	1	-	2/6/23/26	0/1/1/1
6	GOL	A	411	-	-	4/4/4/4	-
6	GOL	B	204	-	-	4/4/4/4	-
5	NAG	A	402	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401	NAG	C1-O5-C5	2.96	116.20	112.19
5	A	402	NAG	O5-C1-C2	-2.88	106.75	111.29
5	A	402	NAG	C1-O5-C5	2.68	115.82	112.19
5	A	408	NAG	O7-C7-C8	-2.03	118.28	122.06
5	A	409	NAG	O5-C5-C6	2.01	110.35	107.20

There are no chirality outliers.

All (24) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	B	206	GOL	C1-C2-C3-O3
6	B	205	GOL	O1-C1-C2-C3
5	A	401	NAG	C8-C7-N2-C2
5	A	401	NAG	O7-C7-N2-C2
6	A	411	GOL	O1-C1-C2-C3
6	A	411	GOL	C1-C2-C3-O3
6	B	204	GOL	O1-C1-C2-C3
6	B	204	GOL	O2-C2-C3-O3
6	A	410	GOL	C1-C2-C3-O3
6	B	203	GOL	O1-C1-C2-C3
6	B	202	GOL	O1-C1-C2-C3
6	B	204	GOL	C1-C2-C3-O3
6	A	410	GOL	O2-C2-C3-O3
6	B	203	GOL	O1-C1-C2-O2
6	B	205	GOL	O1-C1-C2-O2
6	A	411	GOL	O2-C2-C3-O3
6	B	204	GOL	O1-C1-C2-O2
6	B	206	GOL	O2-C2-C3-O3
6	B	202	GOL	O1-C1-C2-O2
6	B	202	GOL	O2-C2-C3-O3
6	A	411	GOL	O1-C1-C2-O2
6	B	207	GOL	O2-C2-C3-O3
7	B	201	PEG	O1-C1-C2-O2
6	B	207	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	207	GOL	1	0
7	B	201	PEG	3	0
6	B	206	GOL	1	0
6	B	205	GOL	3	0
6	B	202	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	317/323 (98%)	0.10	21 (6%) 18 24	16, 36, 67, 83	0
2	B	173/176 (98%)	-0.10	3 (1%) 70 77	14, 23, 40, 76	0
All	All	490/499 (98%)	0.03	24 (4%) 29 35	14, 28, 63, 83	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	193	PHE	5.7
1	A	189	LYS	4.5
1	A	142	ARG	4.0
1	A	157	LEU	3.9
1	A	196	ALA	3.9
1	A	223	ILE	3.6
1	A	48[A]	ILE	3.5
1	A	192	ILE	3.1
1	A	173	GLN	3.1
2	B	173	ILE	3.0
1	A	159	PHE	2.9
1	A	158	ASN	2.8
1	A	225	ASN	2.8
1	A	144	ASN	2.6
1	A	163	ALA	2.5
1	A	143	SER	2.5
2	B	2	ILE	2.5
1	A	187	THR	2.4
1	A	186	GLY	2.2
1	A	226	ILE	2.2
1	A	162	PRO	2.2
1	A	156	HIS	2.1
2	B	3	PHE	2.1
1	A	160	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	C	2	14/15	0.77	0.46	88,97,103,105	0
4	BMA	D	3	11/12	0.79	0.32	90,102,107,111	0
4	NAG	D	2	14/15	0.88	0.23	59,67,76,78	0
3	NAG	C	1	14/15	0.89	0.28	58,65,76,79	0
4	NAG	D	1	14/15	0.95	0.15	56,60,71,72	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	409	14/15	0.69	0.25	85,97,106,110	0
5	NAG	A	401	14/15	0.73	0.38	79,93,98,103	0
6	GOL	A	410	6/6	0.77	0.17	30,49,52,55	0
7	PEG	B	201	7/7	0.80	0.19	31,46,60,65	0
6	GOL	B	202	6/6	0.81	0.14	43,60,63,67	0
6	GOL	B	206	6/6	0.85	0.30	35,61,66,74	0
6	GOL	B	205	6/6	0.85	0.22	36,58,63,68	0
6	GOL	A	411	6/6	0.85	0.14	66,70,72,72	0
6	GOL	B	207	6/6	0.86	0.23	30,57,61,62	0
5	NAG	A	402	14/15	0.87	0.18	34,43,50,51	0
6	GOL	B	204	6/6	0.89	0.12	52,61,63,71	0
5	NAG	A	408	14/15	0.92	0.17	31,40,50,54	0
6	GOL	B	203	6/6	0.95	0.22	31,59,65,73	0

## 6.5 Other polymers

There are no such residues in this entry.