



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 7, 2022 – 04:33 PM EDT

PDB ID : 2O6T  
Title : Crystal structure of the PA5185 protein from *Pseudomonas Aeruginosa* strain PAO1- orthorhombic form (P2221).  
Authors : Chruszcz, M.; Koclega, K.D.; Evdokimova, E.; Cymborowski, M.; Kudritska, M.; Savchenko, A.; Edwards, A.; Minor, W.  
Deposited on : 2006-12-08  
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.27
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	<div> <div>2%</div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	C	149	<div> <div>2%</div> <div>81%</div> <div>15%</div> <div>..</div> </div>
1	E	149	<div> <div>3%</div> <div>85%</div> <div>8%</div> <div>6%</div> </div>
1	G	149	<div> <div>3%</div> <div>74%</div> <div>17%</div> <div>.. 5%</div> </div>
1	I	149	<div> <div>3%</div> <div>76%</div> <div>21%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	149	<div><div><div>%</div><div><div></div></div><div>78%</div><div>18%</div><div>.</div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THIOESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	2	0
			1139	726	200	209	4			
1	C	143	Total	C	N	O	S	0	1	0
			1123	715	198	207	3			
1	E	140	Total	C	N	O	S	0	1	0
			1105	704	195	203	3			
1	G	142	Total	C	N	O	S	0	1	0
			1110	709	196	202	3			
1	I	145	Total	C	N	O	S	0	2	0
			1122	718	195	205	4			
1	K	143	Total	C	N	O	S	0	1	0
			1105	708	189	205	3			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q9HU04
A	0	HIS	-	expression tag	UNP Q9HU04
C	-1	GLY	-	expression tag	UNP Q9HU04
C	0	HIS	-	expression tag	UNP Q9HU04
E	-1	GLY	-	expression tag	UNP Q9HU04
E	0	HIS	-	expression tag	UNP Q9HU04
G	-1	GLY	-	expression tag	UNP Q9HU04
G	0	HIS	-	expression tag	UNP Q9HU04
I	-1	GLY	-	expression tag	UNP Q9HU04
I	0	HIS	-	expression tag	UNP Q9HU04
K	-1	GLY	-	expression tag	UNP Q9HU04
K	0	HIS	-	expression tag	UNP Q9HU04

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Cl 1	0	0
2	E	1	Total 1	Cl 1	0	0
2	G	1	Total 1	Cl 1	0	0
2	K	1	Total 1	Cl 1	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	48	Total 48	O 48	0	0
3	C	49	Total 49	O 49	0	0
3	E	42	Total 42	O 42	0	0
3	G	34	Total 34	O 34	0	0
3	I	43	Total 43	O 43	0	0
3	K	35	Total 35	O 35	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: THIOESTERASE




#### • Molecule 1: THIOESTERASE




#### • Molecule 1: THIOESTERASE




#### • Molecule 1: THIOESTERASE



#### • Molecule 1: THIOESTERASE



● Molecule 1: THIOESTERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.92Å 97.08Å 191.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.53 – 2.55 37.47 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.53-2.55) 99.4 (37.47-2.55)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.180 , 0.248 0.199 , 0.256	Depositor DCC
$R_{free}$ test set	1810 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6959	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.89	0/1166	0.82	0/1587
1	C	0.89	0/1150	0.85	1/1564 (0.1%)
1	E	0.94	0/1132	0.88	0/1539
1	G	0.95	0/1137	0.91	3/1547 (0.2%)
1	I	0.85	0/1152	0.85	0/1569
1	K	1.02	0/1132	0.84	1/1541 (0.1%)
All	All	0.93	0/6869	0.86	5/9347 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	4

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	123	ARG	NE-CZ-NH2	-7.37	116.61	120.30
1	G	108	ARG	NE-CZ-NH1	6.29	123.45	120.30
1	C	123	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	G	21	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	K	137	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	143	GLN	Peptide
1	G	144	SER	Peptide
1	G	4	ALA	Peptide
1	G	5	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1139	0	1094	10	0
1	C	1123	0	1083	22	0
1	E	1105	0	1065	6	0
1	G	1110	0	1067	19	0
1	I	1122	0	1077	24	0
1	K	1105	0	1052	23	0
2	A	1	0	0	0	0
2	E	1	0	0	0	0
2	G	1	0	0	0	0
2	K	1	0	0	0	0
3	A	48	0	0	0	0
3	C	49	0	0	0	0
3	E	42	0	0	1	0
3	G	34	0	0	1	0
3	I	43	0	0	2	0
3	K	35	0	0	0	0
All	All	6959	0	6438	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:5:PRO:CB	1:C:6:ARG:HG3	1.61	1.30
1:C:5:PRO:HB2	1:C:6:ARG:CG	1.63	1.28
1:C:5:PRO:CB	1:C:6:ARG:HA	1.71	1.17
1:G:4:ALA:N	1:G:5:PRO:HD3	1.59	1.12

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:5:PRO:HB3	1:C:6:ARG:HA	1.23	1.09

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/149 (97%)	139 (96%)	5 (3%)	1 (1%)	22	30
1	C	141/149 (95%)	139 (99%)	1 (1%)	1 (1%)	22	30
1	E	138/149 (93%)	137 (99%)	1 (1%)	0	100	100
1	G	140/149 (94%)	138 (99%)	1 (1%)	1 (1%)	22	30
1	I	144/149 (97%)	138 (96%)	6 (4%)	0	100	100
1	K	141/149 (95%)	135 (96%)	6 (4%)	0	100	100
All	All	849/894 (95%)	826 (97%)	20 (2%)	3 (0%)	34	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	5	PRO
1	C	5	PRO
1	A	144	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/122 (95%)	112 (97%)	4 (3%)	37	50
1	C	116/122 (95%)	111 (96%)	5 (4%)	29	39
1	E	115/122 (94%)	109 (95%)	6 (5%)	23	30
1	G	113/122 (93%)	106 (94%)	7 (6%)	18	24
1	I	113/122 (93%)	110 (97%)	3 (3%)	44	59
1	K	112/122 (92%)	110 (98%)	2 (2%)	59	74
All	All	685/732 (94%)	658 (96%)	27 (4%)	31	44

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	122	GLU
1	G	91	LEU
1	I	126	SER
1	G	60	GLU
1	G	108	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	143	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	146/149 (97%)	-0.13	3 (2%) 63 70	10, 20, 33, 42	0
1	C	143/149 (95%)	-0.07	3 (2%) 63 70	10, 20, 32, 46	0
1	E	140/149 (93%)	-0.13	4 (2%) 51 59	9, 21, 31, 39	0
1	G	142/149 (95%)	-0.03	4 (2%) 53 60	10, 21, 33, 44	0
1	I	145/149 (97%)	-0.15	4 (2%) 53 60	11, 20, 30, 41	0
1	K	143/149 (95%)	-0.10	1 (0%) 87 90	10, 20, 30, 43	0
All	All	859/894 (96%)	-0.10	19 (2%) 62 68	9, 20, 33, 46	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	GLY	4.6
1	G	4	ALA	4.3
1	G	106	GLY	3.8
1	E	5	PRO	3.5
1	C	4	ALA	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CL	K	201	1/1	0.80	0.35	67,67,67,67	0
2	CL	G	201	1/1	0.84	0.20	63,63,63,63	0
2	CL	E	201	1/1	0.87	0.24	51,51,51,51	0
2	CL	A	201	1/1	0.93	0.39	60,60,60,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.