



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 02:47 pm BST

PDB ID : 2O7G
Title : Crystal structure of the Pribnow Box recognition region of SigC from Mycobacterium tuberculosis
Authors : Thakur, K.G.; Joshi, A.M.; Gopal, B.
Deposited on : 2006-12-11
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	<div> <div>5%</div> <div> <div></div> <div>53%</div> <div>16%</div> <div>9%</div> <div>21%</div> </div> </div>
1	B	112	<div> <div>3%</div> <div> <div></div> <div>58%</div> <div>13%</div> <div>8%</div> <div>21%</div> </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable RNA polymerase sigma-C factor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	88	Total	C	N	O	0	0	0
			670	421	124	125			
1	B	89	Total	C	N	O	0	0	0
			672	422	125	125			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total 13	O 13	0	0
3	B	7	Total 7	O 7	0	0

- Molecule 1: Probable RNA polymerase sigma-C factor



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	85.17Å 85.17Å 79.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.46 – 2.70 33.46 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.46-2.70) 100.0 (33.46-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.2.0009	Depositor
R, R_{free}	0.201 , 0.248 0.220 , 0.266	Depositor DCC
R_{free} test set	418 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.084 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1382	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.55	8/681 (1.2%)	1.42	14/924 (1.5%)
1	B	1.40	2/683 (0.3%)	1.18	7/926 (0.8%)
All	All	1.48	10/1364 (0.7%)	1.30	21/1850 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	43	TYR	CB-CG	-15.10	1.29	1.51
1	A	43	TYR	CD1-CE1	-13.33	1.19	1.39
1	A	43	TYR	CD2-CE2	-10.81	1.23	1.39
1	B	27	GLU	CD-OE2	-6.23	1.18	1.25
1	A	43	TYR	N-CA	-5.76	1.34	1.46
1	A	83	ARG	CG-CD	-5.71	1.37	1.51
1	B	27	GLU	CD-OE1	-5.55	1.19	1.25
1	A	43	TYR	CZ-OH	-5.23	1.28	1.37
1	A	83	ARG	CB-CG	-5.13	1.38	1.52
1	A	43	TYR	CA-CB	-5.12	1.42	1.53

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	83	ARG	NE-CZ-NH1	13.30	126.95	120.30
1	A	83	ARG	NE-CZ-NH1	9.83	125.22	120.30
1	B	83	ARG	NE-CZ-NH2	-9.01	115.80	120.30
1	A	43	TYR	N-CA-CB	-8.81	94.75	110.60
1	A	83	ARG	NE-CZ-NH2	-8.74	115.93	120.30
1	B	39	ARG	NE-CZ-NH1	-8.61	116.00	120.30
1	A	56	GLU	N-CA-C	-8.56	87.90	111.00
1	B	39	ARG	NE-CZ-NH2	8.43	124.51	120.30
1	A	24	ARG	NE-CZ-NH2	8.32	124.46	120.30
1	A	39	ARG	NE-CZ-NH2	-8.27	116.16	120.30
1	B	27	GLU	OE1-CD-OE2	-8.16	113.51	123.30
1	A	39	ARG	NE-CZ-NH1	7.65	124.12	120.30
1	A	60	ARG	NE-CZ-NH2	7.47	124.03	120.30
1	B	60	ARG	NE-CZ-NH2	-7.46	116.57	120.30
1	A	24	ARG	CG-CD-NE	-6.17	98.84	111.80
1	A	60	ARG	NE-CZ-NH1	-6.09	117.25	120.30
1	A	43	TYR	CZ-CE2-CD2	-5.92	114.47	119.80
1	A	43	TYR	CB-CG-CD2	-5.66	117.60	121.00
1	B	60	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	43	TYR	CD1-CG-CD2	5.19	123.61	117.90
1	A	83	ARG	CA-CB-CG	-5.12	102.14	113.40

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	4	THR	Peptide
1	A	55	GLN	Peptide
1	B	3	ALA	Peptide
1	B	90	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	670	0	667	15	0
1	B	672	0	666	18	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	15	0	0	0	0
3	A	13	0	0	1	0
3	B	7	0	0	0	0
All	All	1382	0	1333	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:ALA:O	1:B:6:SER:CB	2.00	1.07
1:B:3:ALA:O	1:B:6:SER:N	2.08	0.87
1:B:3:ALA:O	1:B:6:SER:HB2	1.74	0.86
1:B:3:ALA:O	1:B:6:SER:HB3	1.76	0.84
1:A:88:ASP:HA	1:A:91:ARG:HH11	1.42	0.83
1:A:44:LEU:HB3	1:A:86:VAL:HG11	1.80	0.61
1:B:22:ASN:HD22	1:B:22:ASN:C	2.02	0.61
1:B:22:ASN:HD22	1:B:23:GLY:N	1.98	0.61
1:B:3:ALA:O	1:B:6:SER:CA	2.49	0.60
1:A:80:ALA:O	1:A:84:HIS:HD2	1.85	0.59
1:A:41:VAL:C	1:A:43:TYR:H	2.09	0.56
1:A:4:THR:HA	1:A:7:ASP:HB2	1.88	0.54
1:B:44:LEU:HB3	1:B:86:VAL:HG11	1.90	0.54
1:A:65:ILE:N	1:A:66:PRO:CD	2.69	0.53
1:B:22:ASN:ND2	1:B:22:ASN:C	2.63	0.52
1:B:4:THR:O	1:B:7:ASP:HB2	2.15	0.46
1:B:80:ALA:O	1:B:84:HIS:HD2	1.99	0.46
1:B:60:ARG:CZ	1:B:85:VAL:HG22	2.46	0.45
1:B:80:ALA:HA	1:B:83:ARG:NH1	2.32	0.45
1:A:41:VAL:C	1:A:43:TYR:N	2.70	0.45
1:B:4:THR:O	1:B:7:ASP:CB	2.65	0.45
1:A:37:VAL:O	1:A:41:VAL:HB	2.17	0.45
1:A:55:GLN:HB3	1:A:55:GLN:HE21	1.55	0.45
1:A:39:ARG:O	1:A:43:TYR:HB3	2.16	0.44
1:A:42:ALA:HA	1:A:50:ALA:HB2	2.00	0.43
1:A:60:ARG:HG2	1:A:60:ARG:H	1.68	0.43
1:A:56:GLU:OE2	1:A:60:ARG:HD3	2.19	0.42
1:B:36:ASP:OD2	1:B:75:ARG:NH2	2.53	0.42
1:B:3:ALA:HA	1:B:5:ALA:H	1.84	0.42
1:A:34:GLN:HG2	3:A:119:HOH:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ALA:HA	1:A:83:ARG:HH11	1.85	0.41
1:B:3:ALA:C	1:B:5:ALA:N	2.75	0.41
1:B:3:ALA:C	1:B:6:SER:H	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	86/112 (77%)	80 (93%)	6 (7%)	0	100	100
1	B	87/112 (78%)	83 (95%)	3 (3%)	1 (1%)	14	34
All	All	173/224 (77%)	163 (94%)	9 (5%)	1 (1%)	25	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	4	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/86 (76%)	53 (82%)	12 (18%)	1	4
1	B	65/86 (76%)	55 (85%)	10 (15%)	2	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	130/172 (76%)	108 (83%)	22 (17%)	2 5

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASP
1	A	20	LYS
1	A	24	ARG
1	A	26	LEU
1	A	41	VAL
1	A	44	LEU
1	A	53	LEU
1	A	59	LEU
1	A	60	ARG
1	A	71	ARG
1	A	75	ARG
1	A	91	ARG
1	B	7	ASP
1	B	22	ASN
1	B	24	ARG
1	B	44	LEU
1	B	49	SER
1	B	53	LEU
1	B	71	ARG
1	B	75	ARG
1	B	90	ILE
1	B	91	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	84	HIS
1	B	22	ASN
1	B	34	GLN
1	B	84	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	114	-	4,4,4	0.16	0	6,6,6	0.40	0
2	SO4	B	113	-	4,4,4	0.19	0	6,6,6	0.58	0
2	SO4	B	115	-	4,4,4	0.18	0	6,6,6	0.22	0
2	SO4	A	113	-	4,4,4	0.39	0	6,6,6	0.51	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	88/112 (78%)	0.03	6 (6%) 17 15	20, 34, 47, 64	0
1	B	89/112 (79%)	-0.01	3 (3%) 45 45	20, 33, 51, 62	0
All	All	177/224 (79%)	0.01	9 (5%) 28 26	20, 34, 51, 64	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	SER	3.9
1	B	91	ARG	3.8
1	B	3	ALA	3.8
1	B	6	SER	3.7
1	A	91	ARG	3.6
1	A	5	ALA	3.6
1	A	4	THR	2.9
1	A	71	ARG	2.7
1	A	90	ILE	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	113	5/5	0.75	0.28	113,114,115,115	0
2	SO4	B	115	5/5	0.89	0.34	101,101,102,102	0
2	SO4	B	114	5/5	0.91	0.35	86,87,87,89	0
2	SO4	B	113	5/5	0.92	0.22	100,101,102,102	0

6.5 Other polymers [i](#)

There are no such residues in this entry.