



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:26 pm BST

PDB ID : 4O87
Title : Crystal structure of a N-tagged Nuclease
Authors : Chakravarty, A.K.; Smith, P.; Shuman, S.
Deposited on : 2013-12-26
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	<div> <div>3%</div> <div>87%</div> <div>12%</div> <div>..</div> </div>
1	B	320	<div> <div>8%</div> <div>88%</div> <div>12%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	505	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6166 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-tagged Nuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	27	0
			2666	1678	456	526	6			
1	B	319	Total	C	N	O	S	0	15	0
			2615	1640	450	519	6			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP Q707V3
A	1	MET	-	EXPRESSION TAG	UNP Q707V3
B	0	SER	-	EXPRESSION TAG	UNP Q707V3
B	1	MET	-	EXPRESSION TAG	UNP Q707V3

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	434	Total	O	0	0
			434	434		
4	B	290	Total	O	0	0
			290	290		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: N-tagged Nuclease



• Molecule 1: N-tagged Nuclease



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.87Å 77.43Å 147.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.53 – 1.80 38.53 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.53-1.80) 99.9 (38.53-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 1.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.159 , 0.199 0.162 , 0.199	Depositor DCC
R_{free} test set	2526 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.652	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6166	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8460e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	1/2794 (0.0%)	0.68	0/3753
1	B	0.55	0/2701	0.61	0/3631
All	All	0.60	1/5495 (0.0%)	0.65	0/7384

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	14	TYR	CD2-CE2	6.07	1.48	1.39

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2666	0	2731	33	1
1	B	2615	0	2666	31	0
2	A	13	0	5	2	0
2	B	13	0	5	2	0
3	A	80	0	0	4	0
3	B	55	0	0	3	0
4	A	434	0	0	19	1
4	B	290	0	0	9	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6166	0	5407	69	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:505:SO4:O3	4:B:750:HOH:O	1.89	0.90
1:A:174:LYS:HD3	1:A:265[B]:LEU:HD11	1.61	0.83
1:A:197[B]:ASN:ND2	4:A:936:HOH:O	2.18	0.76
1:A:130:ASN:ND2	4:A:937:HOH:O	2.21	0.73
1:A:285[B]:ARG:NH2	4:A:1031:HOH:O	2.20	0.72
1:A:157[A]:GLN:OE1	4:A:823:HOH:O	2.09	0.70
1:B:97:LYS:N	3:B:505:SO4:O2	2.19	0.70
1:A:184[B]:GLU:OE1	4:A:1025:HOH:O	2.08	0.69
1:A:247[B]:ASN:OD1	4:A:740:HOH:O	2.09	0.68
1:A:157[A]:GLN:NE2	4:A:675:HOH:O	2.25	0.68
1:B:317:CYS:SG	1:B:318:LYS:N	2.68	0.66
1:A:247[B]:ASN:ND2	4:A:1011:HOH:O	2.29	0.66
1:B:32:ASP:OD1	4:B:667:HOH:O	2.15	0.63
1:B:317:CYS:O	1:B:318:LYS:HB2	2.01	0.61
1:B:307:ASN:ND2	4:B:864:HOH:O	2.19	0.60
1:B:241:ASN:OD1	1:B:241:ASN:N	2.32	0.60
1:B:29:THR:OG1	1:B:34:GLU:OE1	2.12	0.60
1:A:174:LYS:CD	1:A:265[B]:LEU:HD11	2.30	0.60
1:A:45:ASP:O	1:A:49:LYS:HG2	2.01	0.60
1:B:29:THR:HB	1:B:316:THR:HG22	1.83	0.59
1:B:135:LYS:HD3	1:B:135:LYS:H	1.66	0.59
1:A:247[B]:ASN:ND2	4:A:979:HOH:O	2.34	0.59
3:A:513:SO4:O1	4:A:998:HOH:O	2.17	0.58
1:B:305:ASN:HD21	1:B:316:THR:HG23	1.70	0.57
1:A:157[B]:GLN:NE2	4:A:834:HOH:O	2.39	0.55
1:A:123[B]:ASN:OD1	4:A:672:HOH:O	2.18	0.54
1:B:206[B]:GLN:HG3	1:B:212:THR:HG22	1.89	0.54
1:B:56:VAL:HG21	1:B:115:LEU:HD21	1.90	0.54
1:A:157[B]:GLN:NE2	4:A:868:HOH:O	2.00	0.51
1:B:81:LYS:HE2	4:B:747:HOH:O	2.10	0.51
1:B:135:LYS:CD	1:B:135:LYS:H	2.23	0.50
1:A:142:LEU:HD23	1:A:290[A]:LEU:HD11	1.93	0.49
1:A:310:SER:HB3	1:A:311:LYS:HA	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:ASN:HB3	4:B:833:HOH:O	2.13	0.49
2:A:501:CIT:O5	4:A:1016:HOH:O	2.18	0.49
1:B:214:ASN:HB3	4:B:746:HOH:O	2.11	0.49
1:A:157[A]:GLN:NE2	4:A:1024:HOH:O	2.10	0.49
1:A:206[B]:GLN:HG3	1:A:210:GLN:O	2.13	0.49
1:B:211:LYS:HE2	1:B:213[B]:GLN:NE2	2.28	0.48
1:A:172:ARG:HB3	4:A:913:HOH:O	2.13	0.48
1:B:206[A]:GLN:HG3	1:B:210:GLN:O	2.13	0.48
1:B:63:ASN:H	2:B:501:CIT:H21	1.79	0.47
1:B:259[B]:GLN:NE2	4:B:874:HOH:O	2.24	0.47
1:A:214:ASN:HB3	4:A:912:HOH:O	2.13	0.47
1:B:278:GLU:CD	1:B:278:GLU:H	2.15	0.47
1:B:206[B]:GLN:HE21	1:B:212:THR:HG23	1.81	0.46
1:B:97:LYS:HB2	3:B:505:SO4:O1	2.15	0.46
3:A:503:SO4:O3	4:A:834:HOH:O	2.12	0.45
1:A:143:LYS:HG3	1:A:293[B]:PHE:CE2	2.52	0.45
1:A:230[A]:ASP:OD1	4:A:820:HOH:O	2.20	0.44
1:B:81:LYS:HD2	4:B:832:HOH:O	2.18	0.44
2:A:501:CIT:O3	2:A:501:CIT:O7	2.34	0.43
1:A:56:VAL:HG21	1:A:115:LEU:HD21	2.00	0.43
1:B:144:LEU:HD13	1:B:317:CYS:SG	2.58	0.43
1:B:318:LYS:HD2	1:B:318:LYS:HA	1.81	0.43
1:B:55:SER:O	1:B:58[B]:ASN:ND2	2.52	0.43
1:A:269[A]:LYS:NZ	1:A:295:GLU:OE1	2.49	0.42
1:A:125:LYS:NZ	3:A:502:SO4:O4	2.49	0.42
1:A:141:LEU:HA	1:A:141:LEU:HD23	1.87	0.42
1:A:116:ASN:HB3	3:A:510:SO4:O1	2.20	0.42
1:A:31:ASN:N	1:A:32:ASP:HA	2.35	0.42
1:A:31:ASN:OD1	1:A:319:VAL:HG12	2.19	0.41
1:A:36:LYS:H	1:A:130:ASN:HD21	1.68	0.41
1:A:238:ASP:HB3	1:A:243:ILE:O	2.21	0.41
1:B:62:GLY:HA3	2:B:501:CIT:H21	2.02	0.41
1:B:150:ASN:ND2	4:B:877:HOH:O	2.06	0.41
1:B:186:ILE:CG2	1:B:191:ALA:HB2	2.51	0.40
1:A:15:MET:HB3	1:A:15:MET:HE2	1.97	0.40
1:B:195:ALA:HB2	1:B:216[B]:MET:SD	2.62	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:LYS:N	4:B:750:HOH:O[1_455]	2.02	0.18
4:A:813:HOH:O	4:A:835:HOH:O[1_455]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/320 (107%)	337 (98%)	6 (2%)	0	100	100
1	B	333/320 (104%)	319 (96%)	13 (4%)	1 (0%)	41	27
All	All	676/640 (106%)	656 (97%)	19 (3%)	1 (0%)	47	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	77	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/292 (108%)	309 (98%)	7 (2%)	52	39
1	B	307/292 (105%)	300 (98%)	7 (2%)	50	37
All	All	623/584 (107%)	609 (98%)	14 (2%)	52	39

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	88	ASP
1	A	211	LYS
1	A	293[A]	PHE
1	A	293[B]	PHE
1	A	311	LYS
1	A	319	VAL
1	B	1	MET
1	B	135	LYS
1	B	237	ASN
1	B	241	ASN
1	B	280	ILE
1	B	281	LYS
1	B	308	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	510	-	4,4,4	0.14	0	6,6,6	0.10	0
3	SO4	B	509	-	4,4,4	0.16	0	6,6,6	0.06	0
3	SO4	A	508	-	4,4,4	0.15	0	6,6,6	0.10	0
3	SO4	A	504	-	4,4,4	0.12	0	6,6,6	0.23	0
2	CIT	A	501	-	3,12,12	1.12	0	3,17,17	1.76	1 (33%)
3	SO4	B	510	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	A	502	-	4,4,4	0.19	0	6,6,6	0.17	0
3	SO4	A	511	-	4,4,4	0.13	0	6,6,6	0.21	0
3	SO4	A	505	-	4,4,4	0.15	0	6,6,6	0.09	0
3	SO4	B	512	-	4,4,4	0.14	0	6,6,6	0.11	0
3	SO4	B	502	-	4,4,4	0.22	0	6,6,6	0.53	0
3	SO4	B	507	-	4,4,4	0.13	0	6,6,6	0.09	0
3	SO4	B	511	-	4,4,4	0.14	0	6,6,6	0.08	0
3	SO4	A	507	-	4,4,4	0.15	0	6,6,6	0.11	0
3	SO4	A	517	-	4,4,4	0.16	0	6,6,6	0.09	0
3	SO4	A	513	-	4,4,4	0.15	0	6,6,6	0.14	0
3	SO4	A	509	-	4,4,4	0.17	0	6,6,6	0.24	0
3	SO4	A	514	-	4,4,4	0.17	0	6,6,6	0.07	0
2	CIT	B	501	-	3,12,12	1.35	0	3,17,17	0.99	0
3	SO4	A	515	-	4,4,4	0.15	0	6,6,6	0.12	0
3	SO4	B	504	-	4,4,4	0.13	0	6,6,6	0.17	0
3	SO4	B	503	-	4,4,4	0.14	0	6,6,6	0.09	0
3	SO4	A	506	-	4,4,4	0.14	0	6,6,6	0.13	0
3	SO4	A	516	-	4,4,4	0.18	0	6,6,6	0.47	0
3	SO4	A	512	-	4,4,4	0.14	0	6,6,6	0.12	0
3	SO4	B	505	-	4,4,4	0.15	0	6,6,6	0.18	0
3	SO4	A	503	-	4,4,4	0.14	0	6,6,6	0.27	0
3	SO4	B	506	-	4,4,4	0.14	0	6,6,6	0.09	0
3	SO4	B	508	-	4,4,4	0.14	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	501	-	-	6/6/16/16	-
2	CIT	B	501	-	-	3/6/16/16	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	501	CIT	C3-C2-C1	-2.35	111.22	114.98

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	CIT	C1-C2-C3-O7
2	A	501	CIT	C1-C2-C3-C6
2	A	501	CIT	C2-C3-C4-C5
2	A	501	CIT	O7-C3-C4-C5
2	A	501	CIT	C6-C3-C4-C5
2	B	501	CIT	C2-C3-C4-C5
2	B	501	CIT	O7-C3-C4-C5
2	B	501	CIT	C6-C3-C4-C5
2	A	501	CIT	C1-C2-C3-C4

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	510	SO4	1	0
2	A	501	CIT	2	0
3	A	502	SO4	1	0
3	A	513	SO4	1	0
2	B	501	CIT	2	0
3	B	505	SO4	3	0
3	A	503	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/320 (99%)	-0.00	10 (3%) 49 43	10, 20, 46, 118	0
1	B	319/320 (99%)	0.37	25 (7%) 13 10	13, 30, 75, 129	0
All	All	637/640 (99%)	0.18	35 (5%) 25 20	10, 25, 65, 129	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	243	ILE	6.1
1	A	306	SER	5.8
1	B	308	SER	5.7
1	A	319	VAL	5.6
1	B	306	SER	5.5
1	B	244	ILE	4.8
1	B	242	LYS	4.6
1	B	245	ASP	4.3
1	B	309	ASP	4.0
1	B	278	GLU	4.0
1	B	310	SER	3.9
1	B	246	LYS	3.8
1	B	132	GLU	3.4
1	A	310	SER	3.4
1	B	279	ASN	3.2
1	B	311	LYS	3.1
1	A	305	ASN	3.1
1	B	133	ASN	3.0
1	B	240	LYS	2.9
1	A	304	ASP	2.8
1	A	311	LYS	2.7
1	B	130	ASN	2.7
1	B	162[A]	ASN	2.7
1	B	318	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	274	LYS	2.6
1	B	312	ILE	2.5
1	B	307	ASN	2.4
1	A	211	LYS	2.4
1	B	241	ASN	2.4
1	A	317	CYS	2.1
1	B	129	SER	2.1
1	B	247	ASN	2.1
1	A	278	GLU	2.0
1	A	133	ASN	2.0
1	B	131	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	513	5/5	0.51	0.33	97,101,102,102	5
2	CIT	B	501	13/13	0.57	0.30	72,88,98,103	0
3	SO4	B	510	5/5	0.65	0.22	124,128,129,131	0
3	SO4	A	512	5/5	0.73	0.36	83,84,87,91	5
3	SO4	B	511	5/5	0.75	0.21	116,119,122,123	0
3	SO4	A	505	5/5	0.75	0.18	99,102,108,109	0
2	CIT	A	501	13/13	0.76	0.31	77,102,115,116	0
3	SO4	B	507	5/5	0.79	0.30	109,112,114,116	0
3	SO4	A	508	5/5	0.81	0.23	81,84,88,94	0
3	SO4	B	508	5/5	0.81	0.24	109,114,116,117	0
3	SO4	A	506	5/5	0.82	0.30	115,115,116,119	0
3	SO4	B	504	5/5	0.83	0.20	71,82,90,96	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	507	5/5	0.85	0.17	94,94,98,100	0
3	SO4	A	515	5/5	0.85	0.19	71,72,79,80	5
3	SO4	A	517	5/5	0.87	0.21	68,73,76,77	5
3	SO4	A	511	5/5	0.87	0.16	80,81,84,93	0
3	SO4	B	506	5/5	0.87	0.38	99,102,107,108	0
3	SO4	A	509	5/5	0.87	0.21	39,57,66,70	5
3	SO4	B	512	5/5	0.88	0.21	90,93,97,100	0
3	SO4	A	510	5/5	0.88	0.22	122,123,125,127	0
3	SO4	B	503	5/5	0.90	0.13	81,84,88,90	0
3	SO4	B	505	5/5	0.91	0.22	45,58,74,78	0
3	SO4	A	514	5/5	0.92	0.41	27,45,51,52	5
3	SO4	B	509	5/5	0.92	0.37	115,115,118,120	0
3	SO4	A	504	5/5	0.93	0.18	51,77,84,85	0
3	SO4	A	516	5/5	0.94	0.20	22,26,36,41	5
3	SO4	B	502	5/5	0.95	0.17	30,38,50,54	0
3	SO4	A	503	5/5	0.98	0.16	36,40,53,64	0
3	SO4	A	502	5/5	0.98	0.09	21,27,36,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.