



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2021 – 04:04 PM EDT

PDB ID : 6OA8  
Title : Superfolder Green Fluorescent Protein with 4-cyano-L-phenylalanine at the chromophore (position 66)  
Authors : Piacentini, J.; Olenginski, G.M.; Brewer, S.H.; Phillips-Piro, C.M.  
Deposited on : 2019-03-15  
Resolution : 1.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2907 (1.40-1.36)
Clashscore	141614	3037 (1.40-1.36)
Ramachandran outliers	138981	2970 (1.40-1.36)
Sidechain outliers	138945	2969 (1.40-1.36)
RSRZ outliers	127900	2846 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4490 atoms, of which 2048 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	233	Total	C	H	N	O	S	0	44	0
			4091	1296	2030	357	401	7			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP A0A059PIQ0
A	1	VAL	-	expression tag	UNP A0A059PIQ0
A	2	SER	-	expression tag	UNP A0A059PIQ0
A	30	ARG	SER	conflict	UNP A0A059PIQ0
A	?	-	THR	deletion	UNP A0A059PIQ0
A	?	-	TYR	deletion	UNP A0A059PIQ0
A	66	M3V	GLY	conflict	UNP A0A059PIQ0
A	72	SER	ALA	conflict	UNP A0A059PIQ0
A	80	ARG	GLN	conflict	UNP A0A059PIQ0
A	206	VAL	ALA	conflict	UNP A0A059PIQ0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O S	0	0
			5	4 1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total Na 4 4	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	325	Total O 325 325	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.27Å 55.27Å 166.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.45 – 1.37 52.45 – 1.37	Depositor EDS
% Data completeness (in resolution range)	99.3 (52.45-1.37) 99.0 (52.45-1.37)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.39 (at 1.37Å)	Xtriage
Refinement program	PHENIX 1.19_4092	Depositor
R, $R_{free}$	0.161 , 0.189 0.161 , 0.189	Depositor DCC
$R_{free}$ test set	2744 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.4	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 51.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4490	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, EDO, SO4, M3V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/2240	0.72	0/3025

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	204[B]	GLN	Mainchain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2061	2030	1853	32	0
2	A	12	18	18	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	40	0	0	2	0
4	A	4	0	0	0	0
5	A	325	0	0	17	1
All	All	2442	2048	1871	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115[A]:GLU:OE2	5:A:401:HOH:O	1.52	1.26
1:A:115[B]:GLU:OE1	5:A:402:HOH:O	1.53	1.26
1:A:105[B]:THR:HG21	5:A:421:HOH:O	1.67	0.94
1:A:17:GLU:OE2	1:A:30[B]:ARG:NH1	2.09	0.84
1:A:153[B]:THR:HG23	2:A:301:EDO:H21	1.73	0.71
1:A:17:GLU:OE1	1:A:122[A]:ARG:NH1	2.23	0.71
1:A:159:ASN:ND2	5:A:404:HOH:O	2.22	0.66
3:A:310:SO4:O2	5:A:403:HOH:O	2.14	0.63
1:A:4:GLY:N	5:A:408:HOH:O	2.33	0.62
1:A:30[B]:ARG:HG3	1:A:49:THR:HG21	1.81	0.62
1:A:4:GLY:N	5:A:409:HOH:O	2.36	0.57
1:A:153[B]:THR:HG23	2:A:301:EDO:C2	2.35	0.56
1:A:184[B]:GLN:OE1	1:A:186[B]:THR:HG23	2.06	0.56
3:A:304:SO4:S	3:A:305:SO4:O4	2.65	0.55
1:A:17:GLU:OE1	1:A:122[A]:ARG:CZ	2.58	0.52
1:A:66:M3V:CN	1:A:203[A]:THR:OG1	2.59	0.51
1:A:105[B]:THR:HG23	5:A:543:HOH:O	2.10	0.51
1:A:38[B]:THR:HG21	5:A:501:HOH:O	2.13	0.49
1:A:105[B]:THR:CG2	5:A:421:HOH:O	2.42	0.47
1:A:30[B]:ARG:HG2	5:A:503:HOH:O	2.14	0.46
1:A:45:LYS:HE2	1:A:47:ILE:HD11	1.97	0.46
1:A:162:LYS:HE2	1:A:184[A]:GLN:OE1	2.17	0.45
1:A:166:LYS:HB3	1:A:178:LEU:HD11	1.98	0.45
1:A:107:LYS:HE2	5:A:696:HOH:O	2.16	0.45
1:A:72[B]:SER:HA	1:A:224[B]:VAL:HG23	1.99	0.44
1:A:99[B]:SER:HA	1:A:105[B]:THR:HG22	1.99	0.43
1:A:224[B]:VAL:HG21	5:A:626:HOH:O	2.18	0.43
1:A:203[A]:THR:HG22	1:A:224[A]:VAL:HG13	2.01	0.43
1:A:26[A]:LYS:NZ	5:A:422:HOH:O	2.53	0.42
1:A:36:ASP:OD1	1:A:38[B]:THR:OG1	2.34	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:ASP:OD2	1:A:184[A]:GLN:NE2	2.44	0.41
1:A:107:LYS:HE2	5:A:567:HOH:O	2.20	0.41
1:A:186[B]:THR:HG21	5:A:590:HOH:O	2.19	0.41
1:A:38[B]:THR:HG23	5:A:471:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:409:HOH:O	5:A:416:HOH:O[5_755]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/237 (116%)	273 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/207 (117%)	241 (100%)	1 (0%)	91	80

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	145	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	M3V	A	66	1	24,24,25	1.66	7 (29%)	31,33,35	2.06	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	M3V	A	66	1	-	0/14/33/34	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	M3V	CG2-CB2	3.49	1.53	1.46
1	A	66	M3V	CA2-C2	3.20	1.51	1.48
1	A	66	M3V	CZ-CN	2.88	1.51	1.44
1	A	66	M3V	C2-N3	2.86	1.46	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	M3V	C1-N2	2.30	1.35	1.32
1	A	66	M3V	C1-N3	2.28	1.41	1.37
1	A	66	M3V	O2-C2	-2.19	1.18	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	M3V	O-C-CA3	-5.42	110.02	126.39
1	A	66	M3V	CA2-N2-C1	4.81	109.31	105.77
1	A	66	M3V	CA-C1-N3	-4.78	119.01	124.75
1	A	66	M3V	CA-C1-N2	3.81	129.22	123.89
1	A	66	M3V	O2-C2-CA2	3.36	132.85	130.96
1	A	66	M3V	CG2-CB2-CA2	-2.31	127.12	129.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	M3V	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	302	-	3,3,3	0.43	0	2,2,2	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	309	-	4,4,4	0.15	0	6,6,6	0.19	0
3	SO4	A	310	-	4,4,4	0.15	0	6,6,6	0.09	0
3	SO4	A	304	4	4,4,4	0.22	0	6,6,6	0.22	0
3	SO4	A	305	4	4,4,4	0.24	0	6,6,6	0.43	0
3	SO4	A	308	-	4,4,4	0.18	0	6,6,6	0.17	0
3	SO4	A	306	-	4,4,4	0.23	0	6,6,6	0.17	0
3	SO4	A	307	-	4,4,4	0.15	0	6,6,6	0.10	0
2	EDO	A	303	-	3,3,3	0.46	0	2,2,2	0.37	0
2	EDO	A	301	-	3,3,3	0.49	0	2,2,2	0.37	0
3	SO4	A	311	-	4,4,4	0.18	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	302	-	-	0/1/1/1	-
2	EDO	A	303	-	-	1/1/1/1	-
2	EDO	A	301	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	EDO	O1-C1-C2-O2
2	A	303	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	310	SO4	1	0
3	A	304	SO4	1	0
3	A	305	SO4	1	0
2	A	301	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/237 (97%)	0.05	4 (1%) 70 72	9, 15, 29, 51	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	211	PRO	4.1
1	A	212	ASN	3.7
1	A	223[A]	PHE	2.7
1	A	214	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	M3V	A	66	23/24	0.93	0.09	12,17,34,34	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	A	303	4/4	0.73	0.32	44,53,55,66	0
4	NA	A	312	1/1	0.74	0.20	54,54,54,54	0
3	SO4	A	311	5/5	0.77	0.23	19,39,49,59	5
3	SO4	A	306	5/5	0.80	0.19	11,24,24,24	5
3	SO4	A	308	5/5	0.81	0.25	38,40,52,54	5
2	EDO	A	302	4/4	0.82	0.26	34,53,64,64	0
3	SO4	A	309	5/5	0.82	0.24	29,31,44,50	5
2	EDO	A	301	4/4	0.92	0.13	21,28,33,34	0
3	SO4	A	305	5/5	0.95	0.15	15,19,25,33	5
3	SO4	A	304	5/5	0.96	0.10	17,19,20,35	5
3	SO4	A	307	5/5	0.96	0.21	28,45,75,79	0
4	NA	A	315	1/1	0.96	0.12	32,32,32,32	1
3	SO4	A	310	5/5	0.97	0.26	39,40,48,57	5
4	NA	A	313	1/1	0.98	0.06	27,27,27,27	0
4	NA	A	314	1/1	0.99	0.08	17,17,17,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.