



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 11:49 AM BST

PDB ID : 5ODI
Title : Heterodisulfide reductase / [NiFe]-hydrogenase complex from Methanothermococcus thermolithotrophicus cocrystallized with CoM-SH
Authors : Wagner, T.; Koch, J.; Ermler, U.; Shima, S.
Deposited on : 2017-07-05
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	654	<div> <div>4%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
1	G	654	<div> <div>10%</div> <div> <div></div> <div>92%</div> <div>6%</div> </div> </div>
2	B	291	<div> <div>24%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
2	H	291	<div> <div>7%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
3	C	184	<div> <div>17%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
3	I	184	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	140	
4	J	140	
5	E	299	
5	K	299	
6	F	473	
6	L	473	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	GOL	H	304	-	-	-	X
12	COM	H	303	-	X	-	-
15	NFU	F	501	-	-	X	-
7	ACT	A	701	-	-	-	X
7	ACT	G	702	-	-	-	X

2 Entry composition [i](#)

There are 18 unique types of molecules in this entry. The entry contains 31986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heterodisulfide reductase, subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	651	Total	C	N	O	S	0	0	0
			4970	3137	844	940	49			
1	G	652	Total	C	N	O	S	0	0	0
			4979	3142	845	943	49			

- Molecule 2 is a protein called Heterodisulfide reductase, subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	291	Total	C	N	O	S	0	0	0
			2236	1420	379	413	24			
2	H	291	Total	C	N	O	S	0	0	0
			2236	1420	379	413	24			

- Molecule 3 is a protein called Heterodisulfide reductase, subunit C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	184	Total	C	N	O	S	0	0	0
			1426	890	247	275	14			
3	I	183	Total	C	N	O	S	0	0	0
			1416	885	246	271	14			

- Molecule 4 is a protein called Methyl-viologen reducing hydrogenase, subunit D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	137	Total	C	N	O	S	0	0	0
			1097	698	187	200	12			
4	J	138	Total	C	N	O	S	0	0	0
			1106	703	188	203	12			


- Molecule 5 is a protein called Methyl-viologen reducing hydrogenase, subunit G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	297	Total	C	N	O	S	0	0	0
			2249	1419	367	444	19			
5	K	297	Total	C	N	O	S	0	1	0
			2257	1424	370	444	19			

- Molecule 6 is a protein called Methyl-viologen reducing hydrogenase, subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	447	Total	C	N	O	S	0	0	0
			3521	2230	600	672	19			
6	L	447	Total	C	N	O	S	0	0	0
			3521	2230	600	672	19			

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	D	1	Total	C	O	0	0
			4	2	2		
7	G	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
8	G	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 9 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	Fe	S	0	0
			8	4	4		
9	A	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total 8	Fe 4	S 4	0	0
9	A	1	Total 8	Fe 4	S 4	0	0
9	A	1	Total 8	Fe 4	S 4	0	0
9	A	1	Total 8	Fe 4	S 4	0	0
9	C	1	Total 8	Fe 4	S 4	0	0
9	C	1	Total 8	Fe 4	S 4	0	0
9	E	1	Total 8	Fe 4	S 4	0	0
9	E	1	Total 8	Fe 4	S 4	0	0
9	E	1	Total 8	Fe 4	S 4	0	0
9	G	1	Total 8	Fe 4	S 4	0	0
9	G	1	Total 8	Fe 4	S 4	0	0
9	G	1	Total 8	Fe 4	S 4	0	0
9	G	1	Total 8	Fe 4	S 4	0	0
9	G	1	Total 8	Fe 4	S 4	0	0
9	G	1	Total 8	Fe 4	S 4	0	0
9	I	1	Total 8	Fe 4	S 4	0	0
9	I	1	Total 8	Fe 4	S 4	0	0
9	K	1	Total 8	Fe 4	S 4	0	0
9	K	1	Total 8	Fe 4	S 4	0	0
9	K	1	Total 8	Fe 4	S 4	0	0

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			6	3	3		
10	A	1	Total	C	O	0	0
			6	3	3		
10	G	1	Total	C	O	0	0
			6	3	3		
10	H	1	Total	C	O	0	0
			6	3	3		
10	K	1	Total	C	O	0	0
			6	3	3		

- Molecule 11 is Non-cubane [4Fe-4S]-cluster (three-letter code: 9S8) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	B	1	Total	Fe	S	0	0
			8	4	4		
11	B	1	Total	Fe	S	0	0
			8	4	4		
11	H	1	Total	Fe	S	0	0
			8	4	4		
11	H	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 12 is 1-THIOETHANESULFONIC ACID (three-letter code: COM) (formula: $C_2H_6O_3S_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
12	B	1	Total	C	O	S	0	0
			7	2	3	2		
12	H	1	Total	C	O	S	0	0
			7	2	3	2		

- Molecule 13 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	D	1	Total	Fe	S	0	0
			4	2	2		
13	J	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 14 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
14	D	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 15 is formyl[bis(hydrocyanato-1kappaC)]ironnickel(Fe-Ni) (three-letter code: NFU) (formula: C_3HFeN_2NiO).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
15	F	1	Total	C	Fe	N	Ni	O	0	0
			8	3	1	2	1	1		
15	L	1	Total	C	Fe	N	Ni	O	0	0
			8	3	1	2	1	1		

- Molecule 16 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	L	1	Total	Fe	0	0
			1	1		
16	F	1	Total	Fe	0	0
			1	1		

- Molecule 17 is 3,6,9,12,15,18,21,24,27,30,33,36,39-TRIDECAOXAHENTETRACONTANE-1,41-DIOL (three-letter code: PE3) (formula: $C_{28}H_{58}O_{15}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
17	G	1	Total	C	O	0	0
			14	9	5		

- Molecule 18 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	A	125	Total	O	0	0
			125	125		
18	B	14	Total	O	0	0
			14	14		
18	C	19	Total	O	0	0
			19	19		
18	D	36	Total	O	0	0
			36	36		
18	E	49	Total	O	0	0
			49	49		
18	F	65	Total	O	0	0
			65	65		
18	G	87	Total	O	0	0
			87	87		
18	H	23	Total	O	0	0
			23	23		
18	I	27	Total	O	0	0
			27	27		
18	J	5	Total	O	0	0
			5	5		
18	K	42	Total	O	0	0
			42	42		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	L	62	Total	O	0	0
			62	62		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Heterodisulfide reductase, subunit A




• Molecule 1: Heterodisulfide reductase, subunit A





• Molecule 2: Heterodisulfide reductase, subunit B









- Molecule 2: Heterodisulfide reductase, subunit B




- Molecule 3: Heterodisulfide reductase, subunit C





- Molecule 3: Heterodisulfide reductase, subunit C




- Molecule 4: Methyl-viologen reducing hydrogenase, subunit D




- Molecule 4: Methyl-viologen reducing hydrogenase, subunit D





- Molecule 5: Methyl-viologen reducing hydrogenase, subunit G





- Molecule 5: Methyl-viologen reducing hydrogenase, subunit G



- Molecule 6: Methyl-viologen reducing hydrogenase, subunit A



- Molecule 6: Methyl-viologen reducing hydrogenase, subunit A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	370.53 Å 97.08 Å 135.15 Å 90.00° 109.43° 90.00°	Depositor
Resolution (Å)	48.54 – 2.40 48.54 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.54-2.40) 99.9 (48.54-2.40)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.39 Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.202 , 0.241 0.230 , 0.273	Depositor DCC
R_{free} test set	8805 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 68.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	0.046 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	31986	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SF4, PE3, NFU, 9S8, FE, ACT, TRS, COM, FAD, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/5062	0.54	0/6841
1	G	0.28	0/5071	0.52	0/6853
2	B	0.20	0/2277	0.41	0/3070
2	H	0.22	0/2277	0.41	0/3070
3	C	0.27	0/1447	0.44	0/1946
3	I	0.27	0/1437	0.44	0/1934
4	D	0.36	0/1123	0.55	0/1508
4	J	0.26	0/1132	0.49	0/1520
5	E	0.31	0/2288	0.53	0/3102
5	K	0.25	0/2299	0.51	0/3116
6	F	0.27	0/3590	0.50	0/4853
6	L	0.25	0/3590	0.48	0/4853
All	All	0.28	0/31593	0.50	0/42666

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4970	0	4967	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	4979	0	4974	38	0
2	B	2236	0	2244	12	0
2	H	2236	0	2243	12	1
3	C	1426	0	1441	7	0
3	I	1416	0	1435	7	0
4	D	1097	0	1068	6	0
4	J	1106	0	1074	9	0
5	E	2249	0	2252	9	1
5	K	2257	0	2265	12	0
6	F	3521	0	3503	13	0
6	L	3521	0	3503	13	0
7	A	4	0	3	1	0
7	D	4	0	3	0	0
7	G	4	0	3	0	0
8	A	53	0	31	0	0
8	G	53	0	31	0	0
9	A	48	0	0	0	0
9	C	16	0	0	0	0
9	E	24	0	0	0	0
9	G	48	0	0	1	0
9	I	16	0	0	0	0
9	K	24	0	0	0	0
10	A	12	0	16	0	0
10	G	6	0	8	0	0
10	H	6	0	8	0	0
10	K	6	0	8	0	0
11	B	16	0	0	0	0
11	H	16	0	0	1	0
12	B	7	0	6	1	0
12	H	7	0	6	2	0
13	D	4	0	0	0	0
13	J	4	0	0	0	0
14	D	8	0	12	0	0
15	F	8	0	0	2	0
15	L	8	0	0	1	0
16	F	1	0	0	0	0
16	L	1	0	0	0	0
17	G	14	0	16	0	0
18	A	125	0	0	1	0
18	B	14	0	0	0	0
18	C	19	0	0	0	0
18	D	36	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
18	E	49	0	0	0	0
18	F	65	0	0	0	0
18	G	87	0	0	1	0
18	H	23	0	0	0	0
18	I	27	0	0	0	0
18	J	5	0	0	0	0
18	K	42	0	0	0	0
18	L	62	0	0	0	0
All	All	31986	0	31120	153	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:442:CYS:SG	6:F:445:CYS:HB2	2.00	1.02
11:H:302:9S8:S5	12:H:303:COM:S1	2.75	0.84
1:G:187:LYS:HD2	1:G:413:LEU:HD11	1.60	0.84
1:G:596:TYR:OH	4:J:74:TYR:HA	1.79	0.82
1:G:595:PRO:CG	4:J:74:TYR:O	2.27	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:201:ASN:N	2:H:248:LYS:O[4_556]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	649/654 (99%)	619 (95%)	29 (4%)	1 (0%)	47	62
1	G	650/654 (99%)	617 (95%)	33 (5%)	0	100	100
2	B	289/291 (99%)	275 (95%)	14 (5%)	0	100	100
2	H	289/291 (99%)	276 (96%)	13 (4%)	0	100	100
3	C	182/184 (99%)	178 (98%)	4 (2%)	0	100	100
3	I	181/184 (98%)	177 (98%)	4 (2%)	0	100	100
4	D	135/140 (96%)	130 (96%)	5 (4%)	0	100	100
4	J	136/140 (97%)	133 (98%)	3 (2%)	0	100	100
5	E	295/299 (99%)	279 (95%)	16 (5%)	0	100	100
5	K	296/299 (99%)	278 (94%)	18 (6%)	0	100	100
6	F	445/473 (94%)	421 (95%)	22 (5%)	2 (0%)	34	48
6	L	445/473 (94%)	421 (95%)	22 (5%)	2 (0%)	34	48
All	All	3992/4082 (98%)	3804 (95%)	183 (5%)	5 (0%)	51	68

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	F	398	ALA
6	L	398	ALA
6	L	291	THR
1	A	70	ALA
6	F	38	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	539/541 (100%)	531 (98%)	8 (2%)	65	80
1	G	540/541 (100%)	528 (98%)	12 (2%)	52	71
2	B	242/242 (100%)	239 (99%)	3 (1%)	71	85
2	H	242/242 (100%)	236 (98%)	6 (2%)	47	67

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	157/157 (100%)	152 (97%)	5 (3%)	39	59
3	I	156/157 (99%)	152 (97%)	4 (3%)	46	66
4	D	116/119 (98%)	116 (100%)	0	100	100
4	J	117/119 (98%)	117 (100%)	0	100	100
5	E	254/256 (99%)	248 (98%)	6 (2%)	49	68
5	K	255/256 (100%)	249 (98%)	6 (2%)	49	68
6	F	386/410 (94%)	378 (98%)	8 (2%)	53	72
6	L	386/410 (94%)	377 (98%)	9 (2%)	50	70
All	All	3390/3450 (98%)	3323 (98%)	67 (2%)	55	74

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	42	TYR
1	G	413	LEU
6	L	259	MET
1	G	96	PHE
1	G	359	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 46 ligands modelled in this entry, 2 are monoatomic - leaving 44 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	GOL	H	304	-	5,5,5	0.34	0	5,5,5	0.15	0
9	SF4	C	201	3	0,12,12	0.00	-	-		
9	SF4	A	703	1	0,12,12	0.00	-	-		
9	SF4	E	302	5	0,12,12	0.00	-	-		
7	ACT	D	202	-	1,3,3	4.67	1 (100%)	0,3,3	0.00	-
9	SF4	A	706	1	0,12,12	0.00	-	-		
9	SF4	I	201	3	0,12,12	0.00	-	-		
10	GOL	K	304	-	5,5,5	0.42	0	5,5,5	0.36	0
13	FES	D	201	4	0,4,4	0.00	-	-		
9	SF4	A	707	1	0,12,12	0.00	-	-		
9	SF4	K	303	5	0,12,12	0.00	-	-		
12	COM	B	303	-	6,6,6	1.40	2 (33%)	7,8,8	3.01	4 (57%)
9	SF4	G	708	1	0,12,12	0.00	-	-		
9	SF4	E	303	5	0,12,12	0.00	-	-		
9	SF4	I	202	3	0,12,12	0.00	-	-		
10	GOL	A	709	-	5,5,5	0.38	0	5,5,5	0.38	0
11	9S8	B	302	2	2,10,10	1.24	0	-		
11	9S8	B	301	2	2,10,10	0.88	0	-		
7	ACT	A	701	-	1,3,3	4.28	1 (100%)	0,3,3	0.00	-
9	SF4	G	704	1	0,12,12	0.00	-	-		
7	ACT	G	702	-	1,3,3	5.38	1 (100%)	0,3,3	0.00	-
12	COM	H	303	-	6,6,6	1.37	2 (33%)	7,8,8	3.48	4 (57%)
8	FAD	A	702	-	51,58,58	1.85	7 (13%)	60,89,89	2.09	12 (20%)
9	SF4	C	202	3	0,12,12	0.00	-	-		
9	SF4	G	707	1	0,12,12	0.00	-	-		
9	SF4	K	301	5	0,12,12	0.00	-	-		
9	SF4	A	708	1	0,12,12	0.00	-	-		
10	GOL	A	710	-	5,5,5	0.35	0	5,5,5	0.45	0
14	TRS	D	203	-	7,7,7	0.39	0	9,9,9	0.63	0
9	SF4	G	706	1	0,12,12	0.00	-	-		
9	SF4	E	301	5	0,12,12	0.00	-	-		
9	SF4	G	709	1	0,12,12	0.00	-	-		
9	SF4	A	705	1	0,12,12	0.00	-	-		
9	SF4	G	705	1	0,12,12	0.00	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	9S8	H	302	2	2,10,10	0.85	0	-		
10	GOL	G	710	-	5,5,5	0.32	0	5,5,5	0.42	0
13	FES	J	200	4	0,4,4	0.00	-	-		
15	NFU	L	501	6	2,7,7	0.95	0	-		
9	SF4	K	302	5	0,12,12	0.00	-	-		
8	FAD	G	703	-	51,58,58	1.90	7 (13%)	60,89,89	2.04	14 (23%)
15	NFU	F	501	6	2,7,7	1.48	0	-		
17	PE3	G	701	-	13,13,42	0.58	0	12,12,41	0.25	0
11	9S8	H	301	2	2,10,10	1.65	1 (50%)	-		
9	SF4	A	704	1	0,12,12	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	GOL	H	304	-	-	0/4/4/4	-
9	SF4	C	201	3	-	-	0/6/5/5
9	SF4	A	703	1	-	-	0/6/5/5
9	SF4	E	302	5	-	-	0/6/5/5
9	SF4	C	202	3	-	-	0/6/5/5
9	SF4	A	706	1	-	-	0/6/5/5
9	SF4	I	201	3	-	-	0/6/5/5
10	GOL	K	304	-	-	2/4/4/4	-
13	FES	D	201	4	-	-	0/1/1/1
9	SF4	A	707	1	-	-	0/6/5/5
9	SF4	K	303	5	-	-	0/6/5/5
12	COM	B	303	-	-	1/4/4/4	-
9	SF4	G	708	1	-	-	0/6/5/5
9	SF4	E	303	5	-	-	0/6/5/5
9	SF4	I	202	3	-	-	0/6/5/5
10	GOL	A	709	-	-	3/4/4/4	-
11	9S8	B	302	2	-	-	0/3/3/3
11	9S8	B	301	2	-	-	0/3/3/3
9	SF4	G	704	1	-	-	0/6/5/5
12	COM	H	303	-	-	3/4/4/4	-
8	FAD	A	702	-	-	3/30/50/50	0/6/6/6
9	SF4	G	707	1	-	-	0/6/5/5
9	SF4	K	301	5	-	-	0/6/5/5
9	SF4	A	708	1	-	-	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	GOL	A	710	-	-	2/4/4/4	-
14	TRS	D	203	-	-	3/9/9/9	-
9	SF4	G	706	1	-	-	0/6/5/5
9	SF4	E	301	5	-	-	0/6/5/5
9	SF4	G	709	1	-	-	0/6/5/5
9	SF4	A	705	1	-	-	0/6/5/5
9	SF4	G	705	1	-	-	0/6/5/5
11	9S8	H	302	2	-	-	0/3/3/3
9	SF4	K	302	5	-	-	0/6/5/5
13	FES	J	200	4	-	-	0/1/1/1
10	GOL	G	710	-	-	4/4/4/4	-
8	FAD	G	703	-	-	1/30/50/50	0/6/6/6
17	PE3	G	701	-	-	6/11/11/40	-
11	9S8	H	301	2	-	-	0/3/3/3
9	SF4	A	704	1	-	-	0/6/5/5

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	G	703	FAD	C4X-C10	9.88	1.48	1.38
8	A	702	FAD	C4X-C10	9.34	1.48	1.38
7	G	702	ACT	CH3-C	5.38	1.55	1.48
7	D	202	ACT	CH3-C	4.67	1.54	1.48
7	A	701	ACT	CH3-C	4.28	1.54	1.48

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	702	FAD	C4-N3-C2	8.08	121.97	115.14
8	G	703	FAD	C4-N3-C2	7.84	121.76	115.14
8	A	702	FAD	C1'-N10-C9A	6.78	123.63	118.29
8	G	703	FAD	C4-C4X-C10	-6.17	115.87	119.95
8	A	702	FAD	C4-C4X-C10	-5.49	116.32	119.95

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	K	304	GOL	C1-C2-C3-O3
10	A	709	GOL	C1-C2-C3-O3
12	H	303	COM	C1-C2-S2-O2S

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
14	D	203	TRS	C2-C-C1-O1
14	D	203	TRS	C3-C-C1-O1


There are no ring outliers.


7 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	B	303	COM	1	0
7	A	701	ACT	1	0
9	G	704	SF4	1	0
12	H	303	COM	2	0
11	H	302	9S8	1	0
15	L	501	NFU	1	0
15	F	501	NFU	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	651/654 (99%)	-0.17	24 (3%) 41 41	15, 43, 106, 161	0
1	G	652/654 (99%)	0.43	63 (9%) 7 7	23, 64, 144, 221	0
2	B	291/291 (100%)	1.24	69 (23%) 0 0	41, 93, 144, 165	0
2	H	291/291 (100%)	0.42	21 (7%) 15 14	37, 68, 105, 132	0
3	C	184/184 (100%)	0.62	32 (17%) 1 1	22, 60, 141, 156	0
3	I	183/184 (99%)	-0.13	2 (1%) 80 79	26, 51, 93, 123	0
4	D	137/140 (97%)	-0.38	2 (1%) 73 72	21, 37, 64, 118	0
4	J	138/140 (98%)	0.90	24 (17%) 1 1	47, 67, 97, 202	0
5	E	297/299 (99%)	-0.21	5 (1%) 70 68	19, 42, 76, 92	0
5	K	297/299 (99%)	0.43	20 (6%) 17 16	28, 62, 122, 165	0
6	F	447/473 (94%)	-0.02	11 (2%) 57 55	26, 52, 85, 102	0
6	L	447/473 (94%)	0.44	25 (5%) 24 23	28, 66, 139, 177	0
All	All	4015/4082 (98%)	0.27	298 (7%) 14 13	15, 58, 123, 221	0

The worst 5 of 298 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	135	ILE	11.3
2	B	136	ALA	10.0
3	C	178	GLU	7.0
1	G	601	LEU	6.9
3	C	181	ALA	6.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	ACT	G	702	4/4	0.66	0.42	108,110,110,110	0
10	GOL	H	304	6/6	0.69	0.44	95,96,99,101	0
7	ACT	A	701	4/4	0.72	0.55	62,66,66,67	0
17	PE3	G	701	14/43	0.74	0.31	50,58,74,76	0
7	ACT	D	202	4/4	0.83	0.34	78,79,80,82	0
10	GOL	K	304	6/6	0.84	0.20	54,55,61,62	0
14	TRS	D	203	8/8	0.86	0.25	41,54,56,61	0
11	9S8	B	302	8/8	0.89	0.14	68,80,86,90	0
10	GOL	A	710	6/6	0.89	0.21	42,49,49,52	0
10	GOL	G	710	6/6	0.91	0.18	46,51,55,59	0
10	GOL	A	709	6/6	0.92	0.15	44,49,50,53	0
12	COM	B	303	7/7	0.92	0.15	62,64,70,75	0
9	SF4	A	707	8/8	0.93	0.09	98,101,104,105	0
9	SF4	G	709	8/8	0.93	0.10	46,52,58,58	0
9	SF4	G	708	8/8	0.94	0.09	61,67,71,72	0
9	SF4	K	303	8/8	0.94	0.11	63,68,75,77	0
9	SF4	G	707	8/8	0.94	0.07	103,106,108,111	0
9	SF4	A	706	8/8	0.94	0.12	34,39,44,45	0
9	SF4	G	704	8/8	0.95	0.11	67,76,84,84	0
9	SF4	K	302	8/8	0.95	0.14	42,50,53,53	0
9	SF4	I	201	8/8	0.95	0.14	25,27,29,29	0
9	SF4	A	708	8/8	0.96	0.09	70,71,79,81	0
11	9S8	H	302	8/8	0.96	0.15	46,51,58,58	0
9	SF4	C	202	8/8	0.96	0.14	29,37,45,47	0
12	COM	H	303	7/7	0.96	0.18	42,46,57,61	0
9	SF4	E	301	8/8	0.96	0.16	18,26,30,32	0
9	SF4	G	706	8/8	0.97	0.07	81,82,85,86	0
9	SF4	I	202	8/8	0.97	0.15	36,37,49,53	0
9	SF4	A	703	8/8	0.97	0.11	13,18,20,24	0
9	SF4	A	705	8/8	0.97	0.12	24,32,36,41	0
9	SF4	G	705	8/8	0.97	0.11	15,17,20,26	0
9	SF4	K	301	8/8	0.97	0.20	37,41,49,50	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	SF4	E	302	8/8	0.97	0.10	13,24,27,29	0
11	9S8	B	301	8/8	0.97	0.11	57,67,69,74	0
8	FAD	G	703	53/53	0.97	0.15	8,32,52,54	0
8	FAD	A	702	53/53	0.97	0.13	11,27,49,53	0
11	9S8	H	301	8/8	0.97	0.11	41,49,55,59	0
9	SF4	A	704	8/8	0.97	0.09	12,21,28,31	0
9	SF4	C	201	8/8	0.98	0.14	24,27,33,40	0
9	SF4	E	303	8/8	0.98	0.09	10,16,24,27	0
15	NFU	F	501	8/8	0.98	0.17	29,36,39,41	0
13	FES	D	201	4/4	0.99	0.15	20,23,33,36	0
13	FES	J	200	4/4	0.99	0.12	52,54,59,61	0
15	NFU	L	501	8/8	0.99	0.14	28,34,37,41	0
16	FE	L	502	1/1	1.00	0.17	44,44,44,44	0
16	FE	F	502	1/1	1.00	0.10	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PE3 G 701:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD G 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9S8 H 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.