



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 05:57 pm BST

PDB ID : 1OE9
Title : Crystal structure of Myosin V motor with essential light chain-nucleotide-free
Authors : Coureux, P.-D.; Wells, A.L.; Menetrey, J.; Yengo, C.M.; Morris, C.A.; Sweeney, H.L.; Houdusse, A.
Deposited on : 2003-03-21
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	795	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>•</div> <div>8%</div> </div> </div>
2	B	151	<div> <div>5%</div> <div> <div></div> <div>77%</div> <div>15%</div> <div>8%</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYOSIN VA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	731	5866	3760	1004	1065	37	0	3	1

- Molecule 2 is a protein called MYOSIN LIGHT CHAIN 1, SLOW-TWITCH MUSCLE A ISOFORM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	139	1059	673	174	206	6	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	314	Total 314	O 314	0	0
4	B	19	Total 19	O 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MYOSIN VA



• Molecule 2: MYOSIN LIGHT CHAIN 1, SLOW-TWITCH MUSCLE A ISOFORM



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.93Å 98.25Å 111.38Å 90.00° 101.43° 90.00°	Depositor
Resolution (Å)	40.00 – 2.05 35.99 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.8 (40.00-2.05) 95.5 (35.99-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.220 , 0.264 0.223 , 0.263	Depositor DCC
R_{free} test set	3680 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7263	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	1/6005 (0.0%)	0.78	15/8108 (0.2%)
2	B	0.47	0/1075	0.68	0/1448
All	All	0.63	1/7080 (0.0%)	0.76	15/9556 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	239	MET	CG-SD	-5.78	1.66	1.81

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	674	ASP	CB-CG-OD2	6.61	124.25	118.30
1	A	301	ASP	CB-CG-OD2	6.59	124.23	118.30
1	A	17	ASP	CB-CG-OD2	5.91	123.62	118.30
1	A	784	ARG	NE-CZ-NH2	-5.75	117.42	120.30
1	A	355	ASP	CB-CG-OD2	5.74	123.46	118.30
1	A	417	ASP	CB-CG-OD2	5.66	123.39	118.30
1	A	344	ASP	CB-CG-OD2	5.64	123.38	118.30
1	A	66	ASP	CB-CG-OD2	5.48	123.23	118.30
1	A	229	ASP	CB-CG-OD2	5.43	123.19	118.30
1	A	719	ASP	CB-CG-OD2	5.42	123.17	118.30
1	A	367	ASP	CB-CG-OD2	5.13	122.92	118.30
1	A	555	ASP	CB-CG-OD2	5.06	122.86	118.30
1	A	666	ASP	CB-CG-OD2	5.06	122.86	118.30
1	A	215	ASP	CB-CG-OD2	5.03	122.83	118.30
1	A	35	ASP	CB-CG-OD2	5.03	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5866	0	5755	40	0
2	B	1059	0	998	13	0
3	A	5	0	0	0	0
4	A	314	0	0	9	0
4	B	19	0	0	2	0
All	All	7263	0	6753	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:ILE:HD13	1:A:592:LEU:HD11	1.66	0.75
1:A:138:HIS:HD2	1:A:140:PHE:H	1.32	0.73
1:A:660:ARG:NH1	4:A:2261:HOH:O	2.20	0.73
1:A:286:HIS:ND1	4:A:2128:HOH:O	2.28	0.66
2:B:113:LEU:HB3	2:B:120:MET:HE3	1.82	0.62
1:A:192:VAL:HG21	1:A:234:ILE:HG22	1.82	0.60
1:A:714:LEU:HD11	1:A:763:ARG:HA	1.83	0.59
1:A:713:VAL:O	1:A:766:LYS:HE2	2.05	0.57
1:A:31:TYR:CE1	1:A:35:ASP:HB2	2.42	0.55
1:A:169:LYS:NZ	4:A:2075:HOH:O	2.41	0.54
1:A:86:LEU:HD13	1:A:679:VAL:HG22	1.90	0.53
1:A:118:ILE:HA	1:A:123:ILE:HD13	1.90	0.52
1:A:138:HIS:CD2	1:A:140:PHE:H	2.21	0.51
1:A:675:GLU:O	1:A:679:VAL:HG23	2.10	0.51
2:B:37:ARG:NH1	4:B:2005:HOH:O	2.38	0.50
1:A:700:SER:HB2	1:A:751:PHE:HB2	1.93	0.50
1:A:685[A]:CYS:SG	4:A:2275:HOH:O	2.60	0.50
1:A:80:LEU:HG	1:A:691:ILE:HG23	1.96	0.48
2:B:72:PRO:HA	2:B:75:GLN:HE21	1.79	0.48
1:A:512:GLU:HG2	1:A:523:TRP:HB2	1.96	0.47
1:A:192:VAL:HG21	1:A:234:ILE:CG2	2.44	0.46
1:A:456:TYR:CD1	1:A:499:ILE:HG21	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:ILE:HD12	4:A:2045:HOH:O	2.16	0.46
1:A:132:MET:HB2	1:A:145:GLU:CG	2.46	0.45
1:A:179:TYR:CZ	1:A:183:VAL:HG11	2.52	0.45
1:A:210:LYS:HB2	1:A:296:ILE:HD11	1.99	0.45
2:B:113:LEU:HB3	2:B:120:MET:CE	2.46	0.44
2:B:90:LEU:HD12	2:B:145:LEU:HD23	1.98	0.44
1:A:471:LYS:NZ	4:A:2191:HOH:O	2.42	0.44
1:A:493:GLN:N	1:A:494:PRO:CD	2.79	0.44
2:B:71:LEU:HB3	2:B:72:PRO:HD3	2.00	0.44
1:A:685[B]:CYS:SG	1:A:687:VAL:HG23	2.58	0.44
1:A:787:TYR:HA	1:A:790:MET:HG2	2.00	0.43
1:A:460:LYS:HG3	1:A:644:LEU:HD21	1.99	0.43
1:A:783:MET:HE2	4:A:2311:HOH:O	2.18	0.43
1:A:257:ARG:HB3	1:A:263:TYR:CE2	2.53	0.43
1:A:130:GLN:NE2	4:A:2055:HOH:O	2.48	0.43
1:A:132:MET:HB2	1:A:145:GLU:HG2	2.01	0.42
2:B:132:HIS:NE2	2:B:147:HIS:CD2	2.87	0.42
1:A:360:PHE:O	1:A:364:MET:HG2	2.20	0.42
2:B:37:ARG:HA	2:B:41:GLN:O	2.20	0.42
1:A:335:LEU:O	1:A:400:ARG:HD2	2.20	0.42
1:A:553:PHE:CE2	1:A:689:GLU:HB3	2.54	0.42
2:B:29:TYR:CD2	2:B:52:LEU:HD22	2.55	0.42
2:B:36:MET:SD	2:B:77:VAL:HG21	2.60	0.42
1:A:781:TRP:CD1	2:B:148:ILE:HA	2.55	0.42
2:B:13:LYS:O	2:B:17:GLU:HG2	2.20	0.41
1:A:497:ASN:ND2	4:A:2200:HOH:O	2.52	0.41
1:A:496:ILE:O	1:A:500:GLU:HG2	2.20	0.41
1:A:362:ASP:OD1	1:A:585:LYS:NZ	2.54	0.41
2:B:37:ARG:HD2	4:B:2005:HOH:O	2.21	0.41
1:A:224:ILE:HD13	1:A:239:MET:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	720/795 (91%)	700 (97%)	17 (2%)	3 (0%)	34	24
2	B	133/151 (88%)	131 (98%)	2 (2%)	0	100	100
All	All	853/946 (90%)	831 (97%)	19 (2%)	3 (0%)	34	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	633	LYS
1	A	671	PHE
1	A	351	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	622/696 (89%)	610 (98%)	12 (2%)	57	53
2	B	108/129 (84%)	104 (96%)	4 (4%)	34	27
All	All	730/825 (88%)	714 (98%)	16 (2%)	52	46

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	LEU
1	A	233	ARG
1	A	277	LEU
1	A	307	ASN
1	A	315	LEU
1	A	439	TYR
1	A	488	ASP
1	A	514	LYS
1	A	534	LYS
1	A	586	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	691	ILE
1	A	719	ASP
2	B	39	LEU
2	B	62	SER
2	B	105	MET
2	B	149	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	HIS
1	A	264	GLN
1	A	406	HIS
1	A	410	ASN
1	A	424	HIS
1	A	430	HIS
1	A	463	GLN
1	A	493	GLN
1	A	497	ASN
2	B	75	GLN
2	B	147	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1796	-	4,4,4	0.16	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	731/795 (91%)	0.26	33 (4%) 33 35	21, 39, 59, 73	0
2	B	139/151 (92%)	0.44	8 (5%) 23 25	40, 54, 67, 70	0
All	All	870/946 (91%)	0.29	41 (4%) 31 33	21, 42, 63, 73	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	387	TYR	4.6
2	B	135	SER	4.3
1	A	380	LEU	4.3
1	A	632	HIS	4.2
1	A	381	ALA	4.0
2	B	88	ASP	3.7
1	A	332	ILE	3.6
1	A	386	THR	3.5
1	A	426	THR	3.5
1	A	722	SER	3.3
1	A	10	TYR	3.2
2	B	87	GLU	3.2
1	A	249	VAL	3.1
2	B	131	GLY	3.0
2	B	136	ASN	3.0
1	A	478	MET	2.9
1	A	262	PHE	2.7
2	B	86	TYR	2.6
1	A	206	ILE	2.6
1	A	333	LEU	2.5
1	A	259	TYR	2.5
1	A	329	LEU	2.5
1	A	330	ALA	2.5
1	A	352	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	794	ASP	2.4
1	A	335	LEU	2.3
1	A	132	MET	2.3
2	B	134	ASP	2.3
1	A	543	LEU	2.3
1	A	154	GLU	2.3
1	A	408	TYR	2.2
1	A	34	GLY	2.2
1	A	57	LYS	2.2
1	A	454	ILE	2.2
1	A	716	LYS	2.1
1	A	244	LEU	2.1
1	A	411	LEU	2.1
1	A	412	PHE	2.1
1	A	791	ARG	2.0
1	A	243	LEU	2.0
2	B	89	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	1796	5/5	0.86	0.23	44,46,49,50	5

6.5 Other polymers ⓘ

There are no such residues in this entry.