



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2020 – 08:14 pm BST

PDB ID : 4OIR
Title : Crystal structure of Thermus thermophilus RNA polymerase transcription initiation complex soaked with GE23077 and rifamycin SV
Authors : Zhang, Y.; Ebright, R.H.; Arnold, E.
Deposited on : 2014-01-20
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	<div> <div>59%</div> <div>16%</div> <div>24%</div> </div>
1	B	305	<div> <div>56%</div> <div>18%</div> <div>26%</div> </div>
2	C	1119	<div> <div>81%</div> <div>17%</div> <div>2%</div> </div>
3	D	1524	<div> <div>79%</div> <div>18%</div> <div>3%</div> </div>
4	E	99	<div> <div>82%</div> <div>12%</div> <div>5%</div> </div>
5	F	443	<div> <div>64%</div> <div>13%</div> <div>22%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	G	21	<div><div></div><div>43%</div><div>33%</div><div>24%</div></div>
7	H	27	<div><div></div><div>22%</div><div>59%</div><div>7%</div><div>11%</div></div>
8	I	7	<div><div></div><div>57%</div><div>43%</div></div>

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 28730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1809	1155	315	337	2			
1	B	227	Total	C	N	O	S	0	0	0
			1789	1143	310	334	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1112	Total	C	N	O	S	0	0	0
			8774	5550	1565	1635	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1485	Total	C	N	O	S	0	1	0
			11739	7441	2069	2193	36			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	94	Total	C	N	O	S	0	0	0
			758	483	132	139	4			

- Molecule 5 is a protein called DNA directed RNA polymerase sigma factor A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	EXPRESSION TAG	UNP Q5SKW1
F	-18	GLY	-	EXPRESSION TAG	UNP Q5SKW1
F	-17	SER	-	EXPRESSION TAG	UNP Q5SKW1
F	-16	SER	-	EXPRESSION TAG	UNP Q5SKW1
F	-15	HIS	-	EXPRESSION TAG	UNP Q5SKW1
F	-14	HIS	-	EXPRESSION TAG	UNP Q5SKW1
F	-13	HIS	-	EXPRESSION TAG	UNP Q5SKW1
F	-12	HIS	-	EXPRESSION TAG	UNP Q5SKW1
F	-11	HIS	-	EXPRESSION TAG	UNP Q5SKW1
F	-10	HIS	-	EXPRESSION TAG	UNP Q5SKW1
F	-9	SER	-	EXPRESSION TAG	UNP Q5SKW1
F	-8	SER	-	EXPRESSION TAG	UNP Q5SKW1
F	-7	GLY	-	EXPRESSION TAG	UNP Q5SKW1
F	-6	LEU	-	EXPRESSION TAG	UNP Q5SKW1
F	-5	VAL	-	EXPRESSION TAG	UNP Q5SKW1
F	-4	PRO	-	EXPRESSION TAG	UNP Q5SKW1
F	-3	ARG	-	EXPRESSION TAG	UNP Q5SKW1
F	-2	GLY	-	EXPRESSION TAG	UNP Q5SKW1
F	-1	SER	-	EXPRESSION TAG	UNP Q5SKW1
F	0	HIS	-	EXPRESSION TAG	UNP Q5SKW1

- Molecule 6 is a DNA chain called 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*GP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	16	Total	C	N	O	P	0	0	0
			328	156	63	94	15			

- Molecule 7 is a DNA chain called 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*C P*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			


- Molecule 8 is a protein called GE23077.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	I	7	Total	C	N	O	0	0	0
			50	26	9	15			

- Molecule 9 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total Mg 1 1	0	0
9	D	2	Total Mg 2 2	0	0
9	F	1	Total Mg 1 1	0	0

- Molecule 10 is rifamycin SV (three-letter code: RFV) (formula: $C_{37}H_{49}NO_{12}$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	C	1	Total C N O 50 37 1 12	0	0

- Molecule 11 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	D	2	Total Zn 2 2	0	0


- Molecule 12 is (2Z)-2-methylbut-2-enoic acid (three-letter code: MB8) (formula: $C_5H_8O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	I	1	Total	C	O	0	0
			2	1	1		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	8	Total	O	0	0
			8	8		
13	B	6	Total	O	0	0
			6	6		
13	C	46	Total	O	0	0
			46	46		
13	D	46	Total	O	0	0
			46	46		
13	E	3	Total	O	0	0
			3	3		
13	F	9	Total	O	0	0
			9	9		
13	G	4	Total	O	0	0
			4	4		
13	H	1	Total	O	0	0
			1	1		



• Molecule 3: DNA-directed RNA polymerase subunit beta'







- Molecule 4: DNA-directed RNA polymerase subunit omega

Chain E: 82% 12% 5%



- Molecule 5: DNA directed RNA polymerase sigma factor A

Chain F: 2% 64% 13% 22%




- Molecule 6: 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*GP*GP*G)-3'

Chain G: 43% 33% 24%



- Molecule 7: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'

Chain H: 22% 59% 7% 11%



- Molecule 8: GE23077

Chain I: 57% 43%

71	74	T5	76	67
----	----	----	----	----

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.70Å 105.01Å 295.36Å 90.00° 98.79° 90.00°	Depositor
Resolution (Å)	47.50 – 3.10 47.50 – 3.11	Depositor EDS
% Data completeness (in resolution range)	82.6 (47.50-3.10) 82.7 (47.50-3.11)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 3.12Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.206 , 0.253 0.208 , 0.255	Depositor DCC
R_{free} test set	4160 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	63.5	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 25.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.018 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.015 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	28730	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FGL, ZN, RFV, 2TL, DVA, MG, 2RA, DSN, MB8, 0QZ, R2T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1841	0.46	0/2504
1	B	0.23	0/1821	0.46	0/2476
2	C	0.24	0/8941	0.45	0/12092
3	D	0.24	0/11948	0.45	0/16152
4	E	0.24	0/772	0.41	0/1040
5	F	0.23	0/2852	0.40	0/3837
6	G	0.48	0/368	1.06	2/567 (0.4%)
7	H	0.50	0/556	1.14	3/858 (0.3%)
All	All	0.25	0/29099	0.49	5/39526 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
8	I	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	5	DC	O4'-C4'-C3'	-9.49	100.30	106.00
7	H	17	DA	O4'-C1'-N9	7.70	113.39	108.00
7	H	23	DG	C4'-C3'-C2'	-5.43	98.22	103.10
7	H	16	DC	O4'-C1'-N1	5.04	111.53	108.00
6	G	5	DC	C4'-C3'-C2'	-5.01	98.59	103.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
8	I	5	2TL	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1863	32	0
1	B	1789	0	1841	33	0
2	C	8774	0	8877	129	0
3	D	11739	0	11975	173	0
4	E	758	0	770	10	0
5	F	2807	0	2882	46	0
6	G	328	0	181	5	0
7	H	495	0	272	12	0
8	I	50	0	37	3	0
9	B	1	0	0	0	0
9	D	2	0	0	0	0
9	F	1	0	0	0	0
10	C	50	0	48	2	0
11	D	2	0	0	0	0
12	I	2	0	0	0	0
13	A	8	0	0	0	0
13	B	6	0	0	0	0
13	C	46	0	0	1	0
13	D	46	0	0	4	0
13	E	3	0	0	0	0
13	F	9	0	0	2	0
13	G	4	0	0	0	0
13	H	1	0	0	0	0
All	All	28730	0	28746	392	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 392 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:321:ILE:HD11	5:F:329:TYR:HA	1.69	0.74
2:C:12:VAL:HG21	2:C:472:ARG:HD3	1.69	0.74
3:D:256:GLU:HG3	3:D:300:LYS:HG3	1.69	0.74
10:C:1201:RFV:H51	10:C:1201:RFV:H48	1.33	0.72
2:C:428:ARG:NH2	2:C:447:ALA:O	2.22	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/305 (75%)	226 (99%)	3 (1%)	0	100	100
1	B	225/305 (74%)	223 (99%)	2 (1%)	0	100	100
2	C	1108/1119 (99%)	1080 (98%)	28 (2%)	0	100	100
3	D	1482/1524 (97%)	1450 (98%)	31 (2%)	1 (0%)	51	83
4	E	92/99 (93%)	90 (98%)	2 (2%)	0	100	100
5	F	344/443 (78%)	339 (98%)	5 (2%)	0	100	100
All	All	3480/3795 (92%)	3408 (98%)	71 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	565	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/264 (76%)	196 (98%)	4 (2%)	55	80
1	B	200/264 (76%)	190 (95%)	10 (5%)	24	57
2	C	936/941 (100%)	906 (97%)	30 (3%)	39	69
3	D	1253/1279 (98%)	1219 (97%)	34 (3%)	44	74
4	E	82/88 (93%)	81 (99%)	1 (1%)	71	88
5	F	301/388 (78%)	290 (96%)	11 (4%)	34	66
All	All	2972/3224 (92%)	2882 (97%)	90 (3%)	41	71

5 of 90 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	1010	THR
3	D	273	ARG
5	F	295	MET
2	C	1057	SER
3	D	80	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	696	HIS
3	D	1195	GLN
3	D	724	GLN
3	D	350	HIS
3	D	714	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	FGL	I	7	8	2,6,7	0.63	0	0,7,9	0.00	-
8	0QZ	I	6	8	4,5,6	1.60	1 (25%)	2,5,7	0.90	0
8	2TL	I	5	8	5,6,7	1.08	0	6,7,9	0.91	0
8	2RA	I	1	8,12	3,5,6	0.58	0	1,5,7	0.45	0
8	R2T	I	4	8	8,10,11	1.92	2 (25%)	6,13,15	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	FGL	I	7	8	-	0/0/6/8	-
8	0QZ	I	6	8	-	1/3/4/6	-
8	2TL	I	5	8	-	1/5/6/8	-
8	2RA	I	1	8,12	-	2/2/4/6	-
8	R2T	I	4	8	-	6/13/14/16	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	I	4	R2T	CD-NE2	4.29	1.43	1.32
8	I	6	0QZ	OB-CA	-3.04	1.38	1.43
8	I	4	R2T	OB1-CB	-2.26	1.37	1.43

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	I	5	2TL	O-C-CA-CB
8	I	1	2RA	C-CA-CB-NG
8	I	1	2RA	N-CA-CB-NG
8	I	6	0QZ	N-C1-CA-C
8	I	4	R2T	OE1-CD-CG-OG1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	I	7	FGL	1	0
8	I	4	R2T	2	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	RFV	C	1201	-	52,53,53	3.52	12 (23%)	72,80,80	2.44	16 (22%)
12	MB8	I	101	8	0,1,6	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	RFV	C	1201	-	-	10/55/70/70	0/3/4/4

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	1201	RFV	C12-C11	-20.64	1.40	1.55
10	C	1201	RFV	C17-C16	6.85	1.54	1.34
10	C	1201	RFV	C15-N1	5.38	1.47	1.35
10	C	1201	RFV	O7-C25	-5.03	1.37	1.44
10	C	1201	RFV	C5-C11	-3.78	1.47	1.50

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	C	1201	RFV	C12-C11-C5	12.67	111.13	101.52
10	C	1201	RFV	C30-C16-C17	-6.44	107.83	123.42
10	C	1201	RFV	O3-C6-C7	5.47	130.54	121.14
10	C	1201	RFV	O4-C11-C12	5.35	122.85	113.41
10	C	1201	RFV	O7-C35-C36	4.23	118.88	111.09

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	C	1201	RFV	C16-C17-C18-C19
10	C	1201	RFV	C26-C27-C28-C29
10	C	1201	RFV	C26-C27-O6-C37
10	C	1201	RFV	C28-C27-O6-C37
10	C	1201	RFV	C18-C19-C20-C31

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	C	1201	RFV	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/305 (75%)	-0.43	4 (1%) 70 49	31, 50, 79, 128	0
1	B	227/305 (74%)	-0.36	2 (0%) 84 69	35, 65, 92, 125	0
2	C	1112/1119 (99%)	-0.35	10 (0%) 84 69	15, 44, 109, 139	0
3	D	1485/1524 (97%)	-0.28	15 (1%) 82 67	15, 50, 111, 156	0
4	E	94/99 (94%)	-0.40	0 100 100	27, 55, 99, 107	0
5	F	346/443 (78%)	-0.17	7 (2%) 65 44	24, 66, 122, 138	0
6	G	16/21 (76%)	-0.32	0 100 100	66, 101, 179, 183	0
7	H	24/27 (88%)	-0.24	0 100 100	58, 110, 171, 185	0
8	I	0/7	-	-	-	-
All	All	3535/3850 (91%)	-0.31	38 (1%) 80 64	15, 52, 112, 185	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	232	ALA	4.0
2	C	766	GLU	3.2
3	D	1281	VAL	2.9
1	A	233	VAL	2.9
5	F	149	GLU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	FGL	I	7	7/8	0.93	0.15	19,25,27,29	0
8	DSN	I	2	6/7	0.95	0.17	20,21,29,31	0
8	R2T	I	4	11/12	0.95	0.16	21,25,32,33	0
8	2RA	I	1	6/7	0.96	0.11	24,25,29,34	0
8	0QZ	I	6	6/7	0.97	0.15	23,25,25,28	0
8	DVA	I	3	7/8	0.97	0.17	13,17,21,22	0
8	2TL	I	5	7/8	0.98	0.18	21,21,24,29	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	MG	D	2004	1/1	0.77	0.18	50,50,50,50	0
10	RFV	C	1201	50/50	0.95	0.24	19,30,47,53	0
9	MG	F	2001	1/1	0.96	0.17	55,55,55,55	0
9	MG	B	401	1/1	0.96	0.25	25,25,25,25	0
12	MB8	I	101	2/7	0.96	0.13	26,26,26,31	0
11	ZN	D	2001	1/1	0.97	0.17	43,43,43,43	0
9	MG	D	2003	1/1	0.97	0.17	14,14,14,14	0
11	ZN	D	2002	1/1	0.97	0.08	79,79,79,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around RFV C 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.