



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2020 – 11:23 AM EDT

PDB ID : 6OJE  
Title : Dimeric structure of LRRK2 GTPase domain  
Authors : Hoang, Q.Q.; Wu, C.X.; Liao, J.; Park, Y.  
Deposited on : 2019-04-11  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.6  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	<div> <div>11%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>10%</div> </div> </div>
1	B	206	<div> <div>12%</div> <div> <div></div> <div>78%</div> <div>11%</div> <div>9%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leucine-rich repeat serine/threonine-protein kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	186	Total	C	N	O	S	8	7	0
			1503	950	269	277	7			
1	B	187	Total	C	N	O	S	0	4	0
			1484	939	266	270	9			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1315	MET	-	initiating methionine	UNP Q5S007
A	1316	GLY	-	expression tag	UNP Q5S007
A	1317	SER	-	expression tag	UNP Q5S007
A	1318	SER	-	expression tag	UNP Q5S007
A	1319	HIS	-	expression tag	UNP Q5S007
A	1320	HIS	-	expression tag	UNP Q5S007
A	1321	HIS	-	expression tag	UNP Q5S007
A	1322	HIS	-	expression tag	UNP Q5S007
A	1323	HIS	-	expression tag	UNP Q5S007
A	1324	HIS	-	expression tag	UNP Q5S007
A	1325	SER	-	expression tag	UNP Q5S007
A	1326	GLN	-	expression tag	UNP Q5S007
A	1327	ASP	-	expression tag	UNP Q5S007
A	1328	PRO	-	expression tag	UNP Q5S007
B	1315	MET	-	initiating methionine	UNP Q5S007
B	1316	GLY	-	expression tag	UNP Q5S007
B	1317	SER	-	expression tag	UNP Q5S007
B	1318	SER	-	expression tag	UNP Q5S007
B	1319	HIS	-	expression tag	UNP Q5S007
B	1320	HIS	-	expression tag	UNP Q5S007
B	1321	HIS	-	expression tag	UNP Q5S007
B	1322	HIS	-	expression tag	UNP Q5S007
B	1323	HIS	-	expression tag	UNP Q5S007
B	1324	HIS	-	expression tag	UNP Q5S007
B	1325	SER	-	expression tag	UNP Q5S007

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1326	GLN	-	expression tag	UNP Q5S007
B	1327	ASP	-	expression tag	UNP Q5S007
B	1328	PRO	-	expression tag	UNP Q5S007

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 28	C 10	N 5	O 11	P 2	0	0
3	B	1	Total 28	C 10	N 5	O 11	P 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	116	Total	O	0	7
			123	123		

Continued on next page...


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	124	Total 126	O 126	0	2

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Leucine-rich repeat serine/threonine-protein kinase 2

Chain A: 



- Molecule 1: Leucine-rich repeat serine/threonine-protein kinase 2

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.65Å 103.69Å 44.59Å 90.00° 101.20° 90.00°	Depositor
Resolution (Å)	43.80 – 1.95 43.80 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.1 (43.80-1.95) 99.0 (43.80-1.95)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	205.22 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, $R_{free}$	0.160 , 0.196 0.169 , 0.199	Depositor DCC
$R_{free}$ test set	1458 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.9	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.089 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3296	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/1534	0.72	0/2068
1	B	0.64	0/1509	0.72	0/2034
All	All	0.65	0/3043	0.72	0/4102

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1503	0	1503	9	0
1	B	1484	0	1472	19	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	28	0	12	0	0
3	B	28	0	12	0	0
4	A	123	0	0	0	0
4	B	126	0	0	4	0
All	All	3296	0	2999	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1352:GLN:O	1:A:1356:LYS:HG3	1.70	0.90
1:A:1358:LYS:HD3	1:A:1361:ASP:HA	1.67	0.75
1:A:1368:THR:HG21	4:B:1762:HOH:O	1.98	0.62
1:A:1503:THR:O	1:A:1507:GLU:HB2	2.05	0.57
1:A:1514:ARG:O	1:A:1517:LEU:HB2	2.04	0.56
1:B:1442:ALA:O	1:B:1445:SER:OG	2.29	0.51
1:B:1493:GLU:OE1	1:B:1501:ARG:NH2	2.43	0.51
1:B:1483:ARG:HG3	1:B:1507:GLU:CD	2.31	0.50
1:B:1357:THR:HG22	1:B:1358:LYS:HD2	1.94	0.50
1:B:1364:MET:SD	1:B:1366:SER:O	2.72	0.48
1:B:1398[A]:ARG:NH2	4:B:1704:HOH:O	2.45	0.47
1:A:1509:LEU:HD23	1:B:1388:LEU:HD13	1.97	0.46
1:A:1505:ILE:HG12	1:B:1378:ILE:HG21	1.98	0.46
1:A:1340:VAL:HB	1:A:1434:TRP:CZ3	2.50	0.46
1:B:1383:LYS:CD	1:B:1383:LYS:H	2.29	0.46
1:B:1357:THR:CG2	1:B:1358:LYS:HD2	2.47	0.44
1:B:1398[B]:ARG:NH1	4:B:1706:HOH:O	2.50	0.44
1:B:1483:ARG:HG3	1:B:1507:GLU:OE2	2.17	0.44
1:B:1357:THR:HG22	1:B:1358:LYS:N	2.33	0.44
1:B:1349:THR:O	1:B:1352:GLN:HG2	2.17	0.43
1:B:1423:LYS:NZ	4:B:1710:HOH:O	2.52	0.42
1:B:1410:THR:O	1:B:1442:ALA:HB2	2.20	0.42
1:B:1513:ILE:O	1:B:1516:GLN:HB3	2.20	0.41
1:A:1505:ILE:HG12	1:B:1378:ILE:HD13	2.02	0.41
1:B:1414:LEU:HD13	1:B:1446:PRO:HG2	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/206 (93%)	180 (94%)	9 (5%)	2 (1%)	15	6
1	B	189/206 (92%)	184 (97%)	4 (2%)	1 (0%)	29	17
All	All	380/412 (92%)	364 (96%)	13 (3%)	3 (1%)	19	9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1358	LYS
1	B	1384	ARG
1	A	1412	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/181 (88%)	158 (99%)	2 (1%)	69	65
1	B	154/181 (85%)	148 (96%)	6 (4%)	32	19
All	All	314/362 (87%)	306 (98%)	8 (2%)	46	38

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1368	THR
1	A	1495	ASP
1	B	1381	ARG
1	B	1383	LYS
1	B	1388	LEU
1	B	1445	SER
1	B	1449	LEU
1	B	1499	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GDP	A	1602	2	24,30,30	1.19	2 (8%)	31,47,47	2.04	6 (19%)
3	GDP	B	1602	2	24,30,30	1.12	2 (8%)	31,47,47	1.93	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	1602	2	-	0/12/32/32	0/3/3/3
3	GDP	B	1602	2	-	1/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1602	GDP	C6-C5	4.04	1.48	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1602	GDP	C6-C5	4.02	1.48	1.41
3	B	1602	GDP	C5-C4	2.28	1.47	1.40
3	A	1602	GDP	C5-C4	2.25	1.46	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1602	GDP	C2-N3-C4	5.91	122.11	115.36
3	A	1602	GDP	C6-C5-C4	-5.06	115.97	120.80
3	B	1602	GDP	C6-C5-C4	-4.85	116.17	120.80
3	B	1602	GDP	C2-N3-C4	4.50	120.50	115.36
3	A	1602	GDP	N3-C2-N1	-4.22	121.59	127.22
3	B	1602	GDP	C6-N1-C2	4.22	122.64	115.93
3	A	1602	GDP	C6-N1-C2	3.90	122.12	115.93
3	B	1602	GDP	N3-C2-N1	-3.85	122.09	127.22
3	B	1602	GDP	C5-C6-N1	-3.78	118.27	123.43
3	A	1602	GDP	C5-C6-N1	-3.15	119.12	123.43
3	A	1602	GDP	C4-C5-N7	-2.49	106.80	109.40
3	B	1602	GDP	C4-C5-N7	-2.13	107.17	109.40
3	B	1602	GDP	C1'-N9-C4	-2.05	123.05	126.64

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1602	GDP	PA-O3A-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	186/206 (90%)	0.54	23 (12%) 4 6	27, 45, 101, 123	0
1	B	187/206 (90%)	0.62	24 (12%) 3 6	27, 43, 110, 137	0
All	All	373/412 (90%)	0.58	47 (12%) 3 6	27, 44, 110, 137	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1360	SER	10.7
1	B	1362	LEU	10.5
1	A	1362	LEU	9.6
1	B	1519	VAL	9.4
1	B	1518	VAL	7.3
1	B	1380	ILE	7.1
1	B	1360	SER	6.9
1	B	1385	LYS	6.4
1	A	1382	ASP	6.2
1	B	1383	LYS	5.8
1	A	1358	LYS	5.7
1	B	1361	ASP	5.7
1	A	1380	ILE	5.5
1	B	1364	MET	5.5
1	B	1357	THR	4.9
1	B	1381	ARG	4.9
1	B	1382	ASP	4.7
1	A	1384	ARG	4.7
1	B	1358	LYS	4.6
1	A	1364	MET	4.6
1	B	1359	LYS	4.4
1	B	1387	ASP	4.2
1	A	1359	LYS	4.0
1	B	1515	ASP	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1517	LEU	3.8
1	A	1514	ARG	3.7
1	A	1515	ASP	3.7
1	A	1517	LEU	3.7
1	A	1357	THR	3.6
1	A	1381	ARG	3.6
1	B	1384	ARG	3.6
1	A	1518	VAL	3.3
1	A	1379	GLN	3.2
1	A	1386	ARG	3.1
1	A	1385	LYS	3.0
1	B	1512	LYS	3.0
1	A	1361	ASP	2.9
1	B	1388	LEU	2.7
1	A	1383	LYS	2.6
1	B	1386	ARG	2.6
1	A	1363	GLY	2.6
1	A	1513	ILE	2.4
1	A	1411	GLN	2.4
1	B	1483	ARG	2.2
1	B	1363	GLY	2.1
1	B	1516	GLN	2.1
1	A	1387	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*


*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	1603	1/1	0.86	0.08	62,62,62,62	0
2	MG	B	1601	1/1	0.97	0.11	28,28,28,28	0
3	GDP	A	1602	28/28	0.98	0.11	24,28,32,34	0
3	GDP	B	1602	28/28	0.98	0.10	25,32,37,37	0
2	MG	B	1603	1/1	0.98	0.14	63,63,63,63	0
2	MG	A	1601	1/1	0.99	0.13	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.