



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 10:15 pm BST

PDB ID : 2OKI  
Title : Crystal structure of dimeric form of PfFabZ in crystal form2  
Authors : Swarnamukhi, P.L.; Sharma, S.K.; Padala, P.; Surolia, N.; Surolia, A.; Suguna, K.  
Deposited on : 2007-01-17  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	136	
1	B	136	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-hydroxyacyl-ACP dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	0	0
			970	639	160	166	5			
1	B	118	Total	C	N	O	S	0	0	0
			874	575	144	151	4			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	O	0	0
			6	6		
2	B	5	Total	O	0	0
			5	5		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Beta-hydroxyacyl-ACP dehydratase

Chain A: 



#### • Molecule 1: Beta-hydroxyacyl-ACP dehydratase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.97Å 81.93Å 91.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.28 – 2.70 23.28 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.4 (23.28-2.70) 96.4 (23.28-2.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.98 (at 2.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.258 , 0.286 0.305 , 0.358	Depositor DCC
$R_{free}$ test set	809 reflections (9.73%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.2	Xtriage
Anisotropy	1.033	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 91.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	1855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/990	0.76	1/1336 (0.1%)
1	B	0.48	0/888	0.77	1/1200 (0.1%)
All	All	0.49	0/1878	0.77	2/2536 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ILE	CG1-CB-CG2	-5.21	99.95	111.40
1	B	130	PHE	N-CA-CB	5.04	119.67	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	970	0	1013	65	0
1	B	874	0	911	62	0
2	A	6	0	0	0	0
2	B	5	0	0	0	0
All	All	1855	0	1924	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:PHE:HB3	1:A:135:PRO:HD2	1.20	1.17
1:A:134:PHE:CB	1:A:135:PRO:HD2	1.95	0.90
1:A:134:PHE:HB3	1:A:135:PRO:CD	2.01	0.90
1:A:143:VAL:HG13	1:B:143:VAL:HB	1.58	0.86
1:A:107:LYS:HZ3	1:A:120:LEU:HD11	1.45	0.81
1:A:117:ILE:HD13	1:A:156:LEU:HB2	1.66	0.78
1:A:176:GLY:H	1:B:175:ASP:CG	1.89	0.76
1:B:117:ILE:HD13	1:B:152:LEU:HD22	1.70	0.74
1:A:174:VAL:HG12	1:B:177:VAL:CG2	2.18	0.74
1:A:117:ILE:CG2	1:A:156:LEU:HG	2.18	0.73
1:B:147:GLU:O	1:B:151:GLN:HG3	1.88	0.73
1:B:109:ILE:HD13	1:B:120:LEU:HG	1.70	0.72
1:A:127:GLU:HG2	1:A:129:PHE:CE2	2.25	0.72
1:A:117:ILE:HG23	1:A:156:LEU:HG	1.72	0.72
1:B:153:ALA:HB1	1:B:207:LEU:HG	1.76	0.68
1:A:99:ARG:CZ	1:A:99:ARG:HB3	2.24	0.67
1:A:111:MET:HG3	1:A:117:ILE:HG22	1.77	0.66
1:B:195:LEU:HD13	1:B:196:ILE:N	2.11	0.66
1:B:124:SER:O	1:B:127:GLU:HG2	1.96	0.66
1:A:167:ASN:OD1	1:A:228:LEU:HD11	1.96	0.65
1:A:123:VAL:HG12	1:A:185:PRO:HA	1.79	0.65
1:A:100:TYR:CD1	1:A:101:PRO:HD2	2.32	0.64
1:B:177:VAL:O	1:B:178:ARG:HD2	1.97	0.64
1:B:139:ILE:HD12	1:B:139:ILE:N	2.12	0.64
1:A:174:VAL:H	1:B:177:VAL:HG22	1.62	0.63
1:B:212:TYR:HA	1:B:216:LYS:O	1.99	0.63
1:A:174:VAL:HG12	1:B:177:VAL:HG22	1.79	0.62
1:B:195:LEU:HA	1:B:207:LEU:HD13	1.81	0.62
1:A:207:LEU:HD12	1:A:207:LEU:N	2.15	0.61
1:B:139:ILE:HG23	1:B:183:VAL:O	1.99	0.61
1:A:218:VAL:HG23	1:A:219:ILE:HG22	1.83	0.60
1:A:174:VAL:HG12	1:B:177:VAL:HG21	1.85	0.59
1:A:177:VAL:HA	1:A:220:ASN:O	2.02	0.58
1:A:122:GLN:HE21	1:A:122:GLN:HA	1.66	0.58
1:B:138:GLN:C	1:B:139:ILE:HD12	2.25	0.58
1:B:169:PHE:HA	1:B:227:ALA:O	2.04	0.58
1:B:193:ALA:HA	1:B:208:SER:O	2.03	0.58
1:A:101:PRO:HG2	1:A:102:PHE:CD2	2.39	0.57
1:B:139:ILE:HG22	1:B:140:MET:N	2.19	0.57
1:A:175:ASP:OD1	1:B:178:ARG:NH2	2.37	0.56
1:A:99:ARG:NH2	1:A:171:PHE:H	2.03	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:LYS:HB3	1:A:218:VAL:HG12	1.87	0.56
1:B:207:LEU:N	1:B:207:LEU:HD22	2.22	0.55
1:A:147:GLU:O	1:A:151:GLN:HG3	2.07	0.55
1:A:123:VAL:CG1	1:A:185:PRO:HA	2.36	0.55
1:A:96:LEU:HB3	1:A:97:PRO:HD2	1.88	0.54
1:B:151:GLN:O	1:B:155:ILE:HG13	2.07	0.54
1:A:117:ILE:HD11	1:A:152:LEU:HD13	1.90	0.54
1:B:123:VAL:HG21	1:B:183:VAL:CG1	2.37	0.54
1:A:175:ASP:HB3	1:B:175:ASP:OD2	2.08	0.54
1:A:193:ALA:HA	1:A:208:SER:O	2.08	0.53
1:A:177:VAL:HG22	1:A:221:ILE:HG12	1.90	0.53
1:B:196:ILE:HB	1:B:206:LYS:O	2.07	0.53
1:A:101:PRO:HG3	1:B:130:PHE:HB3	1.90	0.53
1:A:107:LYS:NZ	1:A:120:LEU:HD11	2.20	0.52
1:A:127:GLU:HG2	1:A:129:PHE:HE2	1.71	0.52
1:B:141:PRO:HD2	1:B:144:LEU:HD12	1.91	0.52
1:A:174:VAL:N	1:B:177:VAL:HG22	2.24	0.52
1:A:104:LEU:HB2	1:A:148:ALA:HA	1.92	0.52
1:B:181:LYS:HG3	1:B:182:PRO:HD2	1.92	0.52
1:B:189:LEU:HD22	1:B:191:MET:HG3	1.91	0.51
1:B:100:TYR:CD1	1:B:101:PRO:HD2	2.46	0.51
1:B:175:ASP:OD2	1:B:176:GLY:N	2.42	0.51
1:B:213:VAL:CG2	1:B:218:VAL:HG11	2.40	0.51
1:A:143:VAL:CG1	1:B:143:VAL:HB	2.36	0.51
1:A:151:GLN:O	1:A:155:ILE:HG13	2.10	0.50
1:A:156:LEU:HD22	1:A:156:LEU:O	2.11	0.49
1:A:132:GLY:O	1:A:133:HIS:HB3	2.12	0.49
1:A:116:THR:C	1:A:117:ILE:CG2	2.80	0.49
1:B:117:ILE:CD1	1:B:152:LEU:HD22	2.42	0.48
1:A:168:LEU:O	1:A:228:LEU:HD22	2.13	0.47
1:B:190:THR:HB	1:B:212:TYR:HB2	1.96	0.47
1:A:107:LYS:HZ3	1:A:107:LYS:HB3	1.79	0.47
1:A:153:ALA:HB1	1:A:207:LEU:HD22	1.96	0.47
1:B:113:PRO:O	1:B:115:LYS:N	2.47	0.47
1:A:142:GLY:O	1:A:145:GLN:HB2	2.15	0.47
1:A:95:ILE:HD13	1:A:95:ILE:H	1.80	0.46
1:A:126:ASN:CG	1:A:126:ASN:O	2.54	0.46
1:B:184:LEU:O	1:B:187:ASP:HB2	2.15	0.46
1:B:207:LEU:O	1:B:223:GLU:HA	2.15	0.46
1:B:122:GLN:HG2	1:B:188:THR:OG1	2.16	0.46
1:B:207:LEU:N	1:B:207:LEU:CD2	2.78	0.45

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:196:ILE:HD11	1:B:208:SER:HB3	1.97	0.45
1:B:109:ILE:HD13	1:B:120:LEU:CG	2.43	0.45
1:B:195:LEU:HD13	1:B:195:LEU:C	2.37	0.45
1:B:167:ASN:HD22	1:B:167:ASN:HA	1.58	0.45
1:B:170:LEU:HD12	1:B:170:LEU:O	2.18	0.44
1:B:206:LYS:NZ	1:B:223:GLU:CB	2.81	0.43
1:A:176:GLY:N	1:B:175:ASP:OD1	2.52	0.43
1:B:101:PRO:HG2	1:B:102:PHE:CD1	2.53	0.43
1:A:143:VAL:HG11	1:B:143:VAL:O	2.18	0.43
1:B:219:ILE:O	1:B:219:ILE:HG23	2.19	0.42
1:B:103:LEU:HD12	1:B:151:GLN:OE1	2.18	0.42
1:A:100:TYR:CG	1:A:101:PRO:HD2	2.54	0.42
1:A:140:MET:HA	1:A:141:PRO:HD3	1.84	0.42
1:A:176:GLY:H	1:B:175:ASP:CB	2.32	0.42
1:B:130:PHE:HB2	1:B:131:ASN:H	1.52	0.42
1:B:123:VAL:HG21	1:B:183:VAL:HG12	2.01	0.42
1:A:116:THR:C	1:A:117:ILE:HG23	2.39	0.42
1:A:117:ILE:HG21	1:A:117:ILE:HD13	1.79	0.42
1:A:206:LYS:C	1:A:207:LEU:HD12	2.40	0.42
1:A:158:LEU:HA	1:A:158:LEU:HD12	1.93	0.42
1:A:124:SER:HB2	1:A:127:GLU:OE1	2.19	0.42
1:B:176:GLY:HA3	1:B:222:SER:OG	2.20	0.42
1:A:117:ILE:CD1	1:A:152:LEU:HD13	2.50	0.41
1:A:109:ILE:O	1:A:110:TYR:HB2	2.20	0.41
1:A:122:GLN:NE2	1:A:122:GLN:HA	2.32	0.41
1:B:177:VAL:HA	1:B:220:ASN:O	2.21	0.41
1:A:129:PHE:CB	1:A:144:LEU:CD1	2.98	0.41
1:B:139:ILE:O	1:B:141:PRO:HD3	2.21	0.40
1:B:139:ILE:CG2	1:B:140:MET:N	2.84	0.40
1:A:134:PHE:CB	1:A:135:PRO:CD	2.77	0.40
1:B:106:ASP:OD2	1:B:121:LYS:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/136 (88%)	102 (85%)	9 (8%)	9 (8%)	1	1
1	B	110/136 (81%)	97 (88%)	11 (10%)	2 (2%)	8	21
All	All	230/272 (85%)	199 (86%)	20 (9%)	11 (5%)	2	4

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	PRO
1	A	136	GLN
1	B	114	ASN
1	A	113	PRO
1	A	114	ASN
1	A	171	PHE
1	A	97	PRO
1	A	132	GLY
1	A	133	HIS
1	B	127	GLU
1	A	219	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/118 (90%)	89 (84%)	17 (16%)	2	6
1	B	93/118 (79%)	84 (90%)	9 (10%)	8	19
All	All	199/236 (84%)	173 (87%)	26 (13%)	4	10

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	LYS
1	A	95	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	99	ARG
1	A	127	GLU
1	A	130	PHE
1	A	136	GLN
1	A	146	ILE
1	A	152	LEU
1	A	156	LEU
1	A	167	ASN
1	A	168	LEU
1	A	175	ASP
1	A	181	LYS
1	A	198	PHE
1	A	207	LEU
1	A	212	TYR
1	A	228	LEU
1	B	99	ARG
1	B	113	PRO
1	B	130	PHE
1	B	152	LEU
1	B	167	ASN
1	B	177	VAL
1	B	180	LYS
1	B	189	LEU
1	B	201	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	GLN
1	A	194	ASN
1	A	220	ASN
1	B	167	ASN
1	B	192	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.