



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 02:00 pm BST

PDB ID : 5OK7
Title : Conservatively refined structure of Gan1D-E170Q, a catalytic mutant of a putative 6-phospho-beta-galactosidase from *Geobacillus stearothermophilus*
Authors : Lansky, S.; Zehavi, A.; Shoham, Y.; Shoham, G.
Deposited on : 2017-07-25
Resolution : 1.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1385 (1.36-1.32)
Clashscore	141614	1417 (1.36-1.32)
Ramachandran outliers	138981	1397 (1.36-1.32)
Sidechain outliers	138945	1397 (1.36-1.32)
RSRZ outliers	127900	1369 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	<div> <div>4%</div> <div> <div></div> <div>92%</div> <div>5%</div> </div> </div>
1	B	485	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>5%</div> </div> </div>
1	C	485	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div>5%</div> </div> </div>
1	D	485	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>5%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18141 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative 6-phospho-beta-galactobiosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	462	Total	C	N	O	S	0	25	0
			3952	2552	674	712	14			
1	B	462	Total	C	N	O	S	0	23	0
			3950	2546	672	718	14			
1	C	462	Total	C	N	O	S	0	22	0
			3932	2537	672	709	14			
1	D	462	Total	C	N	O	S	0	19	0
			3903	2519	670	701	13			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP W8QF82
A	-5	ILE	-	expression tag	UNP W8QF82
A	-4	HIS	-	expression tag	UNP W8QF82
A	-3	HIS	-	expression tag	UNP W8QF82
A	-2	HIS	-	expression tag	UNP W8QF82
A	-1	HIS	-	expression tag	UNP W8QF82
A	0	HIS	-	expression tag	UNP W8QF82
A	1	HIS	-	expression tag	UNP W8QF82
A	170	GLN	GLU	engineered mutation	UNP W8QF82
B	-6	MET	-	initiating methionine	UNP W8QF82
B	-5	ILE	-	expression tag	UNP W8QF82
B	-4	HIS	-	expression tag	UNP W8QF82
B	-3	HIS	-	expression tag	UNP W8QF82
B	-2	HIS	-	expression tag	UNP W8QF82
B	-1	HIS	-	expression tag	UNP W8QF82
B	0	HIS	-	expression tag	UNP W8QF82
B	1	HIS	-	expression tag	UNP W8QF82
B	170	GLN	GLU	engineered mutation	UNP W8QF82
C	-6	MET	-	initiating methionine	UNP W8QF82
C	-5	ILE	-	expression tag	UNP W8QF82
C	-4	HIS	-	expression tag	UNP W8QF82

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	HIS	-	expression tag	UNP W8QF82
C	-2	HIS	-	expression tag	UNP W8QF82
C	-1	HIS	-	expression tag	UNP W8QF82
C	0	HIS	-	expression tag	UNP W8QF82
C	1	HIS	-	expression tag	UNP W8QF82
C	170	GLN	GLU	engineered mutation	UNP W8QF82
D	-6	MET	-	initiating methionine	UNP W8QF82
D	-5	ILE	-	expression tag	UNP W8QF82
D	-4	HIS	-	expression tag	UNP W8QF82
D	-3	HIS	-	expression tag	UNP W8QF82
D	-2	HIS	-	expression tag	UNP W8QF82
D	-1	HIS	-	expression tag	UNP W8QF82
D	0	HIS	-	expression tag	UNP W8QF82
D	1	HIS	-	expression tag	UNP W8QF82
D	170	GLN	GLU	engineered mutation	UNP W8QF82

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		
3	D	1	Total	C	N	0	0
			5	3	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	565	Total 565	O 565	0	0
4	B	579	Total 579	O 579	0	0
4	C	622	Total 622	O 622	0	0
4	D	551	Total 551	O 551	0	0

- Molecule 1: Putative 6-phospho-beta-galactobiosidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.55Å 97.43Å 105.25Å 90.00° 97.82° 90.00°	Depositor
Resolution (Å)	43.91 – 1.34 43.91 – 1.34	Depositor EDS
% Data completeness (in resolution range)	91.4 (43.91-1.34) 91.4 (43.91-1.34)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.34Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.123 , 0.146 0.124 , 0.146	Depositor DCC
R_{free} test set	4096 reflections (0.99%)	wwPDB-VP
Wilson B-factor (Å ²)	10.4	Xtriage
Anisotropy	0.444	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 52.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	18141	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7729e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/4145	0.58	0/5633
1	B	0.36	0/4128	0.59	0/5615
1	C	0.36	0/4124	0.58	0/5608
1	D	0.36	0/4083	0.57	0/5551
All	All	0.36	0/16480	0.58	0/22407

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3952	0	3816	11	0
1	B	3950	0	3778	17	0
1	C	3932	0	3782	18	0
1	D	3903	0	3743	21	0
2	A	24	0	32	3	0
2	B	30	0	40	2	0
2	C	12	0	16	0	0
2	D	6	0	8	0	0
3	A	5	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	5	0	5	0	0
3	D	5	0	5	1	0
4	A	565	0	0	3	0
4	B	579	0	0	6	0
4	C	622	0	0	3	0
4	D	551	0	0	7	0
All	All	18141	0	15230	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:97[A]:ASN:ND2	4:C:601:HOH:O	2.21	0.71
1:D:97[A]:ASN:ND2	4:D:602:HOH:O	2.26	0.69
1:C:214:HIS:HE1	1:D:383:GLU:HG2	1.58	0.69
1:A:97[B]:ASN:ND2	4:A:603:HOH:O	2.27	0.65
1:D:249[A]:PHE:CE2	1:D:333:ILE:HD11	2.39	0.57
1:B:341:ARG:NH2	1:B:351[A]:ASP:OD2	2.36	0.56
1:D:93:ASN:HA	1:D:149:ARG:HH22	1.74	0.53
1:D:438:GLN:HG3	4:D:677:HOH:O	2.08	0.53
1:B:348[B]:THR:HB	1:B:354:ILE:HD11	1.91	0.52
1:C:473:THR:HG21	1:C:477[B]:GLU:HG3	1.92	0.52
1:D:207:LYS:NZ	4:D:603:HOH:O	2.26	0.51
1:A:149:ARG:HD3	4:A:790:HOH:O	2.10	0.51
1:D:149:ARG:NH2	4:D:611:HOH:O	2.43	0.51
1:D:245:ASN:ND2	4:D:612:HOH:O	2.44	0.51
1:A:266:ALA:HB1	2:A:504:GOL:H32	1.93	0.50
1:B:99:LYS:HD2	4:B:742:HOH:O	2.11	0.49
1:A:336:LEU:CD2	2:A:504:GOL:H2	2.42	0.49
1:B:99:LYS:HE3	4:B:1046:HOH:O	2.13	0.49
1:D:93:ASN:CB	1:D:149:ARG:HH12	2.26	0.48
1:B:472:GLU:OE2	4:B:601:HOH:O	2.20	0.48
1:C:249:PHE:CE2	1:C:333:ILE:HD11	2.48	0.47
1:A:378:GLU:HG3	1:A:425:TRP:HB2	1.95	0.47
1:B:378:GLU:HG3	1:B:425:TRP:HB2	1.96	0.47
1:C:312:ASP:N	1:C:312:ASP:OD1	2.47	0.46
1:D:312:ASP:OD1	1:D:312:ASP:N	2.47	0.46
1:D:92:GLY:HA2	1:D:128:PRO:HG2	1.98	0.46
1:D:292:LYS:HB3	4:D:693:HOH:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:112[A]:ARG:HD3	1:D:161:ARG:HB3	1.97	0.45
1:B:425:TRP:HA	1:B:426:SER:HA	1.80	0.45
1:C:153:THR:O	1:C:157[B]:ARG:HG2	2.17	0.45
1:D:378:GLU:HG3	1:D:425:TRP:HB2	1.98	0.45
1:C:378:GLU:HG3	1:C:425:TRP:HB2	1.98	0.44
1:D:157[A]:ARG:HD2	3:D:502:IMD:C4	2.47	0.44
1:B:341:ARG:NH1	1:B:351[A]:ASP:OD1	2.37	0.44
1:C:157[A]:ARG:HG3	1:C:158:PHE:CE1	2.52	0.43
1:D:93:ASN:HB3	1:D:149:ARG:HH12	1.83	0.43
1:B:348[A]:THR:OG1	1:B:352[A]:TRP:HB2	2.19	0.43
1:A:394:ASN:ND2	4:A:620:HOH:O	2.51	0.43
1:B:153:THR:O	1:B:157[B]:ARG:HG2	2.19	0.43
1:B:435:ASN:HB3	2:B:505:GOL:H32	2.01	0.43
1:B:400[A]:ASP:OD1	4:B:602:HOH:O	2.21	0.43
1:C:171:GLN:HA	1:C:174:PHE:CE2	2.54	0.42
1:B:180[B]:ARG:HB2	1:B:194[B]:MET:SD	2.59	0.42
2:B:504:GOL:O1	4:B:603:HOH:O	2.22	0.42
1:D:249[B]:PHE:HZ	1:D:270:TYR:HB3	1.84	0.42
1:C:32:GLY:HA3	1:C:97[B]:ASN:HD21	1.84	0.42
1:D:175:ILE:HG21	1:D:198:ASN:HB2	2.02	0.42
1:C:214:HIS:CE1	1:D:383:GLU:HG2	2.47	0.42
1:A:16:GLY:HA2	1:A:78:ALA:O	2.20	0.42
1:A:336:LEU:HD21	2:A:504:GOL:H2	2.01	0.42
1:A:47:ARG:HA	1:A:47:ARG:HD3	1.87	0.42
1:A:180[A]:ARG:HD3	1:A:191[A]:MET:HG2	2.02	0.41
1:B:149:ARG:HD3	4:B:707:HOH:O	2.19	0.41
1:B:16:GLY:HA2	1:B:78:ALA:O	2.21	0.41
1:C:425:TRP:HA	1:C:426:SER:HA	1.80	0.41
1:C:94:GLY:O	1:C:149:ARG:NH2	2.44	0.41
1:B:341:ARG:HH22	1:B:351[A]:ASP:CG	2.23	0.41
1:C:292:LYS:HB3	4:C:737:HOH:O	2.21	0.41
1:A:350[B]:TRP:HZ2	1:A:384:PHE:HZ	1.69	0.41
1:B:171:GLN:HA	1:B:174:PHE:CE2	2.55	0.41
1:D:310:PRO:HD2	4:D:671:HOH:O	2.21	0.41
1:C:180:ARG:HB2	1:C:194[B]:MET:SD	2.60	0.41
1:C:148:ASP:HA	1:C:208[A]:VAL:HG12	2.03	0.40
1:C:194[B]:MET:HE1	4:C:1029:HOH:O	2.21	0.40
1:C:16:GLY:HA2	1:C:78:ALA:O	2.21	0.40
1:D:171:GLN:HA	1:D:174:PHE:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	484/485 (100%)	469 (97%)	15 (3%)	0	100	100
1	B	482/485 (99%)	466 (97%)	16 (3%)	0	100	100
1	C	481/485 (99%)	465 (97%)	16 (3%)	0	100	100
1	D	477/485 (98%)	463 (97%)	14 (3%)	0	100	100
All	All	1924/1940 (99%)	1863 (97%)	61 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/412 (101%)	413 (99%)	4 (1%)	76	47
1	B	416/412 (101%)	415 (100%)	1 (0%)	93	82
1	C	415/412 (101%)	412 (99%)	3 (1%)	84	62
1	D	409/412 (99%)	406 (99%)	3 (1%)	84	62
All	All	1657/1648 (100%)	1646 (99%)	11 (1%)	88	62

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	312	ASP
1	A	430	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	478[A]	LEU
1	A	478[B]	LEU
1	B	430	LEU
1	C	24[A]	VAL
1	C	24[B]	VAL
1	C	430	LEU
1	D	430	LEU
1	D	452[A]	SER
1	D	452[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	504	-	5,5,5	0.36	0	5,5,5	0.32	0
2	GOL	C	502	-	5,5,5	0.35	0	5,5,5	0.44	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	502	-	5,5,5	0.39	0	5,5,5	0.39	0
2	GOL	D	501	-	5,5,5	0.37	0	5,5,5	0.33	0
3	IMD	A	505	-	3,5,5	0.40	0	4,5,5	0.61	0
2	GOL	B	505	-	5,5,5	0.37	0	5,5,5	0.32	0
2	GOL	C	501	-	5,5,5	0.41	0	5,5,5	0.52	0
2	GOL	B	501	-	5,5,5	0.35	0	5,5,5	0.34	0
2	GOL	A	504	-	5,5,5	0.39	0	5,5,5	0.40	0
2	GOL	B	502	-	5,5,5	0.36	0	5,5,5	0.34	0
3	IMD	D	502	-	3,5,5	0.41	0	4,5,5	0.56	0
3	IMD	C	503	-	3,5,5	0.41	0	4,5,5	0.54	0
2	GOL	A	501	-	5,5,5	0.36	0	5,5,5	0.33	0
2	GOL	A	503	-	5,5,5	0.39	0	5,5,5	0.31	0
2	GOL	B	503	-	5,5,5	0.36	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	504	-	-	2/4/4/4	-
3	IMD	A	505	-	-	-	0/1/1/1
2	GOL	C	502	-	-	1/4/4/4	-
2	GOL	A	502	-	-	0/4/4/4	-
2	GOL	D	501	-	-	2/4/4/4	-
2	GOL	A	501	-	-	0/4/4/4	-
2	GOL	B	505	-	-	2/4/4/4	-
2	GOL	C	501	-	-	2/4/4/4	-
2	GOL	B	501	-	-	0/4/4/4	-
2	GOL	A	504	-	-	2/4/4/4	-
2	GOL	B	502	-	-	0/4/4/4	-
3	IMD	D	502	-	-	-	0/1/1/1
3	IMD	C	503	-	-	-	0/1/1/1
2	GOL	A	503	-	-	1/4/4/4	-
2	GOL	B	503	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	504	GOL	O1-C1-C2-C3
2	B	505	GOL	O1-C1-C2-C3
2	C	501	GOL	O1-C1-C2-O2
2	C	501	GOL	O1-C1-C2-C3
2	A	504	GOL	C1-C2-C3-O3
2	B	504	GOL	O1-C1-C2-O2
2	B	505	GOL	O1-C1-C2-O2
2	D	501	GOL	O1-C1-C2-O2
2	B	503	GOL	O1-C1-C2-O2
2	B	503	GOL	O1-C1-C2-C3
2	A	504	GOL	O2-C2-C3-O3
2	D	501	GOL	O1-C1-C2-C3
2	C	502	GOL	O1-C1-C2-O2
2	A	503	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	504	GOL	1	0
2	B	505	GOL	1	0
2	A	504	GOL	3	0
3	D	502	IMD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	462/485 (95%)	-0.04	19 (4%) 37 43	8, 14, 29, 56	0
1	B	462/485 (95%)	-0.09	16 (3%) 44 50	8, 13, 26, 57	0
1	C	462/485 (95%)	-0.07	15 (3%) 47 54	8, 12, 23, 73	0
1	D	462/485 (95%)	-0.04	15 (3%) 47 54	8, 14, 27, 69	0
All	All	1848/1940 (95%)	-0.06	65 (3%) 44 50	8, 13, 27, 73	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	312	ASP	9.9
1	D	311	PRO	9.0
1	A	350[A]	TRP	8.7
1	B	315	GLY	8.5
1	A	315	GLY	8.3
1	B	329	THR	8.0
1	C	312	ASP	7.8
1	D	329	THR	7.7
1	D	352	TRP	7.6
1	B	312	ASP	7.5
1	A	329	THR	7.4
1	C	313	GLY	7.4
1	D	315	GLY	7.2
1	D	313	GLY	7.2
1	C	311	PRO	7.1
1	A	313	GLY	7.0
1	C	329	THR	6.6
1	A	312	ASP	6.6
1	C	314	VAL	6.2
1	D	314	VAL	5.8
1	C	4	ARG	5.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	310	PRO	5.5
1	C	315	GLY	5.2
1	A	311	PRO	5.1
1	C	5	HIS	5.1
1	D	4	ARG	5.1
1	B	313	GLY	5.0
1	B	311	PRO	5.0
1	A	352	TRP	5.0
1	A	314	VAL	4.6
1	C	310	PRO	4.6
1	A	389	PRO	4.5
1	B	314	VAL	4.4
1	C	352[A]	TRP	4.4
1	B	5	HIS	4.4
1	D	350	TRP	4.2
1	B	352[A]	TRP	4.1
1	D	5	HIS	3.9
1	B	392	ILE	3.2
1	B	310	PRO	3.1
1	A	392	ILE	3.1
1	A	5	HIS	3.1
1	B	389	PRO	3.1
1	B	350[A]	TRP	3.0
1	A	310	PRO	2.9
1	B	478	LEU	2.9
1	A	478[A]	LEU	2.8
1	D	330	SER	2.8
1	D	351	ASP	2.7
1	A	346	ASP	2.7
1	B	4	ARG	2.6
1	C	330	SER	2.6
1	B	12	GLU	2.5
1	B	160	ASP	2.5
1	A	347	THR	2.4
1	C	350	TRP	2.3
1	A	351	ASP	2.3
1	A	384	PHE	2.2
1	A	282	PRO	2.1
1	A	160	ASP	2.1
1	C	94	GLY	2.1
1	D	282	PRO	2.1
1	C	351	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	160	ASP	2.0
1	D	149	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	IMD	D	502	5/5	0.61	0.25	52,53,53,53	5
2	GOL	B	505	6/6	0.68	0.21	48,51,51,52	0
2	GOL	B	504	6/6	0.68	0.19	59,60,60,61	0
3	IMD	C	503	5/5	0.78	0.18	53,53,53,54	0
2	GOL	A	504	6/6	0.80	0.20	45,45,45,45	0
3	IMD	A	505	5/5	0.83	0.11	45,45,45,45	0
2	GOL	B	503	6/6	0.85	0.15	41,41,41,42	0
2	GOL	A	503	6/6	0.86	0.12	41,41,42,42	0
2	GOL	C	501	6/6	0.86	0.27	24,30,33,36	0
2	GOL	D	501	6/6	0.87	0.15	42,42,43,43	0
2	GOL	C	502	6/6	0.93	0.14	37,38,38,39	0
2	GOL	A	501	6/6	0.94	0.09	16,20,21,23	0
2	GOL	B	502	6/6	0.94	0.09	15,20,22,23	0
2	GOL	B	501	6/6	0.95	0.10	15,20,23,25	0
2	GOL	A	502	6/6	0.97	0.07	17,22,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.