



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 09:03 pm BST

PDB ID : 3OLP
Title : Crystal structure of a bacterial phosphoglucomutase, an enzyme important in the virulence of multiple human pathogens
Authors : Beamer, L.J.; Mehra-Chaudhary, R.
Deposited on : 2010-08-26
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	570	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>9% • 5%</div> </div> </div>
1	B	570	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>6% • 5%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8523 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoglucomutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	543	Total	C	N	O	S	0	9	0
			4034	2546	700	772	16			
1	B	543	Total	C	N	O	S	0	3	0
			4012	2527	706	763	16			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	EXPRESSION TAG	UNP Q8ZQW9
A	-22	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
A	-21	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
A	-20	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
A	-19	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
A	-18	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
A	-17	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
A	-16	SER	-	EXPRESSION TAG	UNP Q8ZQW9
A	-15	SER	-	EXPRESSION TAG	UNP Q8ZQW9
A	-14	GLY	-	EXPRESSION TAG	UNP Q8ZQW9
A	-13	VAL	-	EXPRESSION TAG	UNP Q8ZQW9
A	-12	ASP	-	EXPRESSION TAG	UNP Q8ZQW9
A	-11	LEU	-	EXPRESSION TAG	UNP Q8ZQW9
A	-10	GLY	-	EXPRESSION TAG	UNP Q8ZQW9
A	-9	THR	-	EXPRESSION TAG	UNP Q8ZQW9
A	-8	GLU	-	EXPRESSION TAG	UNP Q8ZQW9
A	-7	ASN	-	EXPRESSION TAG	UNP Q8ZQW9
A	-6	LEU	-	EXPRESSION TAG	UNP Q8ZQW9
A	-5	TYR	-	EXPRESSION TAG	UNP Q8ZQW9
A	-4	PHE	-	EXPRESSION TAG	UNP Q8ZQW9
A	-3	GLN	-	EXPRESSION TAG	UNP Q8ZQW9
A	-2	SER	-	EXPRESSION TAG	UNP Q8ZQW9
A	-1	ASN	-	EXPRESSION TAG	UNP Q8ZQW9
A	0	ALA	-	EXPRESSION TAG	UNP Q8ZQW9
B	-23	MET	-	EXPRESSION TAG	UNP Q8ZQW9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
B	-21	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
B	-20	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
B	-19	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
B	-18	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
B	-17	HIS	-	EXPRESSION TAG	UNP Q8ZQW9
B	-16	SER	-	EXPRESSION TAG	UNP Q8ZQW9
B	-15	SER	-	EXPRESSION TAG	UNP Q8ZQW9
B	-14	GLY	-	EXPRESSION TAG	UNP Q8ZQW9
B	-13	VAL	-	EXPRESSION TAG	UNP Q8ZQW9
B	-12	ASP	-	EXPRESSION TAG	UNP Q8ZQW9
B	-11	LEU	-	EXPRESSION TAG	UNP Q8ZQW9
B	-10	GLY	-	EXPRESSION TAG	UNP Q8ZQW9
B	-9	THR	-	EXPRESSION TAG	UNP Q8ZQW9
B	-8	GLU	-	EXPRESSION TAG	UNP Q8ZQW9
B	-7	ASN	-	EXPRESSION TAG	UNP Q8ZQW9
B	-6	LEU	-	EXPRESSION TAG	UNP Q8ZQW9
B	-5	TYR	-	EXPRESSION TAG	UNP Q8ZQW9
B	-4	PHE	-	EXPRESSION TAG	UNP Q8ZQW9
B	-3	GLN	-	EXPRESSION TAG	UNP Q8ZQW9
B	-2	SER	-	EXPRESSION TAG	UNP Q8ZQW9
B	-1	ASN	-	EXPRESSION TAG	UNP Q8ZQW9
B	0	ALA	-	EXPRESSION TAG	UNP Q8ZQW9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	264	Total	O	0	0
			264	264		
4	B	201	Total	O	0	0
			201	201		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphoglucosyltransferase



• Molecule 1: Phosphoglucosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.12Å 85.12Å 111.88Å 90.00° 94.42° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 45.07 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.8 (50.00-1.95) 98.8 (45.07-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.166 , 0.195 0.165 , 0.194	Depositor DCC
R_{free} test set	3925 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.235	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8523	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/4144	0.60	1/5649 (0.0%)
1	B	0.47	0/4104	0.57	1/5590 (0.0%)
All	All	0.51	0/8248	0.59	2/11239 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	349	LEU	CA-CB-CG	5.38	127.68	115.30
1	A	6	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4034	0	3905	38	0
1	B	4012	0	3881	24	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	264	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	201	0	0	2	0
All	All	8523	0	7786	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:LEU:HD11	1:A:287:MET:HE3	1.40	1.00
1:B:236:LEU:HD11	1:B:287:MET:HE1	1.49	0.91
1:B:391:GLU:HG2	1:B:447:ARG:HH21	1.41	0.84
1:A:430[B]:GLN:OE1	4:A:550:HOH:O	1.99	0.79
1:B:221:MET:O	1:B:225:GLN:HG3	1.83	0.78
1:A:430[B]:GLN:NE2	4:A:771:HOH:O	2.11	0.77
1:A:233[B]:VAL:HG12	1:A:301:PHE:HB2	1.69	0.73
1:B:57:ASN:H	1:B:60:HIS:HD2	1.38	0.71
1:B:40:VAL:H	1:B:178:ASN:HD21	1.38	0.70
1:A:233[A]:VAL:CG2	1:A:255:LEU:HD11	2.22	0.70
1:A:233[A]:VAL:HG23	1:A:255:LEU:HD11	1.73	0.69
1:A:40:VAL:H	1:A:178:ASN:HD21	1.38	0.69
1:B:357:ARG:HG2	1:B:440:PHE:CD1	2.28	0.68
1:B:57:ASN:H	1:B:60:HIS:CD2	2.12	0.67
1:B:357:ARG:HG2	1:B:440:PHE:CG	2.30	0.67
1:A:38:HIS:O	1:A:60:HIS:HE1	1.80	0.65
1:A:457:LYS:HG3	1:A:508:PRO:HB3	1.80	0.64
1:A:57:ASN:H	1:A:60:HIS:HD2	1.45	0.62
1:B:287:MET:HA	1:B:287:MET:HE2	1.83	0.61
1:B:236:LEU:HD11	1:B:287:MET:CE	2.26	0.60
1:B:38:HIS:O	1:B:60:HIS:HE1	1.82	0.60
1:A:151:GLU:OE1	4:A:661:HOH:O	2.17	0.59
1:B:225:GLN:HB3	4:B:629:HOH:O	2.02	0.58
1:A:236:LEU:CD1	1:A:287:MET:HE3	2.25	0.57
1:A:307:TYR:HD2	1:A:412:ILE:HD11	1.70	0.56
1:A:307:TYR:CD2	1:A:412:ILE:HD11	2.41	0.56
1:A:6:ARG:NH2	1:A:15:ASP:OD1	2.36	0.55
1:A:57:ASN:H	1:A:60:HIS:CD2	2.24	0.53
1:A:236:LEU:HD21	1:A:287:MET:HE1	1.92	0.52
1:A:40:VAL:N	1:A:178:ASN:HD21	2.07	0.51
1:B:468:VAL:HG23	4:B:701:HOH:O	2.11	0.51
1:B:119:PRO:HB3	1:B:307:TYR:HB2	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:472:THR:HG22	4:A:622:HOH:O	2.10	0.49
1:A:233[A]:VAL:HG23	1:A:255:LEU:CD1	2.41	0.49
1:B:168:ASN:H	1:B:168:ASN:HD22	1.62	0.48
1:B:468:VAL:HG12	1:B:470:ALA:H	1.78	0.48
1:A:147:HIS:CD2	1:A:309:ARG:NH2	2.82	0.48
1:A:349:LEU:HD11	1:A:492:ILE:HG21	1.96	0.48
1:B:156:LYS:NZ	1:B:158:ASN:HD21	2.13	0.47
1:A:233[A]:VAL:HG11	1:A:244:TRP:CD1	2.50	0.46
1:A:430[B]:GLN:HA	1:A:430[B]:GLN:OE1	2.17	0.45
1:A:119:PRO:HB3	1:A:307:TYR:HB2	1.99	0.45
1:B:168:ASN:HD22	1:B:168:ASN:N	2.14	0.45
1:A:464:SER:OG	1:A:467:MET:HB2	2.17	0.44
1:A:452:ALA:O	1:A:513:ASP:OD1	2.35	0.44
1:A:287:MET:HA	1:A:287:MET:CE	2.47	0.44
1:A:542:VAL:HG23	1:A:543:LEU:HD13	2.00	0.44
1:A:99:LEU:HD22	1:A:109:VAL:HG11	2.00	0.43
1:A:473:LEU:HB3	1:A:499:THR:CG2	2.49	0.43
1:B:40:VAL:N	1:B:178:ASN:HD21	2.10	0.42
1:B:333:PHE:CZ	1:B:345:VAL:HG22	2.54	0.42
1:A:147:HIS:CD2	1:A:309:ARG:HH21	2.38	0.42
1:B:463:LEU:HD13	1:B:542:VAL:HG11	2.01	0.42
1:A:235:PRO:HG2	1:A:241:ILE:HD12	2.02	0.42
1:A:373:PHE:CD1	1:A:409:LYS:HG2	2.54	0.42
1:B:349:LEU:CD1	1:B:370:PRO:HA	2.50	0.42
1:B:320[B]:ASN:ND2	1:B:322:ASN:H	2.18	0.41
1:B:99:LEU:HD22	1:B:109:VAL:HG11	2.02	0.41
1:A:46:GLY:HA3	1:A:155:ILE:O	2.22	0.40
1:A:44:THR:HG23	4:A:802:HOH:O	2.20	0.40
1:A:208:VAL:HG22	1:A:243:TYR:CE1	2.56	0.40
1:A:543:LEU:O	1:A:544:LYS:C	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/570 (96%)	540 (98%)	9 (2%)	1 (0%)	47	38
1	B	544/570 (95%)	530 (97%)	14 (3%)	0	100	100
All	All	1094/1140 (96%)	1070 (98%)	23 (2%)	1 (0%)	51	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/444 (92%)	399 (98%)	10 (2%)	49	40
1	B	404/444 (91%)	391 (97%)	13 (3%)	39	27
All	All	813/888 (92%)	790 (97%)	23 (3%)	42	33

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ILE
1	A	241	ILE
1	A	278	ARG
1	A	356	ASP
1	A	391	GLU
1	A	454	SER
1	A	466	GLU
1	A	467	MET
1	A	483	THR
1	A	491	SER
1	B	168	ASN
1	B	190	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	273	LYS
1	B	301	PHE
1	B	349	LEU
1	B	356	ASP
1	B	357	ARG
1	B	390	GLU
1	B	391	GLU
1	B	409	LYS
1	B	454	SER
1	B	509	SER
1	B	511	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	147	HIS
1	A	158	ASN
1	A	178	ASN
1	A	434	ASN
1	B	60	HIS
1	B	158	ASN
1	B	168	ASN
1	B	178	ASN
1	B	250	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	B	714	-	4,4,4	0.92	0	6,6,6	0.51	0
3	PO4	A	713	-	4,4,4	0.90	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	543/570 (95%)	-0.01	21 (3%)	39 49	15, 25, 53, 76	1 (0%)
1	B	543/570 (95%)	0.14	31 (5%)	23 32	17, 32, 65, 103	0
All	All	1086/1140 (95%)	0.06	52 (4%)	30 40	15, 29, 57, 103	1 (0%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	464	SER	7.3
1	B	542	VAL	7.3
1	A	511	THR	7.2
1	B	466	GLU	6.3
1	B	545	ASN	5.9
1	B	468	VAL	5.9
1	A	542	VAL	5.8
1	B	467	MET	5.5
1	B	469	SER	5.3
1	B	470	ALA	5.2
1	B	462	LYS	5.2
1	B	134	GLY	5.0
1	B	465	PRO	5.0
1	B	463	LEU	4.8
1	A	467	MET	4.2
1	A	465	PRO	4.1
1	B	32	GLU	3.9
1	B	461	SER	3.8
1	B	511	THR	3.8
1	A	2	ALA	3.7
1	B	495	LEU	3.6
1	B	34	GLY	3.4
1	B	452	ALA	3.4
1	A	134	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	463	LEU	3.2
1	A	452	ALA	3.2
1	B	459	ALA	3.2
1	A	543	LEU	3.1
1	A	513	ASP	3.1
1	A	544	LYS	3.1
1	A	458	ALA	3.0
1	A	453	THR	2.9
1	A	450	ALA	2.9
1	B	460	LEU	2.9
1	A	515	TYR	2.9
1	B	438	ALA	2.8
1	B	455	ALA	2.8
1	B	544	LYS	2.5
1	B	512	GLU	2.5
1	B	492	ILE	2.4
1	B	539	VAL	2.4
1	A	510	GLY	2.4
1	A	514	ALA	2.4
1	A	460	LEU	2.3
1	A	464	SER	2.3
1	A	34	GLY	2.3
1	B	458	ALA	2.3
1	B	475	GLY	2.2
1	B	453	THR	2.2
1	B	510	GLY	2.1
1	B	315	PRO	2.0
1	A	512	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	714	5/5	0.82	0.18	96,96,96,96	0
3	PO4	A	713	5/5	0.85	0.17	119,119,119,119	0
2	MG	B	547	1/1	0.98	0.09	27,27,27,27	0
2	MG	A	547	1/1	0.99	0.12	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.