



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 08:12 am BST

PDB ID : 5OLC  
Title : Crystal structure of the 3,6-anhydro-D-galactonate cycloisomerase from *Zobellia galactanivorans*  
Authors : Michel, G.; Czjzek, M.; Jam, M.  
Deposited on : 2017-07-27  
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	
1	B	396	
1	C	396	
1	D	396	
1	E	396	
1	F	396	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	396	<div><div></div><div>4%</div><div>68%</div><div>17%</div><div>•</div><div>11%</div></div>
1	H	396	<div><div></div><div>4%</div><div>69%</div><div>17%</div><div>•</div><div>11%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 21906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galactonate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	0	0
			2721	1749	446	513	13			
1	B	351	Total	C	N	O	S	0	0	0
			2709	1742	444	510	13			
1	C	351	Total	C	N	O	S	0	0	0
			2717	1748	446	510	13			
1	D	351	Total	C	N	O	S	0	0	0
			2711	1745	443	510	13			
1	E	351	Total	C	N	O	S	0	0	0
			2735	1758	451	513	13			
1	F	351	Total	C	N	O	S	0	0	0
			2741	1764	451	513	13			
1	G	351	Total	C	N	O	S	0	0	0
			2730	1755	449	513	13			
1	H	351	Total	C	N	O	S	0	0	0
			2745	1767	452	513	13			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP G0L7B8
A	-6	HIS	-	expression tag	UNP G0L7B8
A	-5	HIS	-	expression tag	UNP G0L7B8
A	-4	HIS	-	expression tag	UNP G0L7B8
A	-3	HIS	-	expression tag	UNP G0L7B8
A	-2	HIS	-	expression tag	UNP G0L7B8
A	-1	HIS	-	expression tag	UNP G0L7B8
A	0	GLY	-	expression tag	UNP G0L7B8
A	1	SER	-	expression tag	UNP G0L7B8
B	-7	MET	-	initiating methionine	UNP G0L7B8
B	-6	HIS	-	expression tag	UNP G0L7B8
B	-5	HIS	-	expression tag	UNP G0L7B8
B	-4	HIS	-	expression tag	UNP G0L7B8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	HIS	-	expression tag	UNP G0L7B8
B	-2	HIS	-	expression tag	UNP G0L7B8
B	-1	HIS	-	expression tag	UNP G0L7B8
B	0	GLY	-	expression tag	UNP G0L7B8
B	1	SER	-	expression tag	UNP G0L7B8
C	-7	MET	-	initiating methionine	UNP G0L7B8
C	-6	HIS	-	expression tag	UNP G0L7B8
C	-5	HIS	-	expression tag	UNP G0L7B8
C	-4	HIS	-	expression tag	UNP G0L7B8
C	-3	HIS	-	expression tag	UNP G0L7B8
C	-2	HIS	-	expression tag	UNP G0L7B8
C	-1	HIS	-	expression tag	UNP G0L7B8
C	0	GLY	-	expression tag	UNP G0L7B8
C	1	SER	-	expression tag	UNP G0L7B8
D	-7	MET	-	initiating methionine	UNP G0L7B8
D	-6	HIS	-	expression tag	UNP G0L7B8
D	-5	HIS	-	expression tag	UNP G0L7B8
D	-4	HIS	-	expression tag	UNP G0L7B8
D	-3	HIS	-	expression tag	UNP G0L7B8
D	-2	HIS	-	expression tag	UNP G0L7B8
D	-1	HIS	-	expression tag	UNP G0L7B8
D	0	GLY	-	expression tag	UNP G0L7B8
D	1	SER	-	expression tag	UNP G0L7B8
E	-7	MET	-	initiating methionine	UNP G0L7B8
E	-6	HIS	-	expression tag	UNP G0L7B8
E	-5	HIS	-	expression tag	UNP G0L7B8
E	-4	HIS	-	expression tag	UNP G0L7B8
E	-3	HIS	-	expression tag	UNP G0L7B8
E	-2	HIS	-	expression tag	UNP G0L7B8
E	-1	HIS	-	expression tag	UNP G0L7B8
E	0	GLY	-	expression tag	UNP G0L7B8
E	1	SER	-	expression tag	UNP G0L7B8
F	-7	MET	-	initiating methionine	UNP G0L7B8
F	-6	HIS	-	expression tag	UNP G0L7B8
F	-5	HIS	-	expression tag	UNP G0L7B8
F	-4	HIS	-	expression tag	UNP G0L7B8
F	-3	HIS	-	expression tag	UNP G0L7B8
F	-2	HIS	-	expression tag	UNP G0L7B8
F	-1	HIS	-	expression tag	UNP G0L7B8
F	0	GLY	-	expression tag	UNP G0L7B8
F	1	SER	-	expression tag	UNP G0L7B8
G	-7	MET	-	initiating methionine	UNP G0L7B8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	-6	HIS	-	expression tag	UNP G0L7B8
G	-5	HIS	-	expression tag	UNP G0L7B8
G	-4	HIS	-	expression tag	UNP G0L7B8
G	-3	HIS	-	expression tag	UNP G0L7B8
G	-2	HIS	-	expression tag	UNP G0L7B8
G	-1	HIS	-	expression tag	UNP G0L7B8
G	0	GLY	-	expression tag	UNP G0L7B8
G	1	SER	-	expression tag	UNP G0L7B8
H	-7	MET	-	initiating methionine	UNP G0L7B8
H	-6	HIS	-	expression tag	UNP G0L7B8
H	-5	HIS	-	expression tag	UNP G0L7B8
H	-4	HIS	-	expression tag	UNP G0L7B8
H	-3	HIS	-	expression tag	UNP G0L7B8
H	-2	HIS	-	expression tag	UNP G0L7B8
H	-1	HIS	-	expression tag	UNP G0L7B8
H	0	GLY	-	expression tag	UNP G0L7B8
H	1	SER	-	expression tag	UNP G0L7B8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	H	1	Total Mg 1 1	0	0
2	G	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	13	Total O 13 13	0	0
3	B	9	Total O 9 9	0	0
3	C	12	Total O 12 12	0	0
3	D	8	Total O 8 8	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	9	Total 9	O 9	0	0
3	F	11	Total 11	O 11	0	0
3	G	19	Total 19	O 19	0	0
3	H	11	Total 11	O 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Galactonate dehydratase




#### • Molecule 1: Galactonate dehydratase



#### • Molecule 1: Galactonate dehydratase




Chain C: 




• Molecule 1: Galactonate dehydratase

Chain D: 




• Molecule 1: Galactonate dehydratase


Chain E: 




- Molecule 1: Galactonate dehydratase



- Molecule 1: Galactonate dehydratase



- Molecule 1: Galactonate dehydratase



D364	K369	Y373	K376	GLU	ILE	HIS	LYS	ASP	GLY	ALA	THR	LYS	ILE	ARG	SER
------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.84Å 154.07Å 150.87Å 90.00° 104.38° 90.00°	Depositor
Resolution (Å)	49.17 – 2.79 49.17 – 2.79	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.17-2.79) 99.1 (49.17-2.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.48 (at 2.81Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.231 , 0.266 0.263 , 0.299	Depositor DCC
$R_{free}$ test set	4656 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.3	Xtriage
Anisotropy	0.412	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 74.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	21906	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/2777	0.72	0/3766
1	B	0.52	0/2765	0.73	0/3752
1	C	0.50	0/2773	0.72	0/3760
1	D	0.51	0/2767	0.72	0/3753
1	E	0.50	0/2791	0.71	0/3781
1	F	0.51	0/2797	0.73	0/3789
1	G	0.51	0/2786	0.72	0/3777
1	H	0.52	0/2801	0.73	0/3793
All	All	0.51	0/22257	0.72	0/30171

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2721	0	2689	31	0
1	B	2709	0	2668	34	0
1	C	2717	0	2690	38	0
1	D	2711	0	2679	26	0
1	E	2735	0	2722	36	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2741	0	2740	37	0
1	G	2730	0	2709	35	0
1	H	2745	0	2751	35	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	13	0	0	0	0
3	B	9	0	0	0	0
3	C	12	0	0	1	0
3	D	8	0	0	0	0
3	E	9	0	0	2	0
3	F	11	0	0	0	0
3	G	19	0	0	0	0
3	H	11	0	0	0	0
All	All	21906	0	21648	257	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 257 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:333:THR:O	1:F:338:ARG:NH2	2.01	0.93
1:F:33:ILE:HD12	1:F:302:ILE:HD12	1.61	0.82
1:H:1:SER:HB3	1:H:38:LEU:HD13	1.62	0.81
1:H:169:LEU:HD22	1:H:173:GLN:HG2	1.64	0.78
1:H:136:LEU:HB3	1:H:166:LYS:HD3	1.68	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/396 (87%)	322 (94%)	21 (6%)	0	100	100
1	B	343/396 (87%)	322 (94%)	21 (6%)	0	100	100
1	C	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
1	D	343/396 (87%)	323 (94%)	20 (6%)	0	100	100
1	E	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
1	F	343/396 (87%)	318 (93%)	24 (7%)	1 (0%)	41	72
1	G	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
1	H	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
All	All	2744/3168 (87%)	2569 (94%)	174 (6%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	335	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/337 (84%)	250 (88%)	34 (12%)	5	15
1	B	281/337 (83%)	247 (88%)	34 (12%)	5	15
1	C	283/337 (84%)	248 (88%)	35 (12%)	4	14
1	D	282/337 (84%)	249 (88%)	33 (12%)	5	16
1	E	287/337 (85%)	254 (88%)	33 (12%)	5	17
1	F	289/337 (86%)	259 (90%)	30 (10%)	7	21
1	G	286/337 (85%)	250 (87%)	36 (13%)	4	14
1	H	290/337 (86%)	257 (89%)	33 (11%)	5	18
All	All	2282/2696 (85%)	2014 (88%)	268 (12%)	5	16

5 of 268 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	205	ILE
1	E	118	SER
1	H	88	TYR
1	D	256	ARG
1	E	1	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	C	95	GLN
1	F	158	GLN
1	G	95	GLN
1	H	95	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	351/396 (88%)	0.46	37 (10%) 6 3	38, 69, 101, 116	0
1	B	351/396 (88%)	0.47	28 (7%) 12 6	39, 68, 99, 117	0
1	C	351/396 (88%)	0.49	25 (7%) 16 9	38, 68, 103, 132	0
1	D	351/396 (88%)	0.31	12 (3%) 45 35	39, 69, 102, 118	0
1	E	351/396 (88%)	0.30	11 (3%) 49 39	35, 64, 94, 118	0
1	F	351/396 (88%)	0.36	17 (4%) 30 21	35, 63, 93, 123	0
1	G	351/396 (88%)	0.42	15 (4%) 35 25	38, 64, 94, 106	0
1	H	351/396 (88%)	0.39	17 (4%) 30 21	38, 64, 94, 108	0
All	All	2808/3168 (88%)	0.40	162 (5%) 23 15	35, 66, 99, 132	0

The worst 5 of 162 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	169	LEU	6.3
1	C	137	TYR	5.9
1	B	159	GLY	5.0
1	B	137	TYR	5.0
1	C	177	TYR	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	F	401	1/1	0.81	0.13	64,64,64,64	0
2	MG	H	401	1/1	0.81	0.11	73,73,73,73	0
2	MG	C	401	1/1	0.82	0.14	59,59,59,59	0
2	MG	D	401	1/1	0.90	0.15	100,100,100,100	0
2	MG	G	401	1/1	0.93	0.06	61,61,61,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.