



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2022 – 03:15 am GMT

PDB ID : 7OLJ
Title : Tankyrase 2 in complex with an inhibitor (OUL219)
Authors : Sowa, S.T.; Lehtio, L.
Deposited on : 2021-05-20
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.27
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	171	<div> <div>0%</div> <div>90%</div> <div>6%</div> </div>
1	BBB	171	<div> <div>2%</div> <div>91%</div> <div>5%</div> </div>
2	A	48	<div> <div>10%</div> <div>90%</div> <div>6%</div> </div>
2	B	48	<div> <div>10%</div> <div>85%</div> <div>12%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly [ADP-ribose] polymerase tankyrase-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	161	Total	C	N	O	S	0	2	0
			1308	821	242	237	8			
1	BBB	163	Total	C	N	O	S	0	3	0
			1317	827	243	239	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	944	SER	-	expression tag	UNP Q9H2K2
AAA	945	MET	-	expression tag	UNP Q9H2K2
BBB	944	SER	-	expression tag	UNP Q9H2K2
BBB	945	MET	-	expression tag	UNP Q9H2K2

- Molecule 2 is a protein called Poly [ADP-ribose] polymerase tankyrase-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	46	Total	C	N	O	S	0	1	0
			378	241	68	67	2			
2	B	47	Total	C	N	O	S	0	0	0
			373	238	65	68	2			

- Molecule 3 is 8-propan-2-yloxy-4 {H}-thieno[2,3-c]isoquinolin-5-one (three-letter code: VJN) (formula: C₁₄H₁₃NO₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	AAA	1	Total	C	N	O	S	0	0
			18	14	1	2	1		
3	BBB	1	Total	C	N	O	S	0	0
			18	14	1	2	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	Zn	0	0
			1	1		
4	BBB	1	Total	Zn	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	BBB	1	Total	O	S	0	0
			5	4	1		
5	BBB	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	AAA	141	Total	O	0	0
			141	141		
7	A	20	Total	O	0	0
			20	20		
7	BBB	110	Total	O	0	0
			110	110		
7	B	17	Total	O	0	0
			17	17		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Poly [ADP-ribose] polymerase tankyrase-2





- Molecule 1: Poly [ADP-ribose] polymerase tankyrase-2



- Molecule 2: Poly [ADP-ribose] polymerase tankyrase-2



- Molecule 2: Poly [ADP-ribose] polymerase tankyrase-2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	90.93Å 98.53Å 119.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.55 – 1.80 44.51 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.55-1.80) 99.8 (44.51-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.191 , 0.213 0.199 , 0.217	Depositor DCC
R_{free} test set	2493 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtrriage
Anisotropy	0.327	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3728	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, SO4, VJN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.65	0/1342	0.74	0/1799
1	BBB	0.65	0/1357	0.73	0/1820
2	A	0.62	0/389	0.73	0/529
2	B	0.62	0/385	0.69	0/525
All	All	0.64	0/3473	0.73	0/4673

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1308	0	1255	5	0
1	BBB	1317	0	1266	6	0
2	A	378	0	364	3	0
2	B	373	0	357	4	0
3	AAA	18	0	0	0	0
3	BBB	18	0	0	0	0
4	AAA	1	0	0	0	0
4	BBB	1	0	0	0	0
5	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	AAA	5	0	0	0	0
5	BBB	10	0	0	0	0
6	A	6	0	8	0	0
7	A	20	0	0	0	0
7	AAA	141	0	0	1	0
7	B	17	0	0	0	0
7	BBB	110	0	0	3	0
All	All	3728	0	3250	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:980:ARG:NH2	7:BBB:1303:HOH:O	2.39	0.54
1:AAA:1108[A]:LEU:HD11	2:A:1128[A]:ARG:HE	1.78	0.48
1:BBB:1055:PHE:HB2	2:B:1140:VAL:HG21	1.94	0.48
1:AAA:1054:MET:HG3	7:AAA:1419:HOH:O	2.13	0.48
1:BBB:1037:ASN:ND2	7:BBB:1303:HOH:O	2.38	0.48
1:AAA:1055:PHE:HB2	2:A:1140:VAL:HG21	1.96	0.48
1:BBB:972:MET:HG3	1:BBB:992:TYR:CE2	2.50	0.47
1:BBB:1095:GLN:HA	2:B:1153:ILE:O	2.16	0.45
2:B:1118:SER:HG	2:B:1142:TYR:HD2	1.65	0.45
1:AAA:1095:GLN:HA	2:A:1153:ILE:O	2.19	0.42
1:BBB:1054:MET:HG3	7:BBB:1393:HOH:O	2.20	0.41
1:BBB:1000:VAL:HB	2:B:1149:PRO:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	161/171 (94%)	160 (99%)	1 (1%)	0	100	100
1	BBB	164/171 (96%)	162 (99%)	2 (1%)	0	100	100
2	A	43/48 (90%)	42 (98%)	1 (2%)	0	100	100
2	B	45/48 (94%)	44 (98%)	1 (2%)	0	100	100
All	All	413/438 (94%)	408 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	139/146 (95%)	137 (99%)	2 (1%)	67	59
1	BBB	140/146 (96%)	140 (100%)	0	100	100
2	A	39/39 (100%)	39 (100%)	0	100	100
2	B	39/39 (100%)	38 (97%)	1 (3%)	46	32
All	All	357/370 (96%)	354 (99%)	3 (1%)	78	78

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	1004	LYS
1	AAA	1037	ASN
2	B	1128	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	1201	-	5,5,5	0.09	0	5,5,5	0.31	0
5	SO4	A	1202	-	4,4,4	0.39	0	6,6,6	0.10	0
3	VJN	AAA	1201	-	17,20,20	1.02	1 (5%)	22,29,29	2.13	2 (9%)
5	SO4	BBB	1203	-	4,4,4	0.39	0	6,6,6	0.14	0
3	VJN	BBB	1201	-	17,20,20	0.92	1 (5%)	22,29,29	2.09	3 (13%)
5	SO4	BBB	1204	-	4,4,4	0.40	0	6,6,6	0.06	0
5	SO4	AAA	1203	-	4,4,4	0.44	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	1201	-	-	3/4/4/4	-
3	VJN	BBB	1201	-	-	0/4/4/4	0/3/3/3
3	VJN	AAA	1201	-	-	0/4/4/4	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	1201	VJN	C8-N1	2.83	1.38	1.33
3	BBB	1201	VJN	C8-N1	2.65	1.37	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	1201	VJN	C7-C8-N1	-8.53	118.45	124.40
3	BBB	1201	VJN	C7-C8-N1	-8.23	118.66	124.40
3	AAA	1201	VJN	C9-N1-C8	3.52	122.13	116.88
3	BBB	1201	VJN	C9-N1-C8	3.28	121.78	116.88
3	BBB	1201	VJN	C4-O1-C2	2.63	122.73	119.37

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	1201	GOL	C1-C2-C3-O3
6	A	1201	GOL	O2-C2-C3-O3
6	A	1201	GOL	O1-C1-C2-O2

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand VJN AAA 1201



Ligand VJN BBB 1201



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	161/171 (94%)	-0.04	2 (1%) 79 76	24, 33, 46, 69	0
1	BBB	163/171 (95%)	0.07	4 (2%) 57 52	26, 35, 50, 73	0
2	A	46/48 (95%)	0.67	5 (10%) 5 4	26, 33, 59, 74	0
2	B	47/48 (97%)	0.61	5 (10%) 6 4	27, 38, 71, 80	0
All	All	417/438 (95%)	0.15	16 (3%) 40 35	24, 34, 59, 80	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	1132	ASN	6.8
2	A	1131	VAL	6.0
1	BBB	951	SER	4.1
2	B	1119	PRO	3.7
1	BBB	950	GLY	3.6
2	B	1117	HIS	3.4
1	BBB	1112	ALA	3.1
2	B	1116	ALA	3.1
1	AAA	951	SER	2.8
1	AAA	950	GLY	2.6
2	A	1161	GLU	2.5
2	B	1120	PRO	2.5
2	B	1121	GLY	2.3
1	BBB	1057	ALA	2.3
2	A	1117	HIS	2.2
2	A	1120	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	GOL	A	1201	6/6	0.71	0.33	43,48,49,50	6
3	VJN	BBB	1201	18/18	0.92	0.12	29,32,38,39	0
5	SO4	A	1202	5/5	0.96	0.21	63,64,64,67	0
3	VJN	AAA	1201	18/18	0.96	0.09	27,29,33,34	0
5	SO4	BBB	1204	5/5	0.97	0.20	59,59,60,62	0
4	ZN	AAA	1202	1/1	0.98	0.06	36,36,36,36	0
5	SO4	BBB	1203	5/5	0.98	0.10	36,40,42,43	0
4	ZN	BBB	1202	1/1	0.98	0.06	33,33,33,33	0
5	SO4	AAA	1203	5/5	0.98	0.09	33,38,38,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around VJN BBB 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around VJN AAA 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.