



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2020 – 09:29 AM BST

PDB ID : 6OM2
Title : Crystal structure of atypical integrin alphaV beta8 with proTGF-beta1 ligand peptide
Authors : Wang, J.C.; Springer, T.A.
Deposited on : 2019-04-17
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	598	<div> <div>6%</div> <div>92%</div> <div>8%</div> </div>
1	C	598	<div> <div>4%</div> <div>88%</div> <div>11%</div> </div>
2	B	421	<div> <div>26%</div> <div>63%</div> <div>10%</div> <div>26%</div> </div>
2	D	421	<div> <div>22%</div> <div>66%</div> <div>12%</div> <div>22%</div> </div>
3	E	11	<div> <div>36%</div> <div>55%</div> <div>36%</div> <div>9%</div> </div>
3	F	11	<div> <div>27%</div> <div>64%</div> <div>9%</div> <div>9%</div> <div>18%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	G	3	 100%
4	K	3	 67% 33%
4	S	3	 100%
5	H	7	 14% 86%
5	L	7	 29% 43% 29%
5	O	7	 29% 71%
6	I	5	 40% 40% 20%
7	J	4	 25% 25% 50%
8	M	2	 50% 50%
8	P	2	 100%
8	Q	2	 100%
8	R	2	 50% 50%
9	N	6	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
14	GLY	C	601	-	-	-	X
4	NAG	K	1	-	-	-	X
4	NAG	K	2	-	-	-	X
4	NAG	S	2	-	-	-	X
4	BMA	S	3	-	-	-	X
5	MAN	O	6	-	-	-	X
8	NAG	M	1	-	-	X	-
8	NAG	M	2	-	-	X	X
8	NAG	Q	2	-	-	-	X

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 28790 atoms, of which 13455 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-V.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	598	Total	C	H	N	O	S	0	0	0
			8991	2927	4371	785	887	21			
1	C	596	Total	C	H	N	O	S	0	0	0
			8981	2923	4369	783	885	21			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	400	GLY	-	insertion	UNP P06756
A	401	CYS	MET	engineered mutation	UNP P06756
A	596	THR	-	expression tag	UNP P06756
A	597	GLY	-	expression tag	UNP P06756
A	598	GLY	-	expression tag	UNP P06756
C	400	GLY	-	insertion	UNP P06756
C	401	CYS	MET	engineered mutation	UNP P06756
C	596	THR	-	expression tag	UNP P06756
C	597	GLY	-	expression tag	UNP P06756
C	598	GLY	-	expression tag	UNP P06756

- Molecule 2 is a protein called Integrin beta-8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	310	Total	C	H	N	O	S	0	0	0
			4747	1551	2301	419	460	16			
2	D	330	Total	C	H	N	O	S	0	0	0
			5003	1638	2414	446	489	16			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	259	CYS	VAL	engineered mutation	UNP P26012
D	259	CYS	VAL	engineered mutation	UNP P26012

- Molecule 3 is a protein called proTGF-beta1 RGD peptide.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	10	Total	C	N	O	0	0	1
			68	40	16	12			
3	F	9	Total	C	N	O	0	0	1
			64	38	15	11			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	K	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	S	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	7	Total	C	N	O	0	0	0
			83	46	2	35			
5	L	7	Total	C	N	O	0	0	0
			83	46	2	35			
5	O	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	I	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	J	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 8 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	M	2	Total	C	N	O	0	0	0
			28	16	2	10			
8	P	2	Total	C	N	O	0	0	0
			28	16	2	10			
8	Q	2	Total	C	N	O	0	0	0
			28	16	2	10			
8	R	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	N	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 10 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	1	Total	Ca	0	0
			1	1		
10	A	4	Total	Ca	0	0
			4	4		
10	D	1	Total	Ca	0	0
			1	1		
10	C	4	Total	Ca	0	0
			4	4		

- Molecule 11 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	N	O	0	0
			14	8	1	5		
11	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 12 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	C	O	0	0
			7	4	3		
12	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 13 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	B	1	Total	Mg	0	0
			1	1		
13	D	1	Total	Mg	0	0
			1	1		

- Molecule 14 is GLYCINE (three-letter code: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
14	C	1	Total	C	N	O	0	0
			4	2	1	1		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	75	Total	O	0	0
			75	75		
15	B	11	Total	O	0	0
			11	11		
15	C	97	Total	O	0	0
			97	97		
15	D	19	Total	O	0	0
			19	19		
15	E	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrin alpha-V




- Molecule 1: Integrin alpha-V




- Molecule 2: Integrin beta-8






• Molecule 2: Integrin beta-8



• Molecule 3: proTGF-beta1 RGD peptide



• Molecule 3: proTGF-beta1 RGD peptide



• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  67% 33%

MAG1
MAG2
BMA3

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain S:  100%

MAG1
MAG2
BMA3

- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  14% 86%

MAG1
MAG2
BMA3
MAN4
MAN5
MAN6
MAN7

- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  29% 43% 29%

MAG1
MAG2
BMA3
MAN4
MAN5
MAN6
MAN7

- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:  29% 71%

MAG1
MAG2
BMA3
MAN4
MAN5
MAN6
MAN7

- Molecule 6: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  40% 40% 20%



- Molecule 7: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 9: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	161.19Å 53.85Å 176.62Å 90.00° 111.47° 90.00°	Depositor
Resolution (Å)	47.43 – 2.77 47.43 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.5 (47.43-2.77) 97.0 (47.43-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.251 , 0.282 0.251 , 0.280	Depositor DCC
R_{free} test set	2000 reflections (2.62%)	wwPDB-VP
Wilson B-factor (Å ²)	75.3	Xtriage
Anisotropy	0.395	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	28790	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BMA, NAG, CA, PEG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/4724	0.46	0/6395
1	C	0.27	0/4716	0.48	1/6385 (0.0%)
2	B	0.24	0/2489	0.45	0/3361
2	D	0.25	0/2639	0.44	0/3571
3	E	0.20	0/67	0.41	0/88
3	F	0.20	0/63	0.40	0/83
All	All	0.26	0/14698	0.46	1/19883 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	303	ARG	NE-CZ-NH2	5.81	123.20	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4620	4371	4468	34	0
1	C	4612	4369	4462	48	0
2	B	2446	2301	2412	30	0
2	D	2589	2414	2554	37	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	68	0	70	3	0
3	F	64	0	67	1	0
4	G	39	0	34	2	0
4	K	39	0	34	0	0
4	S	39	0	34	5	0
5	H	83	0	70	1	0
5	L	83	0	70	3	0
5	O	83	0	70	7	0
6	I	61	0	52	4	0
7	J	50	0	43	4	0
8	M	28	0	25	11	0
8	P	28	0	25	0	0
8	Q	28	0	25	2	0
8	R	28	0	25	5	0
9	N	72	0	61	0	0
10	A	4	0	0	0	0
10	B	1	0	0	0	0
10	C	4	0	0	0	0
10	D	1	0	0	0	0
11	A	28	0	26	0	0
11	D	14	0	13	1	0
12	A	7	0	10	0	0
12	C	7	0	10	1	0
13	B	1	0	0	0	0
13	D	1	0	0	0	0
14	C	4	0	2	1	0
15	A	75	0	0	6	0
15	B	11	0	0	0	0
15	C	97	0	0	3	0
15	D	19	0	0	2	0
15	E	1	0	0	0	0
All	All	15335	13455	14662	179	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:233:GLU:OE1	8:R:2:NAG:O4	1.88	0.89
1:A:121:GLU:OE1	2:B:164:ARG:NH2	2.12	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:289:ASP:OD2	5:H:2:NAG:H5	1.80	0.81
8:M:1:NAG:H62	8:M:2:NAG:N2	1.96	0.80
5:O:3:BMA:H2	5:O:7:MAN:H2	1.66	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	596/598 (100%)	551 (92%)	45 (8%)	0	100	100
1	C	594/598 (99%)	562 (95%)	31 (5%)	1 (0%)	47	76
2	B	292/421 (69%)	264 (90%)	28 (10%)	0	100	100
2	D	320/421 (76%)	295 (92%)	23 (7%)	2 (1%)	25	54
3	E	8/11 (73%)	7 (88%)	1 (12%)	0	100	100
3	F	7/11 (64%)	7 (100%)	0	0	100	100
All	All	1817/2060 (88%)	1686 (93%)	128 (7%)	3 (0%)	47	76

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	388	PHE
1	C	303	ARG
2	D	368	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	488/488 (100%)	482 (99%)	6 (1%)	71	90
1	C	488/488 (100%)	477 (98%)	11 (2%)	50	79
2	B	277/369 (75%)	276 (100%)	1 (0%)	91	96
2	D	291/369 (79%)	290 (100%)	1 (0%)	92	97
3	E	6/7 (86%)	6 (100%)	0	100	100
3	F	6/7 (86%)	5 (83%)	1 (17%)	2	6
All	All	1556/1728 (90%)	1536 (99%)	20 (1%)	69	89

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	114	TRP
1	C	155	CYS
1	C	586	ASN
1	C	73	ASP
1	C	100	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

53 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	1	1,4	14,14,15	0.36	0	17,19,21	0.45	0
4	NAG	G	2	4	14,14,15	0.41	0	17,19,21	0.50	0
4	BMA	G	3	4	11,11,12	0.73	0	15,15,17	0.87	0
5	NAG	H	1	1,5	14,14,15	0.19	0	17,19,21	0.48	0
5	NAG	H	2	5	14,14,15	0.54	0	17,19,21	0.45	0
5	BMA	H	3	5	11,11,12	0.82	0	15,15,17	1.00	1 (6%)
5	MAN	H	4	5	11,11,12	1.18	1 (9%)	15,15,17	0.98	1 (6%)
5	MAN	H	5	5	11,11,12	0.60	0	15,15,17	1.05	2 (13%)
5	MAN	H	6	5	11,11,12	1.35	1 (9%)	15,15,17	1.11	1 (6%)
5	MAN	H	7	5	11,11,12	0.94	1 (9%)	15,15,17	0.87	1 (6%)
6	NAG	I	1	1,6	14,14,15	0.48	0	17,19,21	0.49	0
6	NAG	I	2	6	14,14,15	0.19	0	17,19,21	0.50	0
6	BMA	I	3	6	11,11,12	0.74	0	15,15,17	0.84	0
6	MAN	I	4	6	11,11,12	0.71	0	15,15,17	1.02	2 (13%)
6	MAN	I	5	6	11,11,12	0.68	0	15,15,17	1.00	2 (13%)
7	NAG	J	1	1,7	14,14,15	0.18	0	17,19,21	0.42	0
7	NAG	J	2	7	14,14,15	0.25	0	17,19,21	0.39	0
7	BMA	J	3	7	11,11,12	1.03	1 (9%)	15,15,17	1.46	1 (6%)
7	MAN	J	4	7	11,11,12	0.78	0	15,15,17	1.39	3 (20%)
4	NAG	K	1	2,4	14,14,15	0.74	1 (7%)	17,19,21	1.11	2 (11%)
4	NAG	K	2	4	14,14,15	0.30	0	17,19,21	0.36	0
4	BMA	K	3	4	11,11,12	0.56	0	15,15,17	0.95	0
5	NAG	L	1	1,5	14,14,15	0.45	0	17,19,21	0.43	0
5	NAG	L	2	5	14,14,15	0.33	0	17,19,21	0.40	0
5	BMA	L	3	5	11,11,12	0.70	0	15,15,17	0.89	0
5	MAN	L	4	5	11,11,12	0.94	1 (9%)	15,15,17	1.56	3 (20%)
5	MAN	L	5	5	11,11,12	0.89	1 (9%)	15,15,17	1.67	4 (26%)
5	MAN	L	6	5	11,11,12	0.84	0	15,15,17	1.27	3 (20%)
5	MAN	L	7	5	11,11,12	0.63	0	15,15,17	1.12	2 (13%)
8	NAG	M	1	1,8	14,14,15	0.44	0	17,19,21	0.41	0
8	NAG	M	2	8	14,14,15	0.49	0	17,19,21	0.63	1 (5%)
9	NAG	N	1	1,9	14,14,15	0.32	0	17,19,21	0.42	0
9	NAG	N	2	9	14,14,15	0.19	0	17,19,21	0.46	0
9	BMA	N	3	9	11,11,12	0.68	0	15,15,17	0.82	0
9	MAN	N	4	9	11,11,12	0.87	1 (9%)	15,15,17	0.90	1 (6%)
9	MAN	N	5	9	11,11,12	0.73	0	15,15,17	1.06	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	MAN	N	6	9	11,11,12	0.75	1 (9%)	15,15,17	0.95	2 (13%)
5	NAG	O	1	1,5	14,14,15	0.18	0	17,19,21	0.47	0
5	NAG	O	2	5	14,14,15	0.28	0	17,19,21	0.51	0
5	BMA	O	3	5	11,11,12	1.10	0	15,15,17	1.25	2 (13%)
5	MAN	O	4	5	11,11,12	0.89	1 (9%)	15,15,17	1.39	3 (20%)
5	MAN	O	5	5	11,11,12	0.74	0	15,15,17	1.61	3 (20%)
5	MAN	O	6	5	11,11,12	0.76	1 (9%)	15,15,17	1.49	3 (20%)
5	MAN	O	7	5	11,11,12	0.75	0	15,15,17	1.16	2 (13%)
8	NAG	P	1	1,8	14,14,15	0.42	0	17,19,21	0.51	0
8	NAG	P	2	8	14,14,15	0.35	0	17,19,21	0.52	0
8	NAG	Q	1	1,8	14,14,15	0.49	0	17,19,21	0.44	0
8	NAG	Q	2	8	14,14,15	0.23	0	17,19,21	0.43	0
8	NAG	R	1	8,2	14,14,15	0.65	0	17,19,21	0.69	1 (5%)
8	NAG	R	2	8	14,14,15	0.25	0	17,19,21	0.66	0
4	NAG	S	1	2,4	14,14,15	0.30	0	17,19,21	0.52	0
4	NAG	S	2	4	14,14,15	0.25	0	17,19,21	0.35	0
4	BMA	S	3	4	11,11,12	0.60	0	15,15,17	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
5	NAG	H	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1
5	BMA	H	3	5	-	1/2/19/22	0/1/1/1
5	MAN	H	4	5	-	2/2/19/22	0/1/1/1
5	MAN	H	5	5	-	2/2/19/22	0/1/1/1
5	MAN	H	6	5	-	0/2/19/22	0/1/1/1
5	MAN	H	7	5	-	0/2/19/22	0/1/1/1
6	NAG	I	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	I	2	6	-	0/6/23/26	0/1/1/1
6	BMA	I	3	6	-	0/2/19/22	0/1/1/1
6	MAN	I	4	6	-	1/2/19/22	0/1/1/1
6	MAN	I	5	6	-	1/2/19/22	0/1/1/1
7	NAG	J	1	1,7	-	1/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	J	2	7	-	2/6/23/26	0/1/1/1
7	BMA	J	3	7	-	0/2/19/22	0/1/1/1
7	MAN	J	4	7	-	0/2/19/22	1/1/1/1
4	NAG	K	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	K	2	4	-	2/6/23/26	0/1/1/1
4	BMA	K	3	4	-	1/2/19/22	0/1/1/1
5	NAG	L	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	L	2	5	-	2/6/23/26	0/1/1/1
5	BMA	L	3	5	-	2/2/19/22	0/1/1/1
5	MAN	L	4	5	-	0/2/19/22	0/1/1/1
5	MAN	L	5	5	-	0/2/19/22	0/1/1/1
5	MAN	L	6	5	-	1/2/19/22	0/1/1/1
5	MAN	L	7	5	-	2/2/19/22	0/1/1/1
8	NAG	M	1	1,8	-	2/6/23/26	0/1/1/1
8	NAG	M	2	8	-	4/6/23/26	0/1/1/1
9	NAG	N	1	1,9	-	2/6/23/26	0/1/1/1
9	NAG	N	2	9	-	2/6/23/26	0/1/1/1
9	BMA	N	3	9	-	0/2/19/22	0/1/1/1
9	MAN	N	4	9	-	0/2/19/22	0/1/1/1
9	MAN	N	5	9	-	0/2/19/22	0/1/1/1
9	MAN	N	6	9	-	0/2/19/22	0/1/1/1
5	NAG	O	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	O	2	5	-	0/6/23/26	0/1/1/1
5	BMA	O	3	5	-	2/2/19/22	0/1/1/1
5	MAN	O	4	5	-	0/2/19/22	0/1/1/1
5	MAN	O	5	5	-	1/2/19/22	0/1/1/1
5	MAN	O	6	5	-	0/2/19/22	0/1/1/1
5	MAN	O	7	5	-	0/2/19/22	1/1/1/1
8	NAG	P	1	1,8	-	2/6/23/26	0/1/1/1
8	NAG	P	2	8	-	2/6/23/26	0/1/1/1
8	NAG	Q	1	1,8	-	2/6/23/26	0/1/1/1
8	NAG	Q	2	8	-	2/6/23/26	0/1/1/1
8	NAG	R	1	8,2	-	3/6/23/26	0/1/1/1
8	NAG	R	2	8	-	4/6/23/26	0/1/1/1
4	NAG	S	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	S	2	4	-	3/6/23/26	0/1/1/1
4	BMA	S	3	4	-	2/2/19/22	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	6	MAN	O5-C1	-3.76	1.37	1.43
5	H	4	MAN	O5-C1	-3.52	1.38	1.43
5	H	7	MAN	O5-C1	-2.61	1.39	1.43
5	L	5	MAN	C1-C2	2.45	1.57	1.52
7	J	3	BMA	O3-C3	2.42	1.48	1.43

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	O	5	MAN	C1-O5-C5	4.18	117.86	112.19
7	J	3	BMA	O3-C3-C2	3.99	117.64	109.99
7	J	4	MAN	C1-O5-C5	3.64	117.13	112.19
5	L	5	MAN	C1-O5-C5	3.54	116.98	112.19
5	L	4	MAN	C1-O5-C5	3.45	116.86	112.19

There are no chirality outliers.

5 of 67 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	H	4	MAN	C4-C5-C6-O6
8	M	2	NAG	C1-C2-N2-C7
8	R	2	NAG	C1-C2-N2-C7
8	P	2	NAG	O5-C5-C6-O6
9	N	1	NAG	O5-C5-C6-O6

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	J	4	MAN	C1-C2-C3-C4-C5-O5
5	O	7	MAN	C1-C2-C3-C4-C5-O5

28 monomers are involved in 44 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	4	MAN	2	0
5	H	2	NAG	1	0
4	G	1	NAG	1	0
6	I	3	BMA	2	0
4	S	1	NAG	3	0
4	S	2	NAG	4	0
5	O	5	MAN	2	0


Continued on next page...





Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	I	5	MAN	2	0
8	M	1	NAG	7	0
5	O	6	MAN	1	0
5	O	7	MAN	3	0
7	J	4	MAN	3	0
5	O	4	MAN	3	0
8	Q	2	NAG	2	0
4	G	2	NAG	2	0
7	J	1	NAG	1	0
5	L	1	NAG	1	0
8	Q	1	NAG	2	0
4	G	3	BMA	1	0
7	J	3	BMA	2	0
5	L	6	MAN	2	0
5	O	3	BMA	2	0
5	O	2	NAG	1	0
4	S	3	BMA	2	0
8	M	2	NAG	9	0
5	O	1	NAG	1	0
8	R	1	NAG	4	0
8	R	2	NAG	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.








Oligosaccharide Chain H**Bond lengths****Bond angles****Torsions****Rings**




















5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	NAG	A	2025	1	14,14,15	0.24	0	17,19,21	0.44	0
12	PEG	C	632	-	6,6,6	0.50	0	5,5,5	0.32	0
11	NAG	D	3201	2	14,14,15	0.25	0	17,19,21	0.33	0
12	PEG	A	2026	-	6,6,6	0.49	0	5,5,5	0.24	0
11	NAG	A	2024	1	14,14,15	0.26	0	17,19,21	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	NAG	A	2025	1	-	2/6/23/26	0/1/1/1
12	PEG	C	632	-	-	1/4/4/4	-
11	NAG	D	3201	2	-	4/6/23/26	0/1/1/1
12	PEG	A	2026	-	-	1/4/4/4	-
11	NAG	A	2024	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	D	3201	NAG	C1-C2-N2-C7
11	A	2024	NAG	O5-C5-C6-O6
11	A	2025	NAG	C4-C5-C6-O6
11	A	2024	NAG	C4-C5-C6-O6
11	A	2025	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	C	632	PEG	1	0
11	D	3201	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	598/598 (100%)	0.74	34 (5%) 23 18	59, 93, 156, 225	0
1	C	596/598 (99%)	0.64	21 (3%) 44 38	58, 87, 133, 223	0
2	B	310/421 (73%)	1.85	109 (35%) 0 0	70, 159, 223, 275	0
2	D	330/421 (78%)	1.49	92 (27%) 0 0	70, 138, 201, 261	0
3	E	10/11 (90%)	2.15	4 (40%) 0 0	102, 118, 135, 135	0
3	F	9/11 (81%)	1.48	3 (33%) 0 0	102, 115, 151, 153	0
All	All	1853/2060 (89%)	1.04	263 (14%) 2 1	58, 103, 196, 275	0

The worst 5 of 263 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	358	TYR	13.7
2	D	354	VAL	13.4
2	B	137	PHE	11.3
2	B	388	PHE	11.2
2	B	361	ILE	10.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BMA	K	3	11/12	0.37	0.37	207,209,211,211	0
4	BMA	S	3	11/12	0.42	0.48	182,186,187,188	0
8	NAG	M	2	14/15	0.51	0.49	155,156,158,159	0
5	MAN	L	4	11/12	0.55	0.29	151,152,153,154	0
5	MAN	L	5	11/12	0.63	0.28	143,151,154,154	0
4	NAG	K	2	14/15	0.64	0.51	215,219,221,223	0
5	MAN	O	6	11/12	0.67	0.41	132,138,140,141	0
8	NAG	M	1	14/15	0.69	0.34	122,133,145,152	0
4	NAG	S	2	14/15	0.71	0.41	192,197,201,203	0
5	MAN	O	7	11/12	0.71	0.26	117,123,125,126	0
8	NAG	R	2	14/15	0.72	0.25	199,201,204,206	0
8	NAG	Q	1	14/15	0.74	0.37	136,157,164,171	0
4	NAG	S	1	14/15	0.75	0.27	184,201,212,214	0
8	NAG	Q	2	14/15	0.77	0.64	175,178,180,180	0
5	MAN	O	5	11/12	0.78	0.24	123,124,125,125	0
5	MAN	H	6	11/12	0.78	0.29	83,90,94,95	0
8	NAG	R	1	14/15	0.79	0.23	206,211,213,214	0
6	BMA	I	3	11/12	0.80	0.14	100,102,105,109	0
5	MAN	O	4	11/12	0.80	0.24	121,122,124,127	0
4	NAG	K	1	14/15	0.80	0.57	215,227,231,231	0
4	BMA	G	3	11/12	0.81	0.14	135,136,138,139	0
5	BMA	O	3	11/12	0.81	0.21	109,112,120,120	0
5	MAN	L	7	11/12	0.81	0.21	152,153,154,154	0
7	BMA	J	3	11/12	0.82	0.22	115,116,118,118	0
9	MAN	N	4	11/12	0.83	0.14	107,111,115,118	0
6	NAG	I	2	14/15	0.83	0.21	91,95,97,99	0
6	MAN	I	4	11/12	0.84	0.23	112,114,115,117	0
7	MAN	J	4	11/12	0.85	0.20	113,116,120,120	0
8	NAG	P	2	14/15	0.86	0.27	109,110,111,113	0
5	BMA	H	3	11/12	0.87	0.14	85,87,91,93	0
9	MAN	N	5	11/12	0.87	0.22	121,124,126,127	0
5	NAG	O	2	14/15	0.87	0.20	109,114,117,118	0
5	NAG	L	2	14/15	0.87	0.23	126,129,135,139	0
7	NAG	J	1	14/15	0.88	0.19	101,111,115,119	0
8	NAG	P	1	14/15	0.88	0.20	99,105,110,111	0
5	NAG	H	2	14/15	0.88	0.16	85,88,90,91	0
4	NAG	G	2	14/15	0.88	0.19	133,135,137,137	0
5	MAN	H	7	11/12	0.88	0.23	95,96,97,97	0
5	MAN	L	6	11/12	0.89	0.27	151,152,153,155	0
5	NAG	H	1	14/15	0.89	0.19	82,91,96,99	0
5	BMA	L	3	11/12	0.89	0.15	143,145,150,151	0
4	NAG	G	1	14/15	0.89	0.25	122,127,127,132	0
5	MAN	H	4	11/12	0.90	0.21	80,84,87,88	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	NAG	N	1	14/15	0.90	0.20	75,80,83,88	0
6	MAN	I	5	11/12	0.91	0.14	94,97,100,100	0
6	NAG	I	1	14/15	0.91	0.24	76,84,90,97	0
9	BMA	N	3	11/12	0.92	0.11	97,103,107,108	0
9	MAN	N	6	11/12	0.92	0.12	108,109,111,111	0
7	NAG	J	2	14/15	0.92	0.18	111,113,114,115	0
5	MAN	H	5	11/12	0.92	0.22	88,90,94,96	0
5	NAG	O	1	14/15	0.92	0.21	83,99,107,108	0
9	NAG	N	2	14/15	0.92	0.16	82,90,94,95	0
5	NAG	L	1	14/15	0.92	0.25	110,116,119,122	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around Chain S:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain L:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain O:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain P:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain Q:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain R:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
14	GLY	C	601	4/5	0.45	0.42	102,102,103,104	0
13	MG	D	3208	1/1	0.55	0.14	106,106,106,106	0
11	NAG	A	2024	14/15	0.63	0.27	125,132,136,137	0
11	NAG	D	3201	14/15	0.86	0.39	189,202,212,212	0
13	MG	B	2001	1/1	0.86	0.09	103,103,103,103	0
12	PEG	C	632	7/7	0.86	0.23	75,77,78,79	0
10	CA	B	2002	1/1	0.87	0.15	94,94,94,94	0
12	PEG	A	2026	7/7	0.87	0.16	84,87,87,88	0
10	CA	D	3204	1/1	0.92	0.15	92,92,92,92	0
11	NAG	A	2025	14/15	0.92	0.21	104,122,126,128	0
10	CA	A	2003	1/1	0.93	0.20	73,73,73,73	0
10	CA	C	603	1/1	0.95	0.13	61,61,61,61	0
10	CA	C	602	1/1	0.96	0.14	73,73,73,73	0
10	CA	A	2001	1/1	0.96	0.17	69,69,69,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
10	CA	C	605	1/1	0.96	0.20	82,82,82,82	0
10	CA	A	2002	1/1	0.97	0.11	70,70,70,70	0
10	CA	C	604	1/1	0.97	0.12	66,66,66,66	0
10	CA	A	2004	1/1	0.97	0.18	78,78,78,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.