



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 08:30 AM BST

PDB ID : 1ON8  
Title : Crystal structure of mouse alpha-1,4-N-acetylhexosaminyltransferase (EXTL2) with UDP and GlcUAb(1-3)Galb(1-O)-naphthalenelmethanol an acceptor substrate analog  
Authors : Pedersen, L.C.; Dong, J.; Taniguchi, F.; Kitagawa, H.; Krahn, J.M.; Pedersen, L.G.; Sugahara, K.; Negishi, M.  
Deposited on : 2003-02-27  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	<div> <div>2%</div> <div> <div></div> <div>64%</div> <div>25%</div> <div>•</div> <div>10%</div> </div> </div>
1	B	293	<div> <div>2%</div> <div> <div></div> <div>61%</div> <div>27%</div> <div>•</div> <div>11%</div> </div> </div>
2	C	2	<div> <div></div> <div>100%</div> </div>
2	D	2	<div> <div></div> <div>100%</div> </div>

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-1,4-N-acetylhexosaminyltransferase EXTL2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	0	0
			2110	1362	354	383	11			
1	B	262	Total	C	N	O	S	0	0	0
			2078	1342	348	377	11			

- Molecule 2 is an oligosaccharide called beta-D-glucopyranuronic acid-(1-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			24	12	12			
2	D	2	Total	C	O	0	0	0
			24	12	12			

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mn	0	0
			1	1		
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>O<sub>12</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
4	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	53	Total 53	O 53	0	0
6	B	44	Total 44	O 44	0	0



GAL1  
BDP2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.16 Å   125.16 Å   83.69 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	19.98 – 2.70 19.98 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.7 (19.98-2.70) 96.8 (19.98-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 2.71 Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.192   ,   0.254 0.186   ,   0.248	Depositor DCC
$R_{free}$ test set	901 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.4	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4393	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, UDP, MN, EDO, BDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2169	0.63	0/2949
1	B	0.36	0/2135	0.60	0/2906
All	All	0.37	0/4304	0.61	0/5855

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2110	0	2051	57	0
1	B	2078	0	2015	63	0
2	C	24	0	17	0	0
2	D	24	0	17	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	25	0	11	0	0
4	B	25	0	11	0	0
5	A	4	0	6	0	0
5	B	4	0	6	0	0
6	A	53	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	44	0	0	4	0
All	All	4393	0	4134	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:GLU:O	1:A:292:GLN:HB2	1.75	0.86
1:B:71:MET:CE	1:B:80:LEU:HD12	2.19	0.72
1:A:318:GLN:HE22	1:B:205:ASN:HB3	1.56	0.71
1:B:238:ILE:HD12	1:B:245:ASP:HA	1.71	0.71
1:A:214:ILE:HG12	1:A:246:ASP:HB3	1.71	0.70
1:A:79:LEU:HD22	1:A:277:THR:HB	1.72	0.70
1:B:226:LEU:HD23	1:B:252:LEU:HD21	1.73	0.70
1:B:64:LEU:O	1:B:64:LEU:HD23	1.92	0.70
1:B:146:ALA:HB2	1:B:168:TRP:CZ3	2.27	0.69
1:B:71:MET:HE3	1:B:80:LEU:HD12	1.76	0.67
1:B:272:ASN:OD1	1:B:274:GLU:HB2	1.94	0.66
1:A:71:MET:CE	1:A:80:LEU:HD12	2.26	0.65
1:B:133:ARG:HG2	6:B:636:HOH:O	1.97	0.65
1:A:147:VAL:HG23	1:A:221:SER:HB3	1.77	0.65
1:B:133:ARG:HD3	1:B:239:ASP:OD1	1.96	0.64
1:B:64:LEU:HD12	1:B:145:ASN:ND2	2.13	0.64
1:A:199:GLN:HE21	1:A:200:THR:HG22	1.64	0.63
1:A:201:PRO:HG3	1:A:311:TYR:CE1	2.34	0.63
1:B:63:ALA:C	1:B:65:ASP:H	2.02	0.62
1:B:146:ALA:HB2	1:B:168:TRP:CH2	2.34	0.62
1:B:182:LYS:HD2	6:B:640:HOH:O	1.98	0.62
1:A:307:MET:HE1	1:A:309:LEU:HB2	1.81	0.61
1:B:90:VAL:HG21	1:B:161:LEU:HD12	1.82	0.61
1:A:238:ILE:HD12	1:A:245:ASP:HA	1.82	0.61
1:A:182:LYS:HG2	1:A:183:HIS:N	2.15	0.60
1:B:87:TYR:O	1:B:93:LEU:HD22	2.02	0.60
1:A:146:ALA:HB2	1:A:168:TRP:CH2	2.37	0.59
1:B:307:MET:HE1	1:B:309:LEU:HB2	1.84	0.59
1:A:95:LYS:HD3	1:A:123:ILE:HD12	1.85	0.58
1:B:243:ASN:O	1:B:244:CYS:HB2	2.03	0.58
1:B:197:GLU:H	1:B:197:GLU:CD	2.07	0.58
1:A:120:ILE:HD13	1:A:120:ILE:H	1.69	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:LYS:HE2	1:A:201:PRO:HD3	1.88	0.56
1:A:146:ALA:HB2	1:A:168:TRP:CZ3	2.41	0.56
1:B:120:ILE:H	1:B:120:ILE:HD13	1.70	0.55
1:B:71:MET:HE2	1:B:80:LEU:HD12	1.88	0.55
1:A:91:PRO:O	1:A:92:SER:HB2	2.08	0.54
1:A:274:GLU:O	1:A:277:THR:HG22	2.09	0.53
1:B:292:GLN:O	1:B:295:TYR:HB3	2.09	0.53
1:A:306:GLY:O	1:A:308:PRO:HD3	2.09	0.53
1:B:133:ARG:NH1	1:B:235:HIS:HB3	2.24	0.53
1:A:168:TRP:HA	1:A:176:ILE:HG13	1.91	0.52
1:B:214:ILE:HG12	1:B:246:ASP:HB3	1.91	0.52
1:B:73:THR:HB	1:B:80:LEU:HD13	1.90	0.52
1:A:201:PRO:HG2	1:A:202:GLY:N	2.25	0.52
1:B:243:ASN:OD1	1:B:244:CYS:N	2.43	0.51
1:A:71:MET:HE3	1:A:80:LEU:HD12	1.92	0.51
1:B:230:GLN:NE2	1:B:304:TYR:OH	2.42	0.51
1:B:272:ASN:OD1	1:B:274:GLU:CB	2.59	0.51
1:B:284:TRP:HA	1:B:289:HIS:CG	2.46	0.51
1:A:137:GLN:HA	1:A:228:GLN:NE2	2.25	0.50
1:A:95:LYS:HD3	1:A:123:ILE:CD1	2.42	0.49
1:A:252:LEU:C	1:A:252:LEU:HD23	2.32	0.49
1:A:252:LEU:HD23	1:A:253:VAL:N	2.27	0.49
1:B:112:TRP:CZ2	1:B:122:VAL:HG21	2.48	0.49
1:A:243:ASN:OD1	1:A:244:CYS:N	2.38	0.49
1:B:79:LEU:HD22	1:B:277:THR:HB	1.94	0.49
1:A:171:PHE:N	1:A:172:PRO:HD3	2.27	0.48
1:B:194:GLY:HA3	1:B:198:LEU:HD12	1.95	0.48
1:B:63:ALA:C	1:B:65:ASP:N	2.64	0.48
1:B:318:GLN:HB3	1:B:321:PHE:HB3	1.95	0.48
1:B:191:TYR:CZ	1:B:307:MET:HG3	2.49	0.47
1:A:86:HIS:CD2	1:A:155:LEU:HD13	2.48	0.47
1:B:168:TRP:HA	1:B:176:ILE:HG13	1.97	0.47
1:A:120:ILE:HD13	1:A:120:ILE:N	2.30	0.47
1:B:243:ASN:O	1:B:244:CYS:CB	2.61	0.47
1:A:150:VAL:HA	1:A:215:GLY:O	2.14	0.47
1:B:150:VAL:HA	1:B:215:GLY:O	2.14	0.47
1:B:64:LEU:HD12	1:B:145:ASN:HD22	1.79	0.47
1:A:297:ILE:O	1:A:301:VAL:HG23	2.15	0.47
1:A:227:PHE:HA	1:A:252:LEU:CD1	2.45	0.46
1:A:163:PHE:CD1	1:A:318:GLN:HG3	2.51	0.46
1:B:180:PRO:HD3	1:B:264:PHE:CE1	2.50	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:SER:HB3	1:A:190:ILE:HB	1.96	0.46
1:A:187:SER:O	1:A:188:SER:C	2.54	0.46
1:B:194:GLY:HA3	1:B:198:LEU:CD1	2.46	0.46
1:B:301:VAL:CG2	1:B:307:MET:HG2	2.46	0.46
1:A:110:GLU:HB3	6:A:647:HOH:O	2.16	0.46
1:B:149:MET:SD	1:B:214:ILE:HD12	2.56	0.46
1:B:297:ILE:O	1:B:301:VAL:HG23	2.16	0.46
1:A:201:PRO:CG	1:A:202:GLY:N	2.79	0.45
1:B:150:VAL:HG12	1:B:151:ASP:O	2.16	0.45
1:A:72:GLN:HA	1:A:99:VAL:HB	1.97	0.45
1:B:214:ILE:HA	1:B:217:SER:OG	2.17	0.45
1:B:125:LYS:HD3	1:B:139:PHE:CD2	2.51	0.45
1:A:272:ASN:OD1	1:A:274:GLU:HG3	2.17	0.44
1:A:125:LYS:HD3	1:A:139:PHE:CE2	2.53	0.44
1:B:222:LYS:HG3	6:B:621:HOH:O	2.18	0.43
1:A:134:ASN:O	1:A:137:GLN:HG3	2.18	0.43
1:B:171:PHE:N	1:B:172:PRO:HD3	2.34	0.43
1:B:162:VAL:HG13	6:B:616:HOH:O	2.17	0.43
1:A:149:MET:HB3	1:A:214:ILE:O	2.19	0.43
1:B:79:LEU:HD13	1:B:276:GLU:O	2.19	0.43
1:A:131:LYS:HB3	1:A:133:ARG:HG2	2.00	0.42
1:B:124:PHE:O	1:B:126:PRO:HD3	2.19	0.42
1:B:238:ILE:HD12	1:B:245:ASP:CA	2.47	0.42
1:A:182:LYS:CG	1:A:183:HIS:N	2.82	0.42
1:A:201:PRO:HG2	1:A:202:GLY:H	1.84	0.42
1:B:252:LEU:HD23	1:B:252:LEU:C	2.40	0.42
1:A:224:LEU:O	1:A:228:GLN:HG2	2.19	0.42
1:B:259:LYS:HB3	1:B:260:PRO:CD	2.50	0.42
1:A:251:PHE:O	1:A:255:ARG:HG3	2.19	0.42
1:B:120:ILE:HD13	1:B:120:ILE:N	2.33	0.42
1:A:228:GLN:HG3	6:A:617:HOH:O	2.19	0.42
1:A:133:ARG:HD2	1:A:239:ASP:OD1	2.20	0.42
1:A:155:LEU:HD12	1:A:155:LEU:C	2.40	0.41
1:A:201:PRO:CD	1:A:202:GLY:H	2.33	0.41
1:B:250:ASN:O	1:B:254:THR:HG22	2.20	0.41
1:A:73:THR:HG21	1:A:80:LEU:HD22	2.03	0.41
1:B:107:GLY:HA2	1:B:108:PRO:HD3	1.86	0.41
1:B:201:PRO:HG2	1:B:202:GLY:H	1.86	0.41
1:B:224:LEU:O	1:B:228:GLN:HG2	2.21	0.41
1:B:201:PRO:HG3	1:B:311:TYR:CE1	2.56	0.41
1:A:205:ASN:HB2	1:B:318:GLN:HE22	1.85	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ARG:NH1	1:A:239:ASP:OD2	2.53	0.40
1:B:176:ILE:HD12	1:B:263:ILE:HB	2.02	0.40
1:A:112:TRP:CZ2	1:A:122:VAL:HG21	2.55	0.40
1:A:201:PRO:CG	1:A:202:GLY:H	2.35	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/293 (90%)	243 (92%)	18 (7%)	2 (1%)	19	43
1	B	258/293 (88%)	234 (91%)	21 (8%)	3 (1%)	13	32
All	All	521/586 (89%)	477 (92%)	39 (8%)	5 (1%)	15	37

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	PRO
1	B	203	PRO
1	A	202	GLY
1	B	204	GLY
1	B	276	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/263 (88%)	218 (95%)	12 (5%)	23	49
1	B	227/263 (86%)	218 (96%)	9 (4%)	31	60
All	All	457/526 (87%)	436 (95%)	21 (5%)	27	54

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	LEU
1	A	120	ILE
1	A	143	GLU
1	A	155	LEU
1	A	182	LYS
1	A	186	THR
1	A	197	GLU
1	A	252	LEU
1	A	268	ILE
1	A	280	TYR
1	A	283	MET
1	A	288	GLU
1	B	84	LEU
1	B	120	ILE
1	B	182	LYS
1	B	186	THR
1	B	197	GLU
1	B	203	PRO
1	B	205	ASN
1	B	268	ILE
1	B	288	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	GLN
1	A	292	GLN
1	A	318	GLN
1	B	75	ASN
1	B	145	ASN
1	B	199	GLN
1	B	230	GLN
1	B	292	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GAL	C	1	2	12,12,12	1.94	2 (16%)	17,17,17	0.72	0
2	BDP	C	2	2	9,12,13	4.96	5 (55%)	12,17,19	1.53	3 (25%)
2	GAL	D	1	2	12,12,12	1.97	1 (8%)	17,17,17	0.69	0
2	BDP	D	2	2	9,12,13	4.71	4 (44%)	12,17,19	1.50	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	C	1	2	-	2/2/22/22	0/1/1/1
2	BDP	C	2	2	-	0/0/21/24	0/1/1/1
2	GAL	D	1	2	-	0/2/22/22	0/1/1/1
2	BDP	D	2	2	-	0/0/21/24	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	BDP	O2-C2	-10.20	1.21	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	BDP	O2-C2	-9.62	1.23	1.43
2	C	2	BDP	O5-C5	8.57	1.52	1.43
2	D	2	BDP	O5-C5	7.66	1.51	1.43
2	D	1	GAL	O3-C3	-6.17	1.28	1.43
2	C	1	GAL	O3-C3	-5.85	1.29	1.43
2	D	2	BDP	O5-C1	5.33	1.52	1.43
2	C	2	BDP	O5-C1	5.10	1.51	1.43
2	D	2	BDP	C4-C5	-3.35	1.46	1.53
2	C	2	BDP	C4-C5	-3.22	1.46	1.53
2	C	1	GAL	C4-C5	2.13	1.57	1.53
2	C	2	BDP	C1-C2	2.03	1.56	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	BDP	C3-C4-C5	2.62	114.40	109.02
2	D	2	BDP	C3-C4-C5	2.41	113.98	109.02
2	C	2	BDP	C2-C3-C4	-2.31	106.89	110.89
2	C	2	BDP	O5-C1-C2	-2.28	107.25	110.77
2	D	2	BDP	C2-C3-C4	-2.26	106.99	110.89

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	GAL	O5-C5-C6-O6
2	C	1	GAL	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## Oligosaccharide Chain C



## Oligosaccharide Chain D



## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	A	401	-	3,3,3	2.15	2 (66%)	2,2,2	0.46	0
4	UDP	A	351	3	20,26,26	1.86	4 (20%)	25,40,40	1.52	4 (16%)
4	UDP	B	352	3	20,26,26	1.86	4 (20%)	25,40,40	1.53	4 (16%)
5	EDO	B	402	-	3,3,3	2.25	2 (66%)	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	401	-	-	0/1/1/1	-
4	UDP	A	351	3	-	2/14/32/32	0/2/2/2
4	UDP	B	352	3	-	2/14/32/32	0/2/2/2
5	EDO	B	402	-	-	0/1/1/1	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	351	UDP	C4-N3	5.13	1.42	1.33
4	B	352	UDP	C4-N3	4.88	1.41	1.33
4	B	352	UDP	C6-N1	3.88	1.40	1.35
4	A	351	UDP	C6-N1	3.78	1.40	1.35
4	B	352	UDP	PB-O1B	3.48	1.61	1.50
4	A	351	UDP	PB-O1B	3.38	1.61	1.50
5	B	402	EDO	O2-C2	3.06	1.57	1.42
5	A	401	EDO	O2-C2	2.85	1.56	1.42
4	B	352	UDP	C2'-C1'	-2.83	1.49	1.53
5	B	402	EDO	O1-C1	2.26	1.53	1.42
5	A	401	EDO	O1-C1	2.22	1.53	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	351	UDP	C2'-C1'	-2.18	1.50	1.53

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	351	UDP	O3B-PB-O1B	3.90	125.95	110.68
4	B	352	UDP	O3B-PB-O1B	3.78	125.47	110.68
4	A	351	UDP	C5-C4-N3	-3.48	115.66	123.31
4	B	352	UDP	C5-C4-N3	-3.46	115.70	123.31
4	B	352	UDP	O5'-C5'-C4'	2.98	119.25	108.99
4	A	351	UDP	O5'-C5'-C4'	2.78	118.55	108.99
4	B	352	UDP	O3A-PB-O1B	-2.23	98.85	111.19
4	A	351	UDP	O3A-PB-O1B	-2.05	99.80	111.19

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	351	UDP	C2'-C1'-N1-C6
4	A	351	UDP	O4'-C1'-N1-C6
4	B	352	UDP	C2'-C1'-N1-C6
4	B	352	UDP	O4'-C1'-N1-C6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	265/293 (90%)	-0.52	7 (2%) 56 57	13, 25, 50, 63	0
1	B	262/293 (89%)	-0.38	6 (2%) 60 62	16, 32, 52, 69	0
All	All	527/586 (89%)	-0.45	13 (2%) 57 59	13, 28, 51, 69	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	203	PRO	4.1
1	A	279	GLY	3.4
1	B	63	ALA	3.4
1	B	203	PRO	3.1
1	B	281	SER	3.0
1	B	188	SER	2.9
1	A	188	SER	2.8
1	B	64	LEU	2.7
1	A	64	LEU	2.5
1	A	280	TYR	2.4
1	A	202	GLY	2.3
1	B	189	GLY	2.2
1	A	128	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAL	D	1	12/12	0.89	0.17	52,55,57,61	0
2	GAL	C	1	12/12	0.92	0.17	41,46,49,52	0
2	BDP	C	2	12/13	0.95	0.17	35,39,43,44	0
2	BDP	D	2	12/13	0.96	0.16	45,48,50,50	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	B	402	4/4	0.95	0.14	25,25,26,28	0
4	UDP	A	351	25/25	0.97	0.11	25,28,32,35	0
4	UDP	B	352	25/25	0.97	0.12	27,33,35,36	0
3	MN	A	600	1/1	0.98	0.04	37,37,37,37	0
5	EDO	A	401	4/4	0.98	0.10	17,17,19,20	0
3	MN	B	601	1/1	0.99	0.07	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around UDP A 351:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around UDP B 352:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.