



wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 02:10 pm BST

PDB ID : 1OPO
Title : THE STRUCTURE OF CARNATION MOTTLE VIRUS
Authors : Morgunova, E.; Dauter, Z.; Fry, E.; Stuart, D.; Stel'mashchuk, V.; Mikhailov, A.M.; Wilson, K.S.; Vainshtein, B.K.
Deposited on : 2003-03-06
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	348	
1	B	348	
1	C	348	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	500	-	-	X	-
2	SO4	B	501	-	X	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6130 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2035	1294	336	398	7			
1	B	267	Total	C	N	O	S	0	0	0
			2035	1294	336	398	7			
1	C	268	Total	C	N	O	S	0	0	0
			2044	1300	338	399	7			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: Coat protein



• Molecule 1: Coat protein



• Molecule 1: Coat protein



ARG	PRO	ARG	MET	THR	ASN	PRO	PRO	PRO	ARG	THR	SER	ARG	ASN	SER	PRO	GLY	GLN	ALA	GLY	K81	S82	M83	T84	M85	S86	K87	T88	E89	I90	L91		T97		P102	S103		P111	R112		A115	V116	F117		L120	S121	L122	L123	A124	T125		M128		R150	V151		T162		T166	K167
V168	G169		Q182		V191	D192		R196	F197	I198	R199	D200	S201		L208	V209		L216	S217	T218	Y219		T226		L231	F232	I233	Q234	Y235	T236	I237	V238	L239	S240	D241	P242	T243	K244	T245	A246	K247	I248		S252		K255	Q256	V257	D258		V264	P265		N270	E271	L272	Q273	L274	R275
V276	V277	A278	A279	G280	K281		I284		R287		V290	F291	G292	G293	F294		P297	T298	L299		I304		A313	R314	P315		C319		T323	Q324	K325		I329	L330	K331	I332	T333	K334	T335	S336	A337	E338	Q339	P340	L341	Q342	K343	V344	V345		K348								

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	382.60 Å 382.60 Å 382.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 3.20	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-3.20)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 2.1	Depositor
R, R_{free}	0.183 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6130	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2074	0.89	1/2823 (0.0%)
1	B	0.61	0/2074	0.86	1/2823 (0.0%)
1	C	0.63	0/2083	0.86	2/2834 (0.1%)
All	All	0.63	0/6231	0.87	4/8480 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	314	ARG	N-CA-C	-6.12	94.48	111.00
1	B	170	PHE	N-CA-C	-5.52	96.11	111.00
1	C	240	SER	N-CA-C	5.47	125.78	111.00
1	A	192	ASP	CB-CG-OD1	5.36	123.13	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2035	0	2053	69	0
1	B	2035	0	2053	73	1
1	C	2044	0	2066	56	1
2	A	5	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	B	1	0	0	0	0
All	All	6130	0	6172	195	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 195 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:ILE:HB	1:A:328:GLN:HG2	1.51	0.93
1:A:279:ALA:HA	1:A:323:THR:HG23	1.50	0.92
1:B:107:TRP:O	1:B:108:VAL:HB	1.74	0.85
1:A:144:SER:O	1:A:147:THR:HG23	1.80	0.81
1:A:145:PHE:HA	1:A:182:GLN:HG2	1.61	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:ASP:OD2	1:C:89:GLU:OE2[2_555]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/348 (76%)	249 (94%)	8 (3%)	8 (3%)	4	28
1	B	265/348 (76%)	244 (92%)	12 (4%)	9 (3%)	3	24
1	C	266/348 (76%)	245 (92%)	17 (6%)	4 (2%)	10	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	796/1044 (76%)	738 (93%)	37 (5%)	21 (3%)	5	31

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	SER
1	A	224	ALA
1	B	108	VAL
1	B	224	ALA
1	B	290	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/294 (78%)	201 (88%)	27 (12%)	5	23
1	B	228/294 (78%)	195 (86%)	33 (14%)	3	15
1	C	229/294 (78%)	198 (86%)	31 (14%)	4	18
All	All	685/882 (78%)	594 (87%)	91 (13%)	4	18

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	231	LEU
1	B	273	GLN
1	C	304	ILE
1	B	249	SER
1	B	264	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	128	ASN
1	B	253	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	119	GLN
1	A	342	GLN
1	B	119	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	501	-	4,4,4	0.48	0	6,6,6	2.69	4 (66%)
2	SO4	C	502	-	4,4,4	0.42	0	6,6,6	1.98	2 (33%)
2	SO4	A	500	-	4,4,4	0.23	0	6,6,6	1.35	1 (16%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	SO4	O4-S-O2	-3.95	88.70	109.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	SO4	O4-S-O3	3.57	124.31	109.06
2	C	502	SO4	O3-S-O2	-3.51	91.00	109.31
2	B	501	SO4	O4-S-O1	-2.63	95.60	109.31
2	B	501	SO4	O3-S-O1	2.62	123.00	109.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.