



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 06:01 am BST

PDB ID : 3OPF
Title : Crystal structure of TTHA0988 in space group P212121
Authors : Jacques, D.A.; Kuramitsu, S.; Yokoyama, S.; Trehella, J.; Guss, J.M.;
RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2010-09-01
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	494	<div> <div>7%</div> <div>90%</div> <div>9%</div> </div>
1	B	494	<div> <div>9%</div> <div>94%</div> <div>5%</div> </div>
1	C	494	<div> <div>6%</div> <div>91%</div> <div>8%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein TTHA0988.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	2	0
			3691	2393	663	632	3			
1	B	493	Total	C	N	O	S	0	0	0
			3680	2386	654	637	3			
1	C	488	Total	C	N	O	S	0	1	0
			3676	2381	658	634	3			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	423	Total	O	0	0
			423	423		
4	B	430	Total	O	0	0
			430	430		
4	C	306	Total	O	0	0
			306	306		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative uncharacterized protein TTHA0988




- Molecule 1: Putative uncharacterized protein TTHA0988



- Molecule 1: Putative uncharacterized protein TTHA0988





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.50Å 132.11Å 177.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.05 – 1.95 33.05 – 1.95	Depositor EDS
% Data completeness (in resolution range)	86.6 (33.05-1.95) 86.6 (33.05-1.95)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.197 , 0.233 0.210 , 0.247	Depositor DCC
R_{free} test set	5577 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.715	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12245	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3804	0.55	0/5196
1	B	0.34	0/3788	0.55	0/5179
1	C	0.33	0/3786	0.55	0/5171
All	All	0.34	0/11378	0.55	0/15546

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3691	0	3763	31	0
1	B	3680	0	3722	12	0
1	C	3676	0	3735	20	0
2	A	12	0	16	0	0
2	B	6	0	8	0	0
2	C	6	0	8	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
4	A	423	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	430	0	0	0	0
4	C	306	0	0	0	0
All	All	12245	0	11252	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LEU:HD21	1:A:208:PRO:HG2	1.44	0.96
1:B:260:LEU:HD11	1:B:381:PHE:HB3	1.57	0.86
1:A:416:ALA:HB2	1:A:426:LEU:HD23	1.74	0.70
1:A:22:LEU:HD21	1:A:192:GLY:HA3	1.78	0.65
1:A:22:LEU:HD11	1:A:191:PRO:HB2	1.77	0.64
1:B:135:LEU:HD12	1:B:136:PRO:HD2	1.80	0.63
1:A:162:LEU:CD2	1:A:208:PRO:HG2	2.26	0.62
1:B:455:VAL:HG11	1:B:480:LEU:CD1	2.30	0.60
1:C:48:LEU:CD1	1:C:64:LEU:HD11	2.31	0.60
1:A:301:LEU:HD11	1:A:324:ARG:NH2	2.21	0.56
1:C:223:LEU:HD11	1:C:367:LEU:HD12	1.88	0.55
1:A:48:LEU:CD1	1:A:64:LEU:HD11	2.37	0.55
1:B:307:ILE:HG23	1:B:311:GLN:HE21	1.72	0.55
1:B:455:VAL:HG11	1:B:480:LEU:HD13	1.88	0.53
1:A:273:LEU:C	1:A:273:LEU:HD12	2.29	0.52
1:C:28:LEU:HD11	1:C:64:LEU:HD21	1.91	0.52
1:C:182:HIS:NE2	1:C:209:ARG:HG3	2.24	0.52
1:C:342:VAL:HG11	1:C:349:ALA:HB1	1.91	0.52
1:C:455:VAL:HG11	1:C:480:LEU:CD1	2.41	0.50
1:C:423:VAL:HG12	1:C:425:LEU:CD1	2.41	0.50
1:A:28:LEU:HD11	1:A:64:LEU:HD21	1.93	0.49
1:A:223:LEU:HD11	1:A:367:LEU:HD12	1.94	0.49
1:A:5:PHE:CE1	1:A:48:LEU:HD12	2.48	0.49
1:C:290:VAL:HG11	1:C:375:VAL:HG13	1.95	0.48
1:C:416:ALA:HB2	1:C:426:LEU:HG	1.96	0.47
1:A:155:VAL:HG22	1:A:155:VAL:O	2.14	0.47
1:A:198:LEU:O	1:A:198:LEU:HD12	2.15	0.47
1:A:234:VAL:HG22	1:A:274[B]:LEU:HD23	1.97	0.47
1:C:117:LEU:HD23	1:C:245:LEU:HA	1.97	0.47
1:A:234:VAL:HG22	1:A:274[B]:LEU:CD2	2.45	0.47
1:A:48:LEU:HD11	1:A:64:LEU:HD11	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LEU:HD21	1:A:208:PRO:CG	2.32	0.46
1:A:426:LEU:HD11	4:A:696:HOH:O	2.15	0.46
1:A:27:ALA:HB1	1:A:67:LEU:HD22	1.97	0.45
1:C:423:VAL:HG12	1:C:425:LEU:HD13	1.99	0.45
1:C:127:ALA:HB1	1:C:128:GLU:HA	1.99	0.45
1:C:48:LEU:HD12	1:C:64:LEU:HD11	1.98	0.45
1:C:282:VAL:HG13	1:C:322:ARG:HG2	1.98	0.45
1:C:120:THR:HG22	1:C:354:ARG:HD3	1.99	0.44
1:B:425:LEU:HD11	1:B:489:ILE:HD12	1.99	0.43
1:B:423:VAL:HG21	1:B:480:LEU:HD22	2.00	0.43
1:A:300:ALA:HB3	1:A:307:ILE:HB	1.99	0.43
1:A:301:LEU:HD11	1:A:324:ARG:CZ	2.47	0.43
1:A:273:LEU:CD1	1:A:273:LEU:C	2.87	0.42
1:B:342:VAL:CG2	1:B:349:ALA:HB1	2.49	0.42
1:B:228:PRO:HD3	1:B:363:ALA:HB2	2.00	0.42
1:A:345:PHE:CE1	1:A:346:LEU:HD13	2.54	0.42
1:C:300:ALA:HB3	1:C:307:ILE:HB	2.01	0.42
1:A:35:GLY:O	1:A:37:LEU:HD13	2.19	0.42
1:A:89:LEU:HB3	1:A:90:PRO:HD3	2.02	0.42
1:A:255:ALA:HB3	1:A:256:PRO:HD3	2.02	0.42
1:A:187:PHE:CE2	1:A:245:LEU:HD13	2.54	0.42
1:C:232:ASP:HB3	1:C:274:LEU:HD11	2.02	0.42
1:C:135:LEU:HD23	1:C:136:PRO:O	2.20	0.42
1:A:231:MET:HE1	1:A:420:ARG:HD3	2.01	0.41
1:B:2:VAL:CG1	1:B:49:GLU:HG2	2.50	0.41
1:A:404:ALA:HB2	4:A:1038:HOH:O	2.21	0.41
1:C:43:TYR:OH	1:C:156:GLN:OE1	2.18	0.41
1:B:425:LEU:HD11	1:B:489:ILE:CD1	2.50	0.40
1:B:223:LEU:HD11	1:B:367:LEU:HD12	2.04	0.40
1:C:477:LEU:HD23	1:C:477:LEU:HA	1.98	0.40
1:A:398:PRO:HG2	1:A:468:LYS:HE2	2.03	0.40
1:A:55:LEU:HD21	1:A:60:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/494 (99%)	478 (98%)	10 (2%)	0	100	100
1	B	491/494 (99%)	480 (98%)	11 (2%)	0	100	100
1	C	485/494 (98%)	475 (98%)	10 (2%)	0	100	100
All	All	1464/1482 (99%)	1433 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/381 (97%)	364 (99%)	5 (1%)	67	62
1	B	365/381 (96%)	355 (97%)	10 (3%)	44	34
1	C	368/381 (97%)	362 (98%)	6 (2%)	62	58
All	All	1102/1143 (96%)	1081 (98%)	21 (2%)	57	50

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	LEU
1	A	123	PHE
1	A	273	LEU
1	A	405	PHE
1	A	425	LEU
1	B	37	LEU
1	B	38	ASP
1	B	48	LEU
1	B	100	LEU
1	B	123	PHE
1	B	273	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	342	VAL
1	B	346	LEU
1	B	396	PRO
1	B	425	LEU
1	C	37	LEU
1	C	38	ASP
1	C	100	LEU
1	C	224	LEU
1	C	273	LEU
1	C	425	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	311	GLN
1	C	169	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	495	-	5,5,5	0.41	0	5,5,5	0.23	0
2	GOL	B	495	-	5,5,5	0.36	0	5,5,5	0.14	0
3	SO4	C	495	-	4,4,4	0.14	0	6,6,6	0.11	0
3	SO4	A	496	-	4,4,4	0.12	0	6,6,6	0.11	0
2	GOL	A	497	-	5,5,5	0.37	0	5,5,5	0.32	0
3	SO4	B	496	-	4,4,4	0.15	0	6,6,6	0.12	0
2	GOL	C	496	-	5,5,5	0.38	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	495	-	-	2/4/4/4	-
2	GOL	A	497	-	-	2/4/4/4	-
2	GOL	C	496	-	-	2/4/4/4	-
2	GOL	B	495	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	495	GOL	C1-C2-C3-O3
2	B	495	GOL	C1-C2-C3-O3
2	A	497	GOL	C1-C2-C3-O3
2	A	497	GOL	O2-C2-C3-O3
2	C	496	GOL	C1-C2-C3-O3
2	B	495	GOL	O2-C2-C3-O3
2	A	495	GOL	O2-C2-C3-O3
2	C	496	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/494 (99%)	0.74	34 (6%) 16 25	24, 34, 55, 82	0
1	B	493/494 (99%)	0.81	44 (8%) 9 15	25, 37, 56, 70	0
1	C	488/494 (98%)	0.89	32 (6%) 18 26	28, 40, 58, 77	0
All	All	1471/1482 (99%)	0.81	110 (7%) 14 22	24, 37, 57, 82	0

All (110) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	72	ALA	10.4
1	C	304	GLY	6.7
1	C	70	GLU	5.8
1	C	67	LEU	5.7
1	A	68	PRO	5.6
1	A	63	LEU	5.1
1	B	75	GLY	4.6
1	C	450	SER	4.6
1	B	63	LEU	4.6
1	A	304	GLY	4.5
1	B	314	LEU	4.4
1	A	67	LEU	4.3
1	C	362	TRP	4.1
1	A	34	PRO	4.1
1	A	69	GLN	4.1
1	C	63	LEU	4.0
1	A	426	LEU	3.8
1	B	362	TRP	3.8
1	B	56	SER	3.5
1	A	53	ARG	3.5
1	B	303	GLU	3.4
1	B	450	SER	3.4
1	B	253	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	3.2
1	C	135	LEU	3.2
1	C	372	LEU	3.2
1	C	391	ARG	3.2
1	B	73	GLU	3.2
1	B	68	PRO	3.1
1	A	1	MET	3.1
1	C	69	GLN	3.1
1	A	71	ARG	3.0
1	B	230	LEU	3.0
1	B	55	LEU	3.0
1	B	69	GLN	3.0
1	B	299	VAL	3.0
1	B	273	LEU	3.0
1	C	68	PRO	3.0
1	C	66	GLY	2.9
1	C	302	LEU	2.9
1	A	52	PRO	2.9
1	B	374	PRO	2.8
1	A	72	ALA	2.8
1	A	429	GLU	2.8
1	C	138	LYS	2.8
1	B	412	PRO	2.8
1	B	414	ARG	2.7
1	A	303	GLU	2.7
1	B	34	PRO	2.7
1	A	64	LEU	2.6
1	A	59	ARG	2.6
1	B	304	GLY	2.6
1	B	67	LEU	2.6
1	C	217	GLU	2.6
1	A	450	SER	2.6
1	B	66	GLY	2.6
1	C	335	ALA	2.5
1	A	462	SER	2.5
1	A	2	VAL	2.5
1	B	411	GLY	2.5
1	B	386	LEU	2.5
1	A	54	ARG	2.5
1	B	333	TYR	2.4
1	A	449	PRO	2.4
1	A	463	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	100	LEU	2.4
1	A	35	GLY	2.4
1	B	490	HIS	2.4
1	C	91	GLU	2.4
1	A	50	TYR	2.4
1	A	66	GLY	2.4
1	A	233	LEU	2.3
1	B	59	ARG	2.3
1	B	458	ALA	2.3
1	B	432	GLY	2.3
1	C	271	ALA	2.3
1	B	276	PHE	2.3
1	B	117	LEU	2.3
1	B	426	LEU	2.3
1	A	51	ASP	2.2
1	B	76	ARG	2.2
1	C	479	LEU	2.2
1	C	43	TYR	2.2
1	B	421	VAL	2.2
1	A	274[A]	LEU	2.2
1	C	96	LEU	2.2
1	C	333	TYR	2.2
1	C	303	GLU	2.2
1	B	252	PRO	2.2
1	B	118	GLY	2.2
1	B	54	ARG	2.2
1	B	389	ALA	2.2
1	C	277	ALA	2.2
1	B	227	GLU	2.2
1	A	440	THR	2.1
1	C	269	ALA	2.1
1	B	40	VAL	2.1
1	B	48	LEU	2.1
1	A	76	ARG	2.1
1	A	277	ALA	2.1
1	A	330	VAL	2.1
1	C	404	ALA	2.1
1	A	33	PRO	2.1
1	B	80	ILE	2.1
1	C	125	PHE	2.0
1	C	261	ALA	2.0
1	C	94	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2	VAL	2.0
1	A	457	LEU	2.0
1	C	273	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	C	496	6/6	0.79	0.21	56,61,62,63	0
2	GOL	A	495	6/6	0.80	0.27	39,46,47,49	0
2	GOL	B	495	6/6	0.81	0.21	45,53,53,60	0
3	SO4	B	496	5/5	0.90	0.25	61,64,64,67	0
2	GOL	A	497	6/6	0.90	0.18	41,48,50,50	0
3	SO4	C	495	5/5	0.91	0.32	66,69,70,71	0
3	SO4	A	496	5/5	0.96	0.17	53,53,55,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.