



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 03:30 am BST

PDB ID : 3OPU
Title : Crystal structure of the C-terminal domain of Streptococcus mutans surface protein SpaP
Authors : Nylander, A.; Persson, K.
Deposited on : 2010-09-02
Resolution : 2.18 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>.</div> </div> </div>
1	B	339	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>15%</div> <div>.</div> </div> </div>
1	C	339	<div> <div>4%</div> <div> <div></div> <div>73%</div> <div>25%</div> <div>..</div> </div> </div>
1	D	339	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>18%</div> <div>.</div> </div> </div>
1	E	339	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>14%</div> <div>..</div> </div> </div>
1	F	339	<div> <div>%</div> <div> <div></div> <div>81%</div> <div>17%</div> <div>.</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SpaP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	0	0
			2670	1691	434	542	3			
1	B	338	Total	C	N	O	S	0	0	0
			2670	1691	434	542	3			
1	C	337	Total	C	N	O	S	0	0	0
			2664	1687	434	540	3			
1	D	339	Total	C	N	O	S	0	0	0
			2678	1695	436	544	3			
1	E	339	Total	C	N	O	S	0	0	0
			2678	1695	436	544	3			
1	F	338	Total	C	N	O	S	0	0	0
			2670	1691	434	542	3			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	2	Total	Ca	0	0
			2	2		
2	E	2	Total	Ca	0	0
			2	2		
2	B	2	Total	Ca	0	0
			2	2		
2	C	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		
2	F	2	Total	Ca	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	96	Total 96	O 96	0	0
3	B	109	Total 109	O 109	0	0
3	C	65	Total 65	O 65	0	0
3	D	112	Total 112	O 112	0	0
3	E	76	Total 76	O 76	0	0
3	F	87	Total 87	O 87	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SpaP




• Molecule 1: SpaP




• Molecule 1: SpaP




• Molecule 1: SpaP



• Molecule 1: SpaP



• Molecule 1: SpaP



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	135.78 Å 238.39 Å 78.49 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.01 – 2.18 29.11 – 2.18	Depositor EDS
% Data completeness (in resolution range)	98.2 (29.01-2.18) 99.7 (29.11-2.18)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.19 (at 2.18 Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.210 , 0.248 0.204 , 0.236	Depositor DCC
R_{free} test set	6762 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16587	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2588e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.66	0/2720	0.75	6/3695 (0.2%)
1	B	0.72	3/2720 (0.1%)	0.68	3/3695 (0.1%)
1	C	0.61	0/2714	0.65	0/3686
1	D	0.66	1/2728 (0.0%)	0.66	1/3706 (0.0%)
1	E	0.71	2/2728 (0.1%)	0.70	2/3706 (0.1%)
1	F	0.70	3/2720 (0.1%)	0.64	0/3695
All	All	0.68	9/16330 (0.1%)	0.68	12/22183 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	2
1	F	0	1
All	All	0	3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	1321	ASN	CB-CG	-6.81	1.35	1.51
1	E	1152	TYR	CD2-CE2	-6.40	1.29	1.39
1	B	1334	LYS	CE-NZ	-6.03	1.33	1.49
1	D	1469	ASN	C-N	-5.90	1.20	1.34
1	F	1468	GLU	C-N	-5.49	1.21	1.34

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1373	SER	O-C-N	-9.04	108.23	122.70
1	A	1374	GLU	O-C-N	8.95	137.02	122.70
1	B	1345	THR	N-CA-C	8.56	134.11	111.00
1	A	1374	GLU	CA-C-N	-7.13	101.52	117.20
1	B	1334	LYS	CD-CE-NZ	6.91	127.58	111.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	1194	SER	Peptide
1	E	1344	ASP	Peptide
1	F	1320	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2670	0	2596	37	0
1	B	2670	0	2596	38	0
1	C	2664	0	2590	62	0
1	D	2678	0	2602	49	0
1	E	2678	0	2603	60	0
1	F	2670	0	2594	56	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	96	0	0	2	0
3	B	109	0	0	2	0
3	C	65	0	0	6	0
3	D	112	0	0	6	0
3	E	76	0	0	2	0
3	F	87	0	0	3	0
All	All	16587	0	15581	297	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 297 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:1194:SER:CA	1:E:1195:SER:HB2	1.55	1.36
1:F:1228:GLY:HA2	1:F:1229:ASN:ND2	1.56	1.20
1:E:1194:SER:HA	1:E:1195:SER:HB2	1.23	1.13
1:E:1194:SER:CB	1:E:1195:SER:HB2	1.82	1.10
1:E:1171:THR:HG22	1:E:1326:ASN:HD21	1.12	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/339 (99%)	325 (97%)	10 (3%)	1 (0%)	41	43
1	B	336/339 (99%)	322 (96%)	12 (4%)	2 (1%)	25	24
1	C	335/339 (99%)	321 (96%)	12 (4%)	2 (1%)	25	24
1	D	337/339 (99%)	326 (97%)	9 (3%)	2 (1%)	25	24
1	E	337/339 (99%)	318 (94%)	13 (4%)	6 (2%)	8	5
1	F	336/339 (99%)	322 (96%)	12 (4%)	2 (1%)	25	24
All	All	2017/2034 (99%)	1934 (96%)	68 (3%)	15 (1%)	22	20

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1227	ASN
1	B	1345	THR
1	E	1195	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	1229	ASN
1	E	1345	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/296 (100%)	285 (97%)	10 (3%)	37	44
1	B	295/296 (100%)	287 (97%)	8 (3%)	44	54
1	C	294/296 (99%)	276 (94%)	18 (6%)	18	19
1	D	296/296 (100%)	287 (97%)	9 (3%)	41	49
1	E	296/296 (100%)	284 (96%)	12 (4%)	30	36
1	F	295/296 (100%)	287 (97%)	8 (3%)	44	54
All	All	1771/1776 (100%)	1706 (96%)	65 (4%)	34	40

5 of 65 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1375	GLU
1	D	1151	ASN
1	F	1255	LYS
1	C	1414	SER
1	C	1436	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1356	ASN
1	D	1151	ASN
1	E	1326	ASN
1	C	1265	GLN
1	E	1356	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/339 (99%)	-0.02	6 (1%) 68 69	12, 21, 39, 53	0
1	B	338/339 (99%)	-0.08	8 (2%) 59 59	12, 22, 37, 61	0
1	C	337/339 (99%)	0.22	13 (3%) 39 40	15, 27, 58, 70	0
1	D	339/339 (100%)	-0.02	9 (2%) 54 55	11, 22, 44, 52	0
1	E	339/339 (100%)	0.15	18 (5%) 26 27	13, 24, 49, 80	0
1	F	338/339 (99%)	-0.06	4 (1%) 79 79	14, 24, 43, 54	0
All	All	2029/2034 (99%)	0.03	58 (2%) 51 52	11, 23, 47, 80	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1152	TYR	5.4
1	F	1152	TYR	4.7
1	E	1227	ASN	4.2
1	A	1152	TYR	4.2
1	E	1345	THR	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	C	1	1/1	0.86	0.08	52,52,52,52	1
2	CA	D	1	1/1	0.88	0.14	46,46,46,46	1
2	CA	F	1	1/1	0.93	0.12	38,38,38,38	1
2	CA	E	2	1/1	0.93	0.12	41,41,41,41	1
2	CA	A	9	1/1	0.95	0.14	33,33,33,33	1
2	CA	B	1	1/1	0.95	0.10	29,29,29,29	1
2	CA	E	8	1/1	0.97	0.08	25,25,25,25	1
2	CA	C	6	1/1	0.97	0.04	28,28,28,28	1
2	CA	F	3	1/1	0.98	0.09	28,28,28,28	1
2	CA	A	4	1/1	0.98	0.11	27,27,27,27	1
2	CA	D	7	1/1	0.99	0.12	25,25,25,25	1
2	CA	B	5	1/1	0.99	0.13	26,26,26,26	1

6.5 Other polymers [i](#)

There are no such residues in this entry.