



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 02:58 pm BST

PDB ID : 5ORG
Title : Structure of the periplasmic binding protein (PBP) OccJ from *A. tumefaciens* B6 in complex with octopine.
Authors : Vigouroux, A.; Morera, S.
Deposited on : 2017-08-16
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div>8%</div> <div>94%</div> <div>6%</div> <div>• •</div> </div>
1	B	263	<div> <div>7%</div> <div>90%</div> <div>6%</div> <div>•</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Octopine-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			1904	1210	309	375	10			
1	B	253	Total	C	N	O	S	0	1	0
			1874	1195	303	367	9			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	initiating methionine	UNP P0A4F8
A	277	HIS	-	expression tag	UNP P0A4F8
A	278	HIS	-	expression tag	UNP P0A4F8
A	279	HIS	-	expression tag	UNP P0A4F8
A	280	HIS	-	expression tag	UNP P0A4F8
A	281	HIS	-	expression tag	UNP P0A4F8
A	282	HIS	-	expression tag	UNP P0A4F8
B	20	MET	-	initiating methionine	UNP P0A4F8
B	277	HIS	-	expression tag	UNP P0A4F8
B	278	HIS	-	expression tag	UNP P0A4F8
B	279	HIS	-	expression tag	UNP P0A4F8
B	280	HIS	-	expression tag	UNP P0A4F8
B	281	HIS	-	expression tag	UNP P0A4F8
B	282	HIS	-	expression tag	UNP P0A4F8

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Na	0	0
			2	2		
3	A	7	Total	Na	0	0
			7	7		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	4	Total	Cl	0	0
			4	4		
4	A	10	Total	Cl	0	0
			10	10		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is octopine (three-letter code: 6DB) (formula: $C_9H_{18}N_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			17	9	4	4		
6	B	1	Total	C	N	O	0	0
			17	9	4	4		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total 4	C 2	O 2	0	0
7	B	1	Total 4	C 2	O 2	0	0
7	B	1	Total 4	C 2	O 2	0	0
7	B	1	Total 4	C 2	O 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	138	Total 138	O 138	0	0
8	B	105	Total 105	O 105	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Octopine-binding periplasmic protein



- Molecule 1: Octopine-binding periplasmic protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	99.49 Å 99.49 Å 157.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.14 – 1.99 46.45 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.14-1.99) 99.6 (46.45-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.98 Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.196 , 0.221 0.201 , 0.224	Depositor DCC
R_{free} test set	2736 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4142	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, NA, EDO, 6DB, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1936	0.66	1/2613 (0.0%)
1	B	0.48	0/1909	0.63	0/2578
All	All	0.50	0/3845	0.64	1/5191 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	PRO	C-N-CA	5.19	133.19	122.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1904	0	1915	3	0
1	B	1874	0	1895	8	0
2	A	8	0	12	0	0
2	B	20	0	30	3	0
3	A	7	0	0	0	0
3	B	2	0	0	0	0
4	A	10	0	0	0	0
4	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	6	0	8	0	0
5	B	6	0	8	2	0
6	A	17	0	0	0	0
6	B	17	0	0	0	0
7	A	12	0	9	1	0
7	B	12	0	9	0	0
8	A	138	0	0	3	0
8	B	105	0	0	0	0
All	All	4142	0	3886	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:492:HOH:O	2:B:304:EDO:H22	1.94	0.65
1:A:122:ALA:HA	1:A:221:VAL:HG11	1.86	0.58
1:B:214:GLU:HG3	2:B:304:EDO:H11	1.85	0.58
1:B:147:ILE:HD11	1:B:223:PRO:CG	2.40	0.51
8:A:496:HOH:O	2:B:304:EDO:H21	2.10	0.51
1:B:130:THR:HG22	1:B:224:LEU:HB2	1.93	0.50
1:B:243:ALA:H	5:B:312:GOL:H31	1.79	0.47
7:A:322:ACT:H2	8:A:479:HOH:O	2.17	0.45
1:B:122:ALA:HA	1:B:221:VAL:HG11	1.99	0.44
1:A:68:ALA:HB1	1:B:213:PRO:HB3	2.00	0.44
1:B:241:ASP:HA	5:B:312:GOL:H2	1.98	0.44
1:B:181:TYR:CD1	1:B:187:HIS:HB3	2.54	0.43
1:A:181:TYR:CD1	1:A:187:HIS:HB3	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/263 (97%)	251 (98%)	3 (1%)	1 (0%)	34	30
1	B	252/263 (96%)	248 (98%)	4 (2%)	0	100	100
All	All	507/526 (96%)	499 (98%)	7 (1%)	1 (0%)	47	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	275	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/204 (97%)	194 (98%)	4 (2%)	55	58
1	B	195/204 (96%)	191 (98%)	4 (2%)	53	57
All	All	393/408 (96%)	385 (98%)	8 (2%)	55	58

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	ASP
1	A	142	LYS
1	A	179	LYS
1	A	257	GLU
1	B	72	ASP
1	B	176	VAL
1	B	179	LYS
1	B	257	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	ASN
1	B	111	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 40 ligands modelled in this entry, 23 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	B	305	-	3,3,3	0.64	0	2,2,2	0.29	0
2	EDO	B	304	-	3,3,3	0.50	0	2,2,2	0.07	0
2	EDO	B	301	-	3,3,3	0.62	0	2,2,2	0.23	0
7	ACT	B	316	-	1,3,3	5.27	1 (100%)	0,3,3	0.00	-
7	ACT	A	324	-	1,3,3	5.41	1 (100%)	0,3,3	0.00	-
5	GOL	B	312	-	5,5,5	0.04	0	5,5,5	0.32	0
7	ACT	A	322	-	1,3,3	2.44	1 (100%)	0,3,3	0.00	-
5	GOL	A	320	-	5,5,5	0.04	0	5,5,5	0.13	0
2	EDO	B	303	-	3,3,3	0.63	0	2,2,2	0.23	0
7	ACT	A	323	3	1,3,3	3.61	1 (100%)	0,3,3	0.00	-
6	6DB	A	321	-	10,16,16	1.16	1 (10%)	10,20,20	1.37	1 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ACT	B	314	-	1,3,3	4.42	1 (100%)	0,3,3	0.00	-
7	ACT	B	315	-	1,3,3	4.42	1 (100%)	0,3,3	0.00	-
2	EDO	A	302	-	3,3,3	0.54	0	2,2,2	0.36	0
2	EDO	B	302	-	3,3,3	0.57	0	2,2,2	0.33	0
6	6DB	B	313	-	10,16,16	1.29	1 (10%)	10,20,20	1.11	1 (10%)
2	EDO	A	301	-	3,3,3	0.62	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	305	-	-	0/1/1/1	-
2	EDO	B	304	-	-	1/1/1/1	-
2	EDO	B	301	-	-	0/1/1/1	-
5	GOL	B	312	-	-	0/4/4/4	-
5	GOL	A	320	-	-	0/4/4/4	-
2	EDO	B	303	-	-	0/1/1/1	-
2	EDO	A	301	-	-	0/1/1/1	-
2	EDO	A	302	-	-	0/1/1/1	-
2	EDO	B	302	-	-	0/1/1/1	-
6	6DB	B	313	-	-	1/11/19/19	-
6	6DB	A	321	-	-	1/11/19/19	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	324	ACT	CH3-C	5.41	1.55	1.48
7	B	316	ACT	CH3-C	5.27	1.55	1.48
7	B	314	ACT	CH3-C	4.42	1.54	1.48
7	B	315	ACT	CH3-C	4.42	1.54	1.48
7	A	323	ACT	CH3-C	3.61	1.53	1.48
6	B	313	6DB	CAP-NAN	2.94	1.39	1.33
6	A	321	6DB	CAP-NAN	2.66	1.38	1.33
7	A	322	ACT	CH3-C	2.44	1.51	1.48

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	321	6DB	CAL-CAM-NAN	-3.29	102.79	112.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	313	6DB	CAL-CAM-NAN	-2.58	104.84	112.21

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	313	6DB	CAE-CAI-N-CA
6	A	321	6DB	CAE-CAI-N-CA
2	B	304	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	304	EDO	3	0
5	B	312	GOL	2	0
7	A	322	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/263 (97%)	0.45	22 (8%) 10 9	25, 40, 70, 106	0
1	B	253/263 (96%)	0.45	18 (7%) 16 15	27, 45, 71, 93	0
All	All	510/526 (96%)	0.45	40 (7%) 13 12	25, 42, 71, 106	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	20	MET	6.0
1	A	276	GLN	5.7
1	B	244	LEU	4.8
1	B	23	LYS	4.2
1	B	63	LYS	3.6
1	A	142	LYS	3.6
1	A	273	VAL	3.3
1	B	58	GLU	3.3
1	A	139	ALA	3.3
1	A	275	PRO	3.3
1	B	61	LYS	3.1
1	A	174	GLY	3.0
1	B	175	SER	2.8
1	B	62	ALA	2.7
1	A	131	TYR	2.7
1	A	256	SER	2.7
1	A	258	ASP	2.6
1	A	257	GLU	2.6
1	B	242	THR	2.5
1	A	262	LYS	2.5
1	A	61	LYS	2.5
1	B	57	CYS	2.4
1	A	146	ASP	2.4
1	B	173	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	135	SER	2.3
1	A	259	GLY	2.3
1	B	174	GLY	2.3
1	A	145	ALA	2.3
1	B	243	ALA	2.3
1	B	24	SER	2.3
1	B	258	ASP	2.2
1	A	274	THR	2.2
1	A	21	GLN	2.2
1	B	246	ALA	2.1
1	A	64	CYS	2.1
1	B	25	ILE	2.1
1	A	175	SER	2.1
1	A	138	ASP	2.0
1	B	210	ILE	2.0
1	B	257	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	A	318	1/1	0.61	0.10	108,108,108,108	0
5	GOL	B	312	6/6	0.67	0.29	86,87,88,89	0
4	CL	A	315	1/1	0.71	0.12	84,84,84,84	0
2	EDO	B	301	4/4	0.75	0.36	76,76,76,77	0
2	EDO	B	305	4/4	0.75	0.21	72,74,74,74	0
4	CL	A	317	1/1	0.76	0.09	99,99,99,99	0
3	NA	A	309	1/1	0.80	0.17	84,84,84,84	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	B	303	4/4	0.81	0.22	81,82,83,83	0
4	CL	A	316	1/1	0.81	0.24	103,103,103,103	0
2	EDO	A	301	4/4	0.82	0.14	69,70,72,73	0
3	NA	B	306	1/1	0.83	0.18	46,46,46,46	0
4	CL	B	311	1/1	0.84	0.23	82,82,82,82	0
7	ACT	A	324	4/4	0.85	0.21	58,59,59,62	0
7	ACT	B	316	4/4	0.85	0.16	58,59,60,62	0
4	CL	A	319	1/1	0.86	0.38	60,60,60,60	0
2	EDO	A	302	4/4	0.86	0.52	87,87,88,88	0
3	NA	A	308	1/1	0.87	0.12	77,77,77,77	0
4	CL	A	311	1/1	0.87	0.07	72,72,72,72	0
3	NA	A	306	1/1	0.87	0.14	50,50,50,50	0
4	CL	B	309	1/1	0.88	0.16	87,87,87,87	0
4	CL	B	308	1/1	0.89	0.07	79,79,79,79	0
2	EDO	B	304	4/4	0.90	0.35	50,51,52,52	0
4	CL	A	314	1/1	0.91	0.16	94,94,94,94	0
4	CL	B	310	1/1	0.91	0.17	71,71,71,71	0
3	NA	A	304	1/1	0.92	0.10	43,43,43,43	0
3	NA	A	305	1/1	0.92	0.13	43,43,43,43	0
5	GOL	A	320	6/6	0.92	0.22	79,79,79,80	0
4	CL	A	310	1/1	0.92	0.07	63,63,63,63	0
2	EDO	B	302	4/4	0.93	0.42	82,82,82,83	0
4	CL	A	313	1/1	0.93	0.05	80,80,80,80	0
7	ACT	B	315	4/4	0.94	0.13	51,54,55,58	0
7	ACT	A	322	4/4	0.94	0.17	48,51,52,54	0
7	ACT	A	323	4/4	0.95	0.25	54,54,55,61	0
6	6DB	A	321	17/17	0.96	0.10	23,27,35,38	0
7	ACT	B	314	4/4	0.97	0.30	58,59,60,61	0
3	NA	B	307	1/1	0.97	0.15	41,41,41,41	0
4	CL	A	312	1/1	0.97	0.26	68,68,68,68	0
6	6DB	B	313	17/17	0.97	0.10	29,33,38,38	0
3	NA	A	307	1/1	0.98	0.15	43,43,43,43	0
3	NA	A	303	1/1	0.98	0.13	37,37,37,37	0

6.5 Other polymers ⓘ

There are no such residues in this entry.